

Research Article

Factors Influencing Acute Stroke Thrombolytic Treatments in Hispanics in the San Diego Region

Chen PM^{1*}, Nguyen DT¹, Ho JP¹, Pirastehfar M¹, Narula R¹, Rapp K¹, Agrawal K¹, Huisa B¹, Modir R¹, Meyer D¹, Hemmen T¹, Kidwell C² and Meyer BC¹

¹Department of Neurosciences, Stroke Center, University of California, USA

²Department of Neurology, University of Arizona, USA

*Corresponding author: Chen PM, Department of Neurosciences, University of California, San Diego, USA

Received: November 28, 2017; Accepted: January 04, 2018; Published: January 11, 2018

Abstract

Background: Since the introduction of recombinant tissue plasminogen activator (rt-PA) for acute ischemic stroke, rt-PA rate and number of stroke centers have increased. Despite this, studies have shown racial and ethnic disparities in stroke care especially in Black and Hispanic populations. What factors are related to the administration of rt-PA within the Hispanic population has to date been unclear.

Methods: We performed a retrospective review of IRB approved, prospectively collected data from the UC San Diego Stroke Registry from 7/2004-7/2016. Patients were included based on the primary diagnosis of Transient Ischemic Attack or Ischemic Stroke. Hispanic vs. non-Hispanic patients were compared to assess for overall rt-PA treatment rates and process of care intervals. For the Hispanic cohort itself, demographics and NIHSS scores were assessed to determine why some Hispanics received rt-PA while others were not.

Results: Overall, 1489 patients (300 hispanic vs. 1189 non-hispanic) were included. Comparing hispanics to non-hispanics, there was no difference in rt-pa rate (35.3% vs. 33.1%; p=0.49). In rt-pa treated patients, "onset to arrival" interval was higher in hispanics (1.03 vs. 0.88 hours; p=0.04), while the "arrival to treatment" interval was not different (1.13 vs. 1.02 hours; p=0.07). When looking at hispanic patients only, there was no difference in baseline characteristics except for initial nihss in treated vs. Non-treated patients (13.27 vs. 7.24; p<0.001).

Conclusion: Our analyses sought to determine the factors important to administration of rt-PA to Hispanic patients. These findings highlight the need for strategies to improve recognition and presentation pathways for Hispanics.

Keywords: Stroke; TPA; Hispanic; Regional stroke differences; Ethnic disparities

Introduction

Since the introduction of recombinant tissue plasminogen activator (rt-PA) for acute ischemic stroke (AIS), we have seen the rate of rt-PA administrations and number of new primary and comprehensive stroke centers increase [1]. The recent positive endovascular trials have also further added to treatment options providers can provide in acute stroke care. Despite the rise of acute stroke treatment utilization, we continue to see racial and ethnic variations in stroke care in both urban and rural settings, especially in Hispanic populations. Among minority groups, Hispanic Americans are the fastest-growing group in the United States and are estimated to represent 15% of the US population [2]. Hispanics also have an increased prevalence of risks factors for strokes when compared to non-Hispanic whites [3].

Whether ethnicity and race is a major barrier for acute stroke care remains controversial. Some reports have shown that ethnic disparities trump other risk factors in determining the delay of rt-PA use in acute ischemic stroke [4]. Several studies have demonstrated Blacks presenting with AIS were significantly less likely to be treated with intravenous rt-PA than whites [5-7]. How Hispanic ethnicity

affects rt-PA evaluations has had mixed results. Two large nationwide population studies found lower rates of rt-PA use in Hispanics while a separate study showed similar rates of rt-PA treatment in Hispanics vs. non-Hispanic whites [8-10].

Among rt-PA treated AIS Hispanic and non-Hispanic patients, differences in demographics, comorbidities, and interval of care processes in this cohort have not been clearly studied. In our patient population, we sought to present rt-PA rates for both Hispanics and non-Hispanics, but more so to report on factors that influence the administration of rt-PA to both Hispanics and non-Hispanics in our San Diego, California region.

Methods

Design

This IRB approved study is a retrospective review prospectively collected data from the UC San Diego Stroke Registry. We included all consecutive acute ischemic strokes patients presenting within 12 hours of symptom onset to our network hospitals from June 2004 to July 2016. 1 of the 3 hospitals is a Joint Commission certified Comprehensive Stroke center and the other 2 are certified Primary

Table 1: rt-PA by Ethnicity.

	Ethnicity			p-value
	Hispanic (n=300)	Non-Hispanic (n=1189)	Overall (n=1489)	
rt-PA(+)	106 (35.3%)	394 (33.1%)	593 (33.6%)	0.494
rt-PA(-)	194 (64.7%)	795 (66.9%)	1173 (66.4%)	

Table 2: Process of care intervals in Hispanics vs non-Hispanic whites treated with rt-PA. Data is reported as the mean hours (standard deviation).

	rt-PA(+)			p-value
	Hispanic (n = 102)	Non-Hispanic (n = 374)		
Onset to Arrival	1.03 (0.08, 3.55)	0.88 (0, 4.67)		0.042
Arrival to Treatment	1.13 (0.47, 2.92)	1.02 (0.02, 4.38)		0.071
Onset to Treatment	2.42 (1.02, 4.38)	2.13 (0.15, 4.67)		0.004

Stroke Centers.

Patient selection

Patients age ≥18 years were included based on the presentation and primary diagnosis of Transient Ischemic Attack (TIA) or Acute Ischemic Stroke (AIS). Transfer patients or in hospital stroke events were excluded from analysis for consistent reporting. Patient specific variables included age, gender, self-reported race/ethnicity, risk factors, blood pressure, National Institutes of Health Stroke Scale (NIHSS) score, and process of care intervals. Race/ethnicity is categorized as American Indian, Asian, Black, Hawaiian/Pacific Islander, Hispanic white, and non-Hispanic white.

Statistical analysis

First, in order to assess rt-PA rates, Hispanic vs. non-Hispanic groups were compared. Second, care intervals were also assessed for the rt-PA subset comparisons and included onset-to-arrival time, arrival-to-decision-time, arrival-to-treatment time, and onset-to-treatment time. Times were reported as means and standard deviations (hours). Third, in order to assess the Hispanic population itself for rt-PA treatment differences, rt-PA+ vs. rt-PA- groups were compared within the Hispanic group itself. Baseline characteristics including risk factors, blood pressure, and presenting NIHSS were compared for Hispanics who received rt-PA (rt-PA+) to Hispanics who didn't receive rt-PA (rt-PA-). Significance was determined by a two sample t-test for continuous variables and Fisher's exact test for categorical variables.

Results

We assessed 1,489 patients with a primary diagnosis of TIA or AIS. Among these patients, 300 were Hispanics (20% of the cohort) and 1189 were non-Hispanic. For the overall analysis, 33.6% of the assessed patients received rt-PA treatment. Among these patients, there was no difference in the rate of rt-PA utilization among Hispanic and non-Hispanic whites (35.3% vs. 33.1%; p=0.49) (Table 1).

In rt-PA treated patient subset, Hispanics and non-Hispanic whites had no significant difference in presenting NIHSS (mean 13.27 vs. 11.84; p=0.151). However, Hispanics had significant longer “onset to arrival” intervals (1.08 vs. 0.90 hours; p=0.024) and overall “onset to treatment” interval (2.42 vs. 2.13 hours p=0.003). Despite these differences, the “door to treatment” interval was not significantly

Table 3: Subset Analysis of Hispanic Patients Only: Baseline demographics and exam difference by rt-PA. Blood pressure and NIHSS is reported as the mean (standard deviation).

	Subset Analysis: Hispanic Patients Only		
	rt-PA(-) n = 194	rt-PA(+) n = 106	p-value
Age	65.7	67.02	0.510
Gender (Male)	112 (57.7%)	53 (50%)	0.225
Race (White)	190 (98.4%)	102 (96.2%)	0.266
Diabetes	80 (41.9%)	44 (42.3%)	1.000
Hypertension	136 (71.2%)	75 (72.1%)	0.893
Hyperlipidemia	56 (30.6%)	31 (30.1%)	1.000
Atrial Fibrillation	35 (18.3%)	28 (27.7%)	0.073
Coronary Artery Disease	35 (18.5%)	23 (21.9%)	0.541
Current Tobacco Use	43 (22.2%)	16 (15.1%)	0.172
Current Alcohol Use	26 (13.4%)	16 (15.1%)	0.729
Current SBP	153.36 (26.51)	146.86 (26.1)	0.091
Current DBP	82.02 (16.22)	81.51 (13.41)	0.775
Initial NIHSS (Total)	7.24 (9.05)	13.27 (9.31)	<.001

different (1.13 vs. 1.02 hours; p=0.07) (Table 2).

In the Hispanic-only subset, comparing rt-PA (+) to rt-PA (-) among Hispanics patients only, there was no difference for male sex (56%), Hypertension (71%), Hyperlipidemia (29%), Diabetes (43%), a trial Fibrillation (23%), Smoking (20%), and blood pressure. However, rt-PA (+) Hispanic patients differed from rt-PA (-) patients by presenting NIHSS at arrival (mean 13.27 vs. 7.24; p<0.001) (Table 3).

Discussion

Our analyses sought to determine the factors important to administration of rt-PA to Hispanic patients. We compared Hispanics to non-Hispanics, assessed the rt-PA (+) subset, and finally analyzed the Hispanic cohort itself for relevant factors. Our primary finding was that the rate of rt-PA treatment overall was not significantly different between Hispanics and non-Hispanics in our patient population. Some studies have reported on overall differences in rt-PA rates. A population study performed by Nasr, et al. demonstrated that white patients with AIS received rt-PA at a significantly higher rate than Hispanics (2.3% vs. 2%) [8]. Another study by A paricio, et al. showed that Hispanics still received rt-PA less often at both primary stroke center and non-primary stroke centers [9]. The major difference between our findings and other reports are both Hispanics and whites had high rt-PA treatment rates with no significant difference in treatment rates. It is affirming to the overall system that rt-PA treatment rates do not differ based on self-report of Hispanic ethnicity. While the national average rt-PA treatment rate is between 3.0% and 8.5% of all stroke admissions with some centers reporting a treatment rate up to 35%, we found that 33.5% of patients in our study received rt-PA [11,12]. Our dataset includes only acutely presenting stroke code patients within 12 hours of symptom onset, and does not report a percentage based on number of discharges. This is limitation to our study and may be a reason why the reported treatment rates are higher than the national average.

Our study also found that for rt-PA treated patients, although “door to treatment” intervals were not significantly different between Hispanics and non-Hispanics, “onset to treatment” intervals were longer, with this finding likely being driven by longer “onset to door” times. The reason for this difference is speculative, but may be due to poorer recognition, cultural differences, EMS activation differences, or even dispatch differences. Few studies have looked at the treatment intervals and delays in arrivals among Hispanics with AIS. Our study adds to the literature in that despite similar rt-PA treatment rates and “door to treatment” intervals, Hispanics patients with AIS are likely to have significant delay in hospital arrival time compared to whites. One hypothesis for this delay in arrival time is the mode of transportation Hispanic patient’s use. Neil, et al. showed Hispanics utilized EMS services less often compared to non-Hispanic whites [13]. However, that analysis did not show significant differences in onset to arrival intervals in Hispanics compared to non-Hispanic whites. Our study may have found a different result due to having a larger sample size. Another study found that Hispanic patients were more likely to present to larger hospitals. This was likely attributed to the urban location of hospitals [14]. Given this notable delay, more research should target the process of outside hospital factors that delay TPA treatment. We point out a need for clinician and policy maker investigation into possible regional and ethnic specific differences in stroke treatment access.

In our cohort of Hispanics only, there were no differences in demographic or co-morbidities between those treated with rt-PA and those who were not treated. This homogeneity is reassuring, showing that there were no systematic reasons why some Hispanics were treated and others were not. We did note that the presenting NIHSS was higher for those treated with rt-PA, which is in many ways an expected finding. Disabling deficits are generally treated to a greater degree than milder deficits [15]. When looking at the non-Hispanic white population, the percentage was similar (33.1%) showing that there was not a disproportionately lower treatment per NIHSS scale score for Hispanics. We plan to further assess this ratio in other race and ethnicity groups in subsequent analyses.

With the expected continued growth of the Hispanic American population, identifying the relevant factors (demographics, comorbidities and potential barriers) that contribute to rt-PA use in the Hispanic population, may improve the delivery of acute stroke care to this population. Our study supports the need to develop culturally relevant education programs to focus on treatment delays and differences in order to optimize rt-PA treatments. We are in the planning stages for tailored interventions which may work to minimize these barriers and optimize acute stroke treatment for this population.

Funding/Support

This study was supported by National Institutes of Health SPOTRIAS grant P50N5044148 and Stroke Net grant 5U10NS086535.

References

- Schwamm LH, Ali SF, Reeves MJ, Smith EE, Saver JL, Messe S, et al. Temporal trends in patient characteristics and treatment with intravenous thrombolysis among acute ischemic stroke patients at Get With The Guidelines-Stroke hospitals. *Circ Cardiovasc Qual Outcomes*. 2013; 6: 543-549.
- Cruz-Flores S, Rabinstein A, Biller J, Elkind MS, Griffith P, Gorelick PB, et al. Racial-ethnic disparities in stroke care: the American experience: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2011; 42: 2091-2116.
- Neyer JR, Greenlund KJ, Denny CH, Keenan NL, Casper M, Labarthe DR, et al. Prevalence of stroke: United States, 2005. *MMWR Morb Mortal Wkly Rep*. 2007; 56: 469-474.
- Siegler JE, Boehme AK, Albright KC, Martin-Schild S. Ethnic disparities trump other risk factors in determining delay to emergency department arrival in acute ischemic stroke. *Ethn Dis*. 2013; 23: 29-34.
- Hsia AW, Edwards DF, Morgenstern LB, Wing JJ, Brown NC, Coles R, et al. Racial disparities in tissue plasminogen activator treatment rate for stroke: a population-based study. *Stroke*. 2011; 42: 2217-2221.
- Boehme AK, Siegler JE, Mullen MT, Albright KC, Lyerly MJ, Monlezun DJ, et al. Racial and gender differences in stroke severity, outcomes, and treatment in patients with acute ischemic stroke. *J Stroke Cerebrovasc Dis*. 2014; 23: e255-e261.
- Bhattacharya P, Mada F, Salowich-Palm L, Hinton S, Millis S, Watson SR, et al. Are racial disparities in stroke care still prevalent in certified stroke centers? *J Stroke Cerebrovasc Dis*. 2013; 22: 383-388.
- Nasr DM, Brinjikji W, Cloft HJ, Rabinstein AA. Racial and ethnic disparities in the use of intravenous recombinant tissue plasminogen activator and outcomes for acute ischemic stroke. *J Stroke Cerebrovasc Dis*. 2013; 22: 154-160.
- Aparicio HJ, Carr BG, Kasner SE, Kallan MJ, Albright KC, Kleindorfer DO, et al. Racial Disparities in Intravenous Recombinant Tissue Plasminogen Activator Use Persist at Primary Stroke Centers. *J Am Heart Assoc*. 2015; 14.
- Schwamm LH, Reeves MJ, Pan W, Smith EE, Frankel MR, Olson D, et al. Race/ethnicity, quality of care, and outcomes in ischemic stroke. *Circulation*. 2010; 121: 1492-1501.
- Reeves MJ, Arora S, Broderick JP, Frankel M, Heinrich J, Hikenbottom S, et al. Acute stroke care in the US: results from 4 pilot prototypes of the paulcoverdell national acute stroke registry. *Stroke*. 2005; 36: 1232-1240.
- Kleindorfer D, Lindsell CJ, Brass L, Koroshetz W, Broderick JP. National US estimates of recombinant tissue plasminogen activator use: ICD-9 codes substantially underestimate. *Stroke*. 2008; 39: 924-928.
- Neil WP, Raman R, Hemmen TM, Ernstrom K, Meyer BC, Meyer DM, et al. Association of Hispanic ethnicity with acute ischemic stroke care processes and outcomes. *Ethn Dis*. 2015; 25:19-23.
- Lacy CR, Suh D, Bueno M, Kostis J. Delay in Presentation and Evaluation for Acute Stroke. *Stroke*. 2001; 32: 63-69.
- Schwamm LH, Ali SF, Reeves J, Smith EE, Saver JL, Messe S, et al. Temporal Trends in Patient Characteristics and Treatment with Intravenous Thrombolysis Among Acute Ischemic Stroke Patients at Get With the Guidelines-Stroke Hospitals. *Circ Cardiovasc Qual Outcomes*. 2015; 6: 543-549.