

## Clinical Image

# Tinea Imbricate

Manas Bajpai<sup>1</sup>, Nilesh Pardhe<sup>1</sup> and Pradkshana Vijay<sup>2\*</sup>

<sup>1</sup>Department of Oral and Maxillofacial Pathology, NIMS Dental College, India

<sup>2</sup>Department of Oral Pathology and Microbiology, KGMU, India

\*Corresponding author: Pradkshana Vijay, Senior Resident, Department of Oral Pathology and Microbiology, KGMU Lucknow, India

Received: November 20, 2017; Accepted: December 27, 2017; Published: December 29, 2017

## Clinical Image

An 8 year old presented with the chief complaint of several itchy hypo pigmented scars on her left side of face since 2 months, but other parts of her body were normal. Skin scrapings were microscopically examined in KOH 10% and numerous irregular, branch and septate hyaline hyphae were observed. The patient was successfully treated with oral terbinafine 250mg/day, topical clotrimazole (1% ointment), topical miconazole (2% cream) two times daily and potassium permanganate for daily washing for four weeks. After 1 year of follow up no complaint, reported about the relapse of infection.

Tinea imbricata is an unusual form of tinea corporis caused by the strictly anthropophilic dermatophyte *T. concentricum*.



Figure 1: Erythematous itchy lesion of face.