

Editorial

Legacy of Maternal Obesity

Torsoni AS and Torsoni MA^{1*}

¹Faculty of Applied Sciences, University of Campinas, Brazil

***Corresponding author:** Torsoni MA, Faculty of Applied Sciences, University of Campinas, Rua Pedro Zaccaria, 1300, Jardim Santa Luiza, Limeira, São Paulo, CEP 13484350, Brazil, Tel: 551937016705; Fax: 55 1937016680; Email: marcio.torsoni@fca.unicamp.br

Received: September 13, 2014; **Accepted:** September 16, 2014; **Published:** September 18, 2014

Abstract

Maternal obesity and consumption of high-fat diet can influence fetal development and growth. Placental inflammation leads to molecular alterations in different tissues in the off spring. Hypothalamic tissues can show early changes in the expression of neuropeptides, which contribute to damages in metabolic control and energy homeostasis. Inflammation of the placenta can affect orexigenic/anorexigenic neuropeptides, reward system, and energy expenditure, which in turn contribute to obesity and diabetes in the offspring. These changes can be caused by epigenetic mechanisms and can be passed on from the mother to the offspring.

Introduction

Obesity constitutes a growing health problem throughout the world. Prevalence of obesity in women of child bearing age [1] and in children is worrying because both increase the risk of metabolic diseases in the later life of the offspring. Human and animal studies have shown that changes in early life can have deleterious effects in adult life. These findings have increased the interest in the effects of maternal obesity on disease risk in the offspring [2-4].

The effect of maternal over nutrition during fetal development increases the risk of diseases in the offspring during its adult life. Different types of stress during critical periods of early development permanently alter an organism's physiology and metabolism. This phenomenon is called metabolic programming and has originated from fetal hypothesis proposed by Barker [5].

However, some important questions need to be answered regarding metabolic imprinting. These include understanding the role of gestation and lactation in the ill outcomes observed in the offspring and whether maternal obesity or High-Fat Diet (HFD) consumption during gestation and/or lactation could affect differently fetal development and result in metabolic damage in the offspring. Most studies have used animal models in which obesity is induced by feeding HFD during gestation and/or lactation. Therefore, it is impossible to establish the direct relation between metabolic damage in the offspring and maternal obesity and HFD consumption during pregnancy. However, despite the difficulty in the interpretation of results from these models, most studies have shown damages in different metabolic pathways and tissues. Maternal obesity and HFD consumption are associated with fatty liver [6,7] micro RNA expression modulation [8,9], insulin resistance [6,10], and cognitive disruption [11,12].

Hypothalamic tissue controls food intake, energy expenditure, reward system, and peripheral metabolism. Molecular changes in fetal and adult hypothalamus are related to maternal HFD consumption during pregnancy. Gupta and colleagues [13] showed increased mRNA levels of proopiomelanocortin, melanocortin receptor-4, neuropeptide Y, and agouti-related polypeptide in fetal hypothalamus from rats [13]. Up regulation of the orexigenic system can increase food intake and body weight. For example, increased

expression of orexigenic neuropeptides in the postnatal period can induce high intake of milk during breastfeeding, thus contributing to excessive weight gain. Moreover, in mice, perinatal exposure to HFD can produce a more deleterious response to HFD challenge in later life even after an interval of normal diet in mice [14], suggesting permanent molecular alterations. Epigenetic modifications in DNA can result in permanent changes in the expression profile of genes related to metabolism and energy homeostasis.

Epigenetic modifications have been described in genes associated with reward, which can affect preference for palatable foods [15]. Expression of both μ -opioid receptor and preproenkephalin was increased in the nucleus accumbens, prefrontal cortex, and hypothalamus of mice born to dams that consumed HFD [15]. Moreover, mice born to HFD-fed dams during pregnancy and lactation showed hypomethylation in the dopamine reuptake transporter promoter region, μ -opioid receptor, and preproenkephalin. Reduced methylation of CpG in the promoter region permits the binding of a transcription factor to DNA and increases gene expression.

Maternal obesity and HFD consumption during pregnancy and/or lactation alter the levels of hormones (leptin and insulin), nutrients (fatty acids and glucose), and inflammatory cytokines [6,16] in the blood. This can affect the environment of the developing offspring. Some authors have shown proinflammatory components in the placenta of humans [17], rats [18], and sheep [19] in response to HFD consumption and obesity. However, uterine changes and embryonic inflammation precedes placentation. A study has shown that in rats, gene expression in the blastocyst at 4.5 days postcoitum is clearly influenced by maternal obesity. The same study has shown that placental inflammation increases due to the accumulation of ectopic lipids and expression of lipid metabolic genes in the uterus [18]. In contrast, a recent study has shown that obese dams show macrophage infiltration in the adipose tissue and liver but reversal of obesity-induced inflammation during gestation [20].

Initially, these findings may seem contradictory. However, pregnancy induces metabolic changes in the mother to ensure passage of nutrients to the fetus; therefore, reducing the availability of fatty acids and glucose in the maternal adipose tissue. HFD consumption during pregnancy increases placental transport of amino acids, glucose, and fatty acids [21], thus contributing to proinflammatory

environment and ectopic lipid accumulation. Saturated fatty acids can activate toll-like receptor 4 (TLR4) and stimulate cytokine expression. Cotyledonary tissue obtained from obese pregnant mothers showed increased expression of TLR4 and macrophage markers (CD11b, CD14 and CD68) [19]. Larger and colleagues [22] have shown that high levels of IL-6 stimulate fatty acid accumulation in human primary trophoblast cells. This can contribute to excessive nutrient transfer in conditions associated with elevated maternal IL-6, such as obesity and gestational diabetes [22]. Although the precise mechanism is unknown, a recent study has shown that palmitic acid-mediated placental inflammation in human placenta choriocarcinoma cell line requires JNK signaling and recruitment of early growth response protein-1 (EGR-1) on cytokine promoters [23].

Although controversial, increased lipid circulation seems to be a mediator of fetal programming [24-26]. Epigenetic modifications (histone acetylation and methylation) and altered expression of enzymes that regulate histone and DNA methylation in the placenta have been observed in the offspring of dams who are fed HFD during pregnancy [26,27]. Monitoring of maternal blood lipid profile during gestation and breast milk composition during lactation is important to avoid permanent changes in the fetus that are caused by placental inflammation because gestational and lactational periods are important for neurogenesis and neuronal plasticity.

In summary monitoring of maternal nutrition and weight gain during the perinatal period is important to prevent metabolic disorders in the offspring. Furthermore, this paves a way for research in appropriate nutrition of obese mothers to prevent placental inflammation and metabolic complications in the offspring.

References

- Guelinckx I, Devlieger R, Beckers K, Vansant G. Maternal obesity: pregnancy complications, gestational weight gain and nutrition. *Obes Rev.* 2008; 9: 140-150.
- Drake AJ, Reynolds RM. Impact of maternal obesity on offspring obesity and cardiometabolic disease risk. *Reproduction.* 2010; 140: 387-398.
- Dyer JS, Rosenfeld CR. Metabolic imprinting by prenatal, perinatal, and postnatal overnutrition: a review. *Semin Reprod Med.* 2011; 29: 266-276.
- King V, Norman JE, Seckl JR, Drake AJ. Post-weaning diet determines metabolic risk in mice exposed to over nutrition in early life. *Reprod Biol Endocrinol.* 2014; 12: 73.
- Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet.* 1986; 1: 1077-1081.
- Ashino NG, Saito KN, Souza FD, Nakutz FS, Roman EA, Velloso LA, et al. Maternal highfat feeding through pregnancy and lactation predisposes mouse offspring to molecular insulin resistance and fatty liver. *The Journal of nutritional biochemistry.* 2012; 23: 341-348.
- McCurdy CE, Bishop JM, Williams SM, Grayson BE, Smith MS, Friedman JE, et al. Maternal high-fat diet triggers lipotoxicity in the fetal livers of nonhuman primates. *J Clin Invest.* 2009; 119: 323-335.
- Benatti RO, Melo AM, Borges FO, Ignacio-Souza LM, Simino LA, Milanski M, et al. Maternal high-fat diet consumption modulates hepatic lipid metabolism and microRNA-122 (miR-122) and microRNA-370 (miR-370) expression in offspring. *Br J Nutr.* 2014; 111: 2112-2122.
- Zhang J, Zhang F, Didelot X, Bruce KD, Cagampang FR, Vatish M, et al. Maternal high fat diet during pregnancy and lactation alters hepatic expression of insulin like growth factor-2 and key micro RNAs in the adult offspring. *BMC Genomics.* 2009; 10: 478.
- Melo AM, Benatti RO, Ignacio-Souza LM, Okino C, Torsoni AS, Milanski M, et al. Hypothalamic endoplasmic reticulum stress and insulin resistance in offspring of mice dams fed highfat diet during pregnancy and lactation. *Metabolism: clinical and experimental.* 2014; 63: 682-692.
- Bilbo SD, Tsang V. Enduring consequences of maternal obesity for brain inflammation and behavior of offspring. *FASEB journal. official publication of the Federation of American Societies for Experimental Biology.* 2010; 24: 2104-2115.
- Tozuka Y, Kumon M, Wada E, Onodera M, Mochizuki H, Wada K. Maternal obesity impairs hippocampal BDNF production and spatial learning performance in young mouse offspring. *Neurochem Int.* 2010; 57: 235-247.
- Gupta A, Srinivasan M, Thamadolok S, Patel MS. Hypothalamic alterations in fetuses of high fat diet-fed obese female rats. *J Endocrinol.* 2009; 200: 293-300.
- Kruse M, Seki Y, Vuguin PM, Du XQ, Fiallo A, Glenn AS, et al. High-fat intake during pregnancy and lactation exacerbates high-fat diet-induced complications in male offspring in mice. *Endocrinology.* 2013; 154: 3565-3576.
- Vucetic Z, Kimmel J, Totoki K, Hollenbeck E, Reyes TM. Maternal high-fat diet alters methylation and gene expression of dopamine and opioid-related genes. *Endocrinology.* 2010; 151: 4756-4764.
- Kępczyńska MA, Wargent ET, Cawthorne MA, Arch JR, O'Dowd JF, Stocker CJ. Circulating levels of the cytokines IL10, IFN β and resistin in an obese mouse model of developmental programming. *J Dev Orig Health Dis.* 2013; 4: 491-498.
- Challier JC, Basu S, Bintein T, Minium J, Hotmire K, Catalano PM, et al. Obesity in pregnancy stimulates macrophage accumulation and inflammation in the placenta. *Placenta.* 2008; 29: 274-281.
- Shankar K, Zhong Y, Kang P, Lau F, Blackburn ML, Chen JR, et al. Maternal obesity promotes a proinflammatory signature in rat uterus and blastocyst. *Endocrinology.* 2011; 152: 4158-4170.
- Zhu MJ, Du M, Nathanielsz PW, Ford SP. Maternal obesity up-regulates inflammatory signaling pathways and enhances cytokine expression in the mid-gestation sheep placenta. *Placenta.* 2010; 31: 387-391.
- Ingvorsen C, Thysen AH, Fernandez-Twinn D, Nordby P, Nielsen KF, Ozanne SE, et al. Effects of pregnancy on obesity-induced inflammation in a mouse model of fetal programming. *Int J Obes (Lond).* 2014.
- Sferruzzi-Perri AN, Vaughan OR, Haro M, Cooper WN, Musial B, Charalambous M, et al. An obesogenic diet during mouse pregnancy modifies maternal nutrient partitioning and the fetal growth trajectory. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology.* 2013; 27: 3928-3937.
- Lager S, Jansson N, Olsson AL, Wennergren M, Jansson T, Powell TL. Effect of IL-6 and TNF- α on fatty acid uptake in cultured human primary trophoblast cells. *Placenta.* 2011; 32: 121-127.
- Saben J, Zhong Y, Gomez-Acevedo H, Thakali KM, Borengasser SJ, Andres A, et al. Early growth response protein-1 mediates lipotoxicity-associated placental inflammation: role in maternal obesity. *American journal of physiology Endocrinology and metabolism.* 2013; 305: E1-14.
- Krasnow SM, Nguyen ML, Marks DL. Increased maternal fat consumption during pregnancy alters body composition in neonatal mice. *Am J Physiol Endocrinol Metab.* 2011; 301: E1243-1253.
- Masuyama H, Hiramatsu Y. Effects of a high-fat diet exposure in utero on the metabolic syndrome-like phenomenon in mouse offspring through epigenetic changes in adipocytokine gene expression. *Endocrinology.* 2012; 153: 2823-2830.
- Strakovsky RS, Zhang X, Zhou D, Pan YX. Gestational high fat diet programs hepatic phosphoenolpyruvate carboxykinase gene expression and histone modification in neonatal offspring rats. *J Physiol.* 2011; 589: 2707-2717.
- Gabory A, Ferry L, Fajardy I, Jouneau L, Gothié JD, Vigé A, et al. Maternal diets trigger sex-specific divergent trajectories of gene expression and epigenetic systems in mouse placenta. *PLoS One.* 2012; 7: e47986.