

## Research Article

# The Role of Health Locus of Control in Walking Time, Hand Grip Strength and Alternate Stepping in a 60-93 Year-Old General Population

Ekström H<sup>1\*</sup>; Elmståhl S<sup>2</sup>; Wranger SL<sup>2</sup><sup>1</sup>The Department of Clinical Sciences in Malmö, Division of Geriatric Medicine, Skåne University Hospital, Lund University, Malmö, Sweden<sup>2</sup>Division of Geriatric Medicine, Department of Clinical Sciences in Malmö, Lund University, Sweden**\*Corresponding author: Ekström H**

Division of Geriatric Medicine, Lund University, Jan Waldenströms gata 35, 205 02 Malmö, Sweden.

Tel: +46 703 92 86 10; Fax: +46 391313

Email: henrik.ekstrom@med.lu.se

**Received:** March 13, 2023**Accepted:** April 19, 2023**Published:** April 26, 2023**Abstract**

Individuals with an internal locus of control have been shown to be more self-confident and to perform better in several areas, compared to those with an external locus of control. Few studies have investigated the relationship between Health Locus of Control (HLoC) and physical tests. The aim of this study was to investigate whether and to what extent internal, chance and external HLoC is associated with the outcome of walking speed, hand grip strength and alternate stepping physical tests.

In this cross-sectional study, a total of 3,819 individuals, aged 60-93 years, constituted the study population. Associations between internal, chance and external HLoC and the results from the physical tests were examined in linear regression models, adjusted for sex, age, education, moving-related pain in upper extremities, back and lower extremities, heart disease, lung disease, depressive mood and cognition.

The regression models indicated that a higher result on the internal scale was associated with faster walking, a stronger hand grip and a greater number of steps, while a higher result on the external scale was associated with slower walking, lower hand grip strength and fewer steps.

The findings suggest that in a clinical setting, assessment of HLoC may contribute to the understanding of physical performance in terms of walking speed, grip strength and alternate stepping among adults 60-93 years old. Modification to ensure a stronger internal control has the potential to improve the performance of these physical tests, which could be worth considering in a medical or physiotherapeutic assessment.

**Keywords:** Health locus of control; Walking speed; Hand grip strength; Alternate stepping; Older adults

**Abbreviations:** LoC; HLoC; WS; HGS; AS; GÅS; SNAC; BMI; MADRS; COPD; MMSE; ICC

**Introduction**

The theory of a control focus (LoC), introduced by Rotter in 1966 within the framework of Social learning theory [1], describes how an individual's life situation can be partly explained on the basis of an internal or external control locus. That is, either taking responsibility for and acting on one's own initiative in conjunction with different life events or believing that other people or external forces such as chance are in control or have the greatest influence [2].

Health Locus of Control (HLoC), a control locus set in a health perspective, was first described by Wallston in 1976 [3]. HLoC refers to the extent an individual believes her/his health to be mostly controlled by an internal, chance or external locus. Individuals with an internal HLoC consider that their own responsibility and actions form the basis for their health and well-being, while those with a chance or external locus believe that their health is largely a matter of luck or in the hands of other people.

ple such as doctors or healthcare professionals. This means that for those with a chance or external locus, health is something outside of themselves that they cannot influence or control [4].

Many studies have shown the importance of HLoC for general well-being and that health can be affected depending on whether an individual has an internal, chance or external health locus. Internal HLoC has been associated with higher quality of life [5], improved smoking habits [6], better self-rated health [7], maintaining physical function after hospitalization [8], less prone to depressive conditions [9], lower prevalence of cardiovascular diseases [10], greater tolerance of pain [11] and better adherence to anti-hypertensive treatment [12]. Furthermore, individuals with an internal HLoC are more likely to participate in health-promoting activities [13]. This differs from external HLoC, which has been shown to be associated with trust in health professionals [14] and often with a more passive attitude to health problems [15].

Walking Speed (WS), Hand Grip Strength (HGS) and Alternate Stepping (AS) have been used to provide an assessment of health status and functional ability in both healthy individuals and those with impaired health [16-20]. These tests examine several organ systems such as muscles, bones, heart-lung and the nervous system [21,22]. WS and HSG are also employed as criteria in estimating frailty [23].

To the authors' knowledge, no previous studies have examined the importance of HLoC in relation to WS and HGS among older adults. If HLoC can affect the ability to perform these tests, it would be an important factor to consider both for the physical performance itself and as an essential underlying personal factor in the assessment of physical health, functional ability or frailty. Thus, the aim of this study was to investigate whether and to what extent internal, chance and external HLoC are associated with the outcome of walking speed, hand grip strength and alternate stepping physical tests in a general population aged 60-93 years, adjusted for socio-demographics and health status.

**Materials and Methods**


**Study Population**

In this cross-sectional study participants were drawn from the longitudinal Good Aging in Skåne (GÅS) Project, which began in 2001 and is a part of the Swedish National Study on Aging and Care (SNAC). The design of the GÅS and SNAC study is described elsewhere [24,25].

Participants randomized from the national population register were invited to take part in the study by letter and written informed consent was obtained. A total of 6,991 eligible individuals were invited in two waves (Figure 1). From the first wave between 2001 and 2004, 2,931 (60.0%) out of 4,893 agreed to participate, while from the second wave between 2006 and 2012, 1,523 (72.6%) out of 2,098 agreed to participate. The first wave included participants in nine age-cohorts (60, 66, 72, 78, 81, 84, 87, 90 and 93 years) and the second wave participants in two age cohorts (60 and 81 years).

A total of 254 individuals from the first wave and 184 individuals from the second wave were excluded, as they did not complete any of the HLoC-scales. An additional 144 individuals from the first wave and 53 from the second wave were excluded, as they did not perform any of the physical tests (walking 15 m at normal speed, HGS or AS). The first and second waves

were then merged, after which the study population consisted of 3,819 participants, 1,725 (45.2%) men and 2,094 (54.8%) women (Figure 1) (Table 1).



**Figure 1:** Flow sheet explaining the enrolment of participants in the first wave 2001-2004 and in the second wave 2006-2012.

**Table 1:** Description of study sample, N=3,819.

|  | Study sample n(%) | Internal Missing n(%) |
|--|-------------------|-----------------------|
| Sex  |                   | -                     |
| Men  | 1,725(45.2)       |                       |
| Women                                      | 2,094(54.8)       |                       |
| Age, Mean (SD)                             | 69.4(9.8)         | -                     |
| Age decade                                 |                   | -                     |
| 60   | 2,335(61.7)       |                       |
| 70   | 57(13.5)          |                       |
| 80   | 817(21.4)         |                       |
| 90   | 130(3.4)          |                       |
| Education                                  |                   | 11(0.3)               |
| Primary school                             | 1,787(46.8)       |                       |
| Secondary school                           | 1,110(29.1)       |                       |
| University                                 | 911(23.9)         |                       |
| BMI  |                   | 82(2.1)               |
| Underweight                                | 139(3.6)          |                       |
| Normal                                     | 1,302(34.1)       |                       |
| Overweight                                 | 1,593(41.7)       |                       |
| Obese                                      | 703(18.4)         |                       |
| Moving-related pain Upper Extremities      |                   | 2(0.1)                |
| Yes  | 1,231(32.2)       |                       |
| No   | 2,586(67.7)       |                       |
| Moving-related pain Back/lower extremities |                   | 5(0.1)                |
| Yes  | 1,891(49.5)       |                       |
| No   | 1,926(50.4)       |                       |
| Heart disease                              |                   | 5(0.1)                |
| Yes  | 688(18.0)         |                       |
| No   | 3,126(81.9)       |                       |
| Lung disease                               |                   | 5(0.1)                |
| Yes  | 476(12.5)         |                       |
| No   | 3,382(87.4)       |                       |
| Depressive mood                            |                   | 111(2.9)              |
| Yes  | 533(14.0)         |                       |
| No   | 3,175(83.1)       |                       |
| MMSE                                       |                   | 59(1.5)               |
| ≤24p                                       | 421(12.6)         |                       |
| >24p                                       | 3,279(85.9)       |                       |

## Socio-Demographics, Health Status, Lifestyle Habits and Health Locus of Control

Structured interviews including questions about health variables were carried out by trained medical staff in accordance with predefined research protocols. Self-reported questionnaires were used to obtain data on sociodemographics, lifestyle habits and HLoC. Assessments took place either at the research centre or, in cases of ill health, in participants' homes.

### Health Variables, Lifestyle Habits

Health variables covered heart disease, lung disease, Body Mass Index (BMI), depressive mood, cognition and experience of pain in the upper extremities or back/lower extremities when moving. Heart disease included myocardial infarction, angina pectoris or clinical heart failure. Lung disease comprised Chronic Obstructive Pulmonary Disease (COPD), asthma or tuberculosis. Moving-related pain from the upper extremities covered pain from the shoulders, elbows, wrists and hands. Pain from the lower extremities or back when moving involved pain from the back, hips, knees and feet.

To check the medical history after the medical examination, reported diseases were verified through the National Diagnosis Registry and medical records, for which the participants' permission had been obtained. Weight (kg) was measured using a balance scale with a precision of  $\pm 50$ g with participants in light clothing but with no shoes [26]. Height was measured without shoes to the nearest 0.1cm using a scale fixed to a wall with the participant standing erect with heels and shoulders against the wall and a straight fixed gaze [26]. Body Mass Index (BMI) was then calculated as weight (kg)/height ( $m^2$ ) and cut off set to  $<18.5$ kg/ $m^2$  underweight,  $18.5$ - $24.9$ kg/ $m^2$  normal (healthy) weight,  $25.0$ - $29.9$ kg/ $m^2$  overweight and  $\geq 30$ kg/ $m^2$  obese [27].

The Montgomery-Åsberg Depression Rating Scale (MADRS) was used as a test of depressive mood. The scale ranges from 0-60 points and a score  $\geq 7$  points was used as a cut off for depressive mood [28]. The MADRS has previously been validated for older adults [29]. The Mini Mental State Examination (MMSE) was used to assess global cognitive function [30]. The scale ranges from 0-30 points and 24 points was used as a cut off for cognitive impairment [31].

### Physical Tests

The physical tests consisted of walking 15m at a normal (comfortable) and maximum speed, followed by the HGS test and the AS stair touching test. These physical tests were chosen in order to gain a broad impression of the participants' physical ability in terms of mobility, muscle strength, balance and coordination [32]. All tests were performed at the Department of Geriatrics, Malmö University Hospital. A trained registered research nurse instructed the participants about how the tests should be performed, monitored the performance but provided no encouragement during the assessment. The participants wore their normal clothes and shoes, and walking aids were allowed [32].

**Walking 15 metres at normal and maximum speed:** WS was used as a measure of functional mobility [33]. Following a dynamic start, the participants were asked to walk 15m at a normal (comfortable) and a maximum speed. The test took place in a hospital corridor and participants were allowed to accelerate and decelerate for several metres before and after the test. To ensure that the participants achieved their individual maximum

speed, the nurse instructed them that the test was to be performed as quickly as possible without running. The time taken to walk was recorded using a digital stopwatch. Each test (normal and maximum WS) was performed once, and participants were allowed to rest for one minute between tests. High intra-class correlation (ICC  $>0.90$ ) has been reported for walking 15m at both normal and maximum speed [34].

**Hand grip strength:** The Grippit<sup>®</sup>, a device that measures HGS, was used for this test [35]. Participants performed a standard testing procedure in accordance with the test leader's instructions. The hand grip device and a forearm support were mounted on a transportable base, ensuring correct arm and hand grip positions, and the participants started to squeeze the handle on command [34]. The test was carried out twice on each hand and the maximum force noted. The best result (maximum force) was used in the analysis. High intraclass correlation has been reported (ICC = 0.97) for both hands [36].

**Alternate stepping:** Participants stood with feet parallel at 5cm from the front of a 7.5cm high block stably positioned against a wall. They were asked to place one foot on the block and then return it to the floor as quickly as possible. A research nurse stood close by for safety but did not assist in the performance. The total number of steps completed in 15 seconds was recorded for both the right and the left lower extremity [37]. The best value from the tests (the dominant lower extremity) was used in the analysis. High intraclass correlation has been reported (ICC=0.78) [38].

### Health Locus of Control

Internal and external HLoC were assessed by the Multidimensional Health Locus of Control (MHLC) questionnaire version B [39]. The questionnaire consists of three subscales for estimating internal, external and chance locus respectively. Each scale ranges from 0 – 39 points and contains 6 positively formulated statements based on a Likert scale with the following 5 response alternatives; agrees exactly = 1, agrees quite well = 2, neither agrees nor disagrees = 3, does not really agree = 4, does not agree at all = 5.

Higher scores on the internal subscale indicate that attitudes and behaviour in relation to health are more a matter of personal beliefs about what one can do oneself to promote better health. In contrast, higher scores on the external and chance scales indicate that health is more in the hands of other people, such as healthcare professionals, or that health cannot be influenced but is mostly a result of chance [39]. Cronbach's alpha for internal, chance and external HLoC was 0.74, 0.81 and 0.76, respectively.

### Statistical Analysis

Descriptive statistics pertaining to the study population are presented in Table 1. Differences in proportions were tested with the chi-square ( $\chi^2$ ) test.

Results of the WS and HGS physical tests according to sex, age, education, co-habiting, BMI, moving-related pain in the upper extremities, moving-related pain in the back/lower extremities, heart disease, lung disease and depressive mood are presented as means and standard deviations. Significance tests for two groups were performed with Student's T-test and for three groups with ANOVA (Table 2). Correlation between physical tests and internal and external HLC was calculated using Pearson's correlation (r) (Table 2).

**Table 2:** Results of the physical tests, mean and SD, according to sex, age, education, smoking habits, BMI, moving-related pain in the upper extremities, moving-related pain in the back/lower extremities, heart disease, lung disease, depressive mood and correlations, Pearson's (r), between physical tests and internal, chance and external HLoC (N=3,819).

| Physical tests<br>Variables                    | Walking 15 metres<br>normal speed |         | Walking 15 metres<br>Maximum speed |         | Hand grip strength<br>maximum force<br>dominant hand |         | Alternate stepping<br>number of steps |         |
|--|-----------------------------------|---------|------------------------------------|---------|--|---------|---------------------------------------|---------|
|  | m/s(SD)                           | p-value | m/s(SD)                            | p-value | Kg(SD)   | p-value | Number(SD)                            | p-value |
| Sex  |                                   |         |                                    |         |  |         |                                       |         |
| Men  | 1.39(0.26)                        | <0.001  | 1.80(0.39)                         | <0.001  | 38.1(10.33)  | <0.001  | 18.0(5.08)                            | <0.001  |
| Women  | 1.30(0.28)                        |         | 1.58(0.36)                         |         | 20.4(6.67)   |         | 15.9(4.89)                            |         |
| Age(decade)                                    |                                   |         |                                    |         |  |         |                                       |         |
| 60   | 1.46(0.22)                        | <0.001  | 1.85(0.32)                         | <0.001  | 31.6(12.54)  | <0.001  | 18.7(4.51)                            | <0.001  |
| 70   | 1.25(0.24)                        |         | 1.54(0.32)                         |         | 26.1(10.11)  |         | 15.0(4.25)                            |         |
| 80   | 1.10(0.24)                        |         | 1.34(0.32)                         |         | 21.6(9.09)   |         | 13.0(4.23)                            |         |
| 90   | 0.95(0.23)                        |         | 1.16(0.39)                         |         | 18.6(8.30)   |         | 11.2(3.61)                            |         |
| Education                                      |                                   |         |                                    |         |  |         |                                       |         |
| Primary school                                 | 1.27(0.27)                        | <0.001  | 1.57(0.37)                         | <0.001  | 26.2(11.89)  | <0.001  | 15.2(4.76)                            | <0.001  |
| Secondary school                               | 1.36(0.27)                        |         | 1.71(0.37)                         |         | 29.2(12.68)  |         | 17.4(4.81)                            |         |
| University                                     | 1.45(0.25)                        |         | 1.88(0.36)                         |         | 31.3(12.02)  |         | 19.4(4.79)                            |         |
| BMI  |                                   |         |                                    |         |  |         |                                       |         |
| Underweight                                    | 1.32(0.31)                        | <0.001  | 1.63(0.39)                         | <0.001  | 22.8(9.71)   | <0.001  | 15.7(5.23)                            | <0.001  |
| Normal   | 1.38(0.28)                        |         | 1.74(0.39)                         |         | 27.4(11.82)  |         | 17.4(5.05)                            |         |
| Overweight                                     | 1.35(0.28)                        |         | 1.71(0.38)                         |         | 30.1(12.66)  |         | 17.1(5.08)                            |         |
| Obese  | 1.26(0.27)                        |         | 1.56(0.38)                         |         | 28.3(12.31)  |         | 15.6(4.81)                            |         |
| Moving-related pain,<br>upper extremities      |                                   |         |                                    |         |  |         |                                       |         |
| Yes  | 1.31(0.27)                        | <0.001  | 1.63(0.38)                         | <0.001  | 25.0(12.08)  | <0.001  | 16.2(4.91)                            | <0.001  |
| No   | 1.36(0.28)                        |         | 1.71(0.39)                         |         | 30.2(12.08)  |         | 17.2(5.13)                            |         |
| Moving-related pain,<br>back/lower extremities |                                   |         |                                    |         |  |         |                                       |         |
| Yes  | 1.29(0.29)                        | <0.001  | 1.61(0.38)                         | <0.001  | 26.3(12.12)  | <0.001  | 16.1(4.95)                            | <0.001  |
| No   | 1.39(0.26)                        |         | 1.76(0.39)                         |         | 30.7(12.3)   |         | 17.6(5.08)                            |         |
| Heart disease                                  |                                   |         |                                    |         |  |         |                                       |         |
| Yes  | 1.17(0.30)                        | <0.001  | 1.46(0.40)                         | <0.001  | 27.1(12.53)  | 0.002   | 14.4(4.97)                            | <0.001  |
| No   | 1.38(0.26)                        |         | 1.73(0.37)                         |         | 28.8(12.24)  |         | 17.4(4.95)                            |         |
| Lung disease                                   |                                   |         |                                    |         |  |         |                                       |         |
| Yes  | 1.35(0.28)                        | 0.003   | 1.61(0.38)                         | <0.001  | 26.4(11.32)  | <0.001  | 16.2(5.05)                            | 0.004   |
| No   | 1.29(0.28)                        |         | 1.70(0.39)                         |         | 28.8(12.42)  |         | 17.0(5.07)                            |         |
| Depressive mood                                |                                   |         |                                    |         |  |         |                                       |         |
| Yes  | 1.23(0.28)                        | <0.001  | 1.52(0.39)                         | <0.001  | 23.9(10.90)  | <0.001  | 14.9(5.02)                            | <0.001  |
| No   | 1.36(0.29)                        |         | 1.72(0.38)                         |         | 29.48(12.30)   |         | 17.3(4.94)                            |         |
| MMSE   |                                   |         |                                    |         |  |         |                                       |         |
| ≤24p   | 1.16(0.29)                        | <0.001  | 1.42(0.38)                         | <0.001  | 23.2(10.66)  | <0.001  | 13.4(4.82)                            | <0.001  |
| >24p   | 1.37(0.27)                        |         | 1.72(0.38)                         |         | 29.3(12.32)  |         | 17.4(4.91)                            |         |
| Internal HLoC(Pearson's [r])                   | 0.126                             | <0.001  | 0.123                              | <0.001  | 0.146  | <0.001  | 0.099                                 | <0.001  |
| Chance HLoC(Pearson's [r])                     | -0.293                            | <0.001  | -0.324                             | <0.001  | -0.164   | <0.001  | -0.313                                | <0.001  |
| External HLoC(Pearson's [r])                   | -0.296                            | <0.001  | -0.333                             | <0.001  | -0.165   | <0.001  | -0.316                                | <0.001  |

Standard multivariate linear regression models were constructed to analyse the associations between the time taken to walk 15m, maximum HGS and number of AS as dependent variables and HLoC as an independent variable, adjusted for sex, age decade, education, BMI, moving-related pain in the upper extremities, moving-related pain in the back/lower extremities, heart disease, lung disease, depressive mood and outcome of the MMSE. The variables were simultaneously entered into the regression models and all variables with the sole exception of the HLoC scales were used as dummies (Table 3). For all regression models assumptions of normality, linearity and homoscedasticity were controlled for by examining the residual scatterplots, i.e., predicted values of dependent variables against residuals, and no deviations were observed [40]. Multicollinearity was tested for and none of the included variables had a variance inflation factor >2.0 [41]. For all models, the explained variance was assessed by the adjusted R-square and overall

significance by the F-test. The level of significance was set to <0.05. An attrition analysis was carried out to compare Participants with non-participants (n=635, 14.2%) and differences in proportions regarding included variables were tested with the chi-square ( $\chi^2$ ) test (Table 4).

Data analysis was performed using SPSS for Windows, version 24.0 (IBM 211 Corporation, Armonk, NY, USA).

### Ethical Considerations

The study was conducted in accordance with the Helsinki Declaration [42] and approved by the regional ethics committee at Lund University 2010-2012, registration no. LU 744-00. All participants provided their written consent and allowed retrieval of information from the National Patient Register medical records. Participants were informed that they could withdraw from the study at any time.

**Table 3:** Adjusted multiple linear regression models with walking speed(m/s), hand grip strength(kg) and alternate stepping(Nb) as dependent variables and HLoC scales as independent variables, adjusted for sex, age, education, BMI, moving-related pain in the upper extremities, moving-related pain in the back/lower extremities, heart disease, lung disease, depressive mood and cognition(MMSE).

| Physical tests                  | Walking 15m at normal speed<br>n =3,288 |                   |         | Walking 15m at maximum speed<br>n =3,274 |                   |         | Maximum hand grip strength dominant hand<br>n = 3,483 |                   |         | Alternate stepping, number of steps<br>n = 3,444 |                 |         |
|---------------------------------|---|-------------------|---------|--|-------------------|---------|---|-------------------|---------|--|-----------------|---------|
|                                 | B <sup>a</sup>                          | 95% CI            | p-value | B <sup>a</sup>                           | 95% CI            | p-value | B <sup>a</sup>  | 95% CI            | p-value | B <sup>a</sup>                                   | 95% CI          | p-value |
| Sex women (ref. men)            | -0.043                                  | -0.058/<br>-0.028 | <0.001  | -0.165                                   | -0.185/<br>-0.145 | <0.001  | -16.63  | -17.14/<br>-16.12 | <0.001  | -1.34  | -1.62/<br>-1.06 | <0.001  |
| Age (ref. decade 60)            |   |                   |         |  |                   |         |   |                   |         |  |                 |         |
| 70                              | -0.158                                  | -0.180/<br>-0.137 | <0.001  | -0.220                                   | -0.249/<br>-0.192 | <0.001  | -4.27   | -5.03/<br>-3.52   | <0.001  | -2.50  | -2.91/<br>-2.08 | <0.001  |
| 80                              | -0.293                                  | -0.313/<br>-0.272 | <0.001  | -0.387                                   | -0.414/<br>0.360- | <0.001  | -7.87   | -8.59/<br>-7.16   | <0.001  | -4.18  | -4.57/<br>-3.80 | <0.001  |
| 90                              | -0.425                                  | -0.469/<br>-0.381 | <0.001  | -0.551                                   | -0.609/<br>-0.493 | <0.001  | -9.96   | -11.50/<br>-8.52  | <0.001  | -5.91  | -6.77/<br>-5.05 | <0.001  |
| Education (ref. primary school) |   |                   |         |  |                   |         |   |                   |         |  |                 |         |
| Secondary school                | 0.022                                   | 0.005/<br>0.039   | 0.011   | 0.030                                    | 0.008/<br>0.053   | 0.008   | -0.20   | -0.79/<br>0.39    | 0.506   | 0.90   | 0.58/<br>1.22   | <0.001  |
| University                      | 0.39                                    | 0.020/<br>0.059   | <0.001  | 0.091                                    | 0.065/<br>0.116   | <0.001  | -0.42   | -1.08/<br>0.24    | 0.216   | 1.57   | 1.20/<br>1.93   | <0.001  |
| BMI (ref. normal BMI)           |   |                   |         |  |                   |         |   |                   |         |  |                 |         |
| Underweight                     | -0.009                                  | -0.049/<br>0.030  | 0.614   | -0.026                                   | -0.079/<br>0.027  | 0.332   | -1.17   | -2.53/<br>0.18    | 0.089   | -0.80  | -1.52/<br>-0.08 | 0.029   |
| Overweight                      | -0.024                                  | -0.040/<br>-0.007 | 0.005   | -0.036                                   | -0.058/<br>-0.014 | 0.001   | 0.94  | 0.38/ 1.51        | 0.001   | -0.39  | -0.64/<br>-0.08 | 0.014   |
| Obese                           | -0.101                                  | -0.121/<br>-0.080 | <0.001  | -0.154                                   | -0.182/<br>-0.127 | <0.001  | 0.71  | -0.01/<br>1.42    | 0.052   | -1.55  | -1.94/<br>-1.16 | <0.001  |
| Moving-related pain             |   |                   |         |  |                   |         |   |                   |         |  |                 |         |
| Upper extremities(yes)          | -0.032                                  | -0.048/<br>-0.015 | <0.001  | -0.047                                   | -0.069/<br>-0.025 | <0.001  | -2.68   | -3.24/<br>-2.12   | <0.001  | -0.70  | -1.00/<br>-0.39 | <0.001  |
| Back/lower extremities(yes)     | -0.041                                  | -0.056/<br>-0.026 | <0.001  | -0.061                                   | -0.081/<br>-0.040 | <0.001  | -0.86   | -1.39/<br>-0.34   | 0.001   | -0.59  | -0.88/<br>-0.31 | <0.001  |
| Heart disease(yes)              | -0.058                                  | -0.078/<br>-0.038 | <0.001  | -0.076                                   | -0.102/<br>-0.049 | <0.001  | -0.20   | -0.89/<br>0.50    | 0.582   | -0.68  | -1.07/<br>-0.30 | <0.001  |
| Lung disease(yes)               | -0.024                                  | -0.046/<br>-0.002 | 0.029   | -0.037                                   | -0.065/<br>-0.008 | 0.012   | -0.34   | -1.09/<br>0.41    | 0.372   | -0.04  | -0.44/<br>0.37  | 0.853   |
| Depressive mood(yes)            | -0.056                                  | -0.077/<br>-0.036 | <0.001  | -0.070                                   | -0.097/<br>-0.043 | <0.001  | -1.29   | -2.01/<br>-0.57   | <0.001  | -1.02  | -1.41/<br>-0.63 | <0.001  |
| MMSE>24p(yes)                   | 0.069                                   | 0.046/<br>0.0092  | <0.001  | 0.107                                    | 0.076/<br>0.137   | <0.001  | 2.41  | 1.62/ 3.20        | <0.001  | 1.64   | 1.20/<br>2.08   | <0.001  |
| Internal HLoC                   | 0.006                                   | 0.004/<br>0.008   | <0.001  | 0.007                                    | 0.005/<br>0.010   | <0.001  | 0.08  | 0.01/ 0.15        | 0.019   | 0.08   | 0.05/<br>1.20   | <0.001  |
| Chance HLoC                     | -0.004                                  | -0.006/<br>-0.002 | <0.001  | -0.008                                   | -0.010/<br>-0.005 | <0.001  | 0.16  | -0.04/<br>0.07    | 0.562   | -0.08  | -0.11/<br>-0.05 | <0.001  |
| External HLoC                   | -0.003                                  | -0.005/<br>-0.002 | <0.001  | -0.006                                   | -0.008/<br>-0.004 | <0.001  | -0.26   | -0.33/<br>-0.20   | <0.001  | -0.10  | -0.13/<br>-0.06 | <0.001  |
| R <sup>2</sup>                  |   | 0.435             |         |  | 0.507             |         |   | 0.645             |         |  | 0.378           |         |
| Model fit                       | F(18, 3269) =141.7, p<0.001             |                   |         | F(18, 3255) =185.6, p<0.001              |                   |         | F(18, 3464) = 353.1, p<0.001                          |                   |         | F(18, 3425) = 117.3 p<0.001                      |                 |         |

Note: <sup>a</sup>Unstandardized regression coefficient

### Results

The study population comprised 3,819 participants. Their mean age was 69.4 years (SD=9.8), and the proportion of women was 54.8%. Moving-related pain from the upper extremities was reported by 32.2% and moving-related pain from the back/lower extremities by 49.5%. Heart disease, including infarction, angina pectoris or heart failure, was reported by 18.0% and lung disease, including asthma, COPD or tuberculosis, by 12.5%. According to the MADRS, 14.0% were categorized as suffering from depressive mood and 12.6% scored ≤24 points on the MMSE scale (Table 1).

Lower mean values in the WS and HGS tests were found in

participants with moving-related pain from the upper or back/lower extremities, heart disease, and lung disease as well as in those who were in a depressive state at the time of the study (Table 2).

All physical performances, walking 15m at normal speed, at maximum speed, HGS and AS, correlated positively with the results of the internal HLoC scale: walking 15m at normal speed  $r=0.12$ ,  $n=3,423$ ,  $p<0.001$ . The tested physical performances correlated negatively with the results of chance HLoC, walking 15m at normal speed  $r=-0.29$ ,  $n=3,425$ ,  $p<0.001$ , and external HLoC, walking 15m at normal speed  $r=-0.30$ ,  $n=3,410$ ,  $p<0.001$  (Table 2).

**Table 4:** Attrition analysis comparing study sample to non-participants.

|                        | Study sample, N=3819<br>n(%) | Non-participants, N=635<br>n(%) | p-value |
|------------------------|------------------------------|---------------------------------|---------|
| Sex                    |                              |                                 |         |
| Men                    | 1725(45.2)                   | 243(38.3)                       | 0.001   |
| Women                  | 2094(54.8)                   | 392(61.7)                       |         |
| Age, Mean(SD)          | 69.4(9.8)                    | 77.6(12.1)                      | <0.001  |
| Age decade             |                              |                                 |         |
| 60                     | 2335(61.7)                   | 224(35.3)                       | <0.001  |
| 70                     | 57(13.5)                     | 42(6.6)                         |         |
| 80                     | 817(21.4)                    | 218(34.3)                       |         |
| 90                     | 130(3.4)                     | 151(23.8)                       |         |
| Education              |                              |                                 |         |
| Primary school         | 1787(46.9)                   | 235(64.0)                       | <0.001  |
| Secondary school       | 1110(29.1)                   | 86(23.4)                        |         |
| University             | 911(23.9)                    | 46(12.5)                        |         |
| BMI                    |                              |                                 |         |
| Underweight            | 139(3.7)                     | 32(8.2)                         | <0.001  |
| Normal                 | 1302(34.8)                   | 146(37.3)                       |         |
| Overweight             | 1593(42.6)                   | 139(35.5)                       |         |
| Obese                  | 703(18.8)                    | 74(18.9)                        |         |
| Moving-related pain    |                              |                                 |         |
| Upper Extremities      |                              |                                 |         |
| Yes                    | 1231(32.3)                   | 129(27.8)                       | 0.052   |
| No                     | 2586(67.7)                   | 335(72.2)                       |         |
| Moving-related pain    |                              |                                 |         |
| Back/lower extremities |                              |                                 |         |
| Yes                    | 1891(49.5)                   | 240(51.7)                       | 0.375   |
| No                     | 1926(50.5)                   | 224(48.3)                       |         |
| Hart disease           |                              |                                 |         |
| Yes                    | 688(18.0)                    | 176(37.1)                       | <0.001  |
| No                     | 3126(82.0)                   | 299(62.9)                       |         |
| Lung disease           |                              |                                 |         |
| Yes                    | 476(12.5)                    | 50(10.6)                        | 0.233   |
| No                     | 3338(87.5)                   | 423(89.4)                       |         |
| Depressive mood        |                              |                                 |         |
| Yes                    | 533(14.4)                    | 88(28.8)                        | <0.001  |
| No                     | 3175(85.6)                   | 218(71.2)                       |         |
| MMSE                   |                              |                                 |         |
| ≤24p                   | 481(12.8)                    | 147(45.7)                       | <0.001  |
| >24p                   | 3279(87.2)                   | 175(54.3)                       |         |

The adjusted multivariate linear regression showed that a higher result on the internal HLoC scale was significantly associated with walking 15m at a normal speed ( $B=0.006$ ,  $p<0.001$ ) as well as at a maximum speed ( $B=0.007$ ,  $p<0.001$ ), while a higher result on the chance and external HLoC was significantly associated with a longer time to walk 15 metres at a normal speed ( $B=-0.004$ ,  $p<0.001$ ) and ( $B=-0.003$ ,  $p<0.001$ ) respectively, and walking 15m at a maximum speed ( $B=-0.008$ ,  $p<0.001$ ) and ( $B=-0.006$ ,  $p<0.001$ ). Furthermore, a higher result on the internal HLoC scale was significantly associated with a stronger hand grip ( $B=0.08$ ,  $p=0.019$ ), while a higher result on the external HLoC scale was significantly associated with a weaker hand grip ( $B=-0.26$ ,  $p<0.001$ ). In the alternating step test; a higher result on the internal HLoC scale was significantly associated with higher number of steps ( $B=0.08$ ,  $p<0.001$ ), while a higher result on the chance and external HLoC was significantly associated with a lower number of steps ( $B=-0.08$ ,  $p<0.001$ ) and ( $B=-0.10$ ,  $p<0.001$ ) respectively (Table 3).

### Discussion

This study showed that internal, chance and external HLoC were significantly associated with normal and maximum WS and HSG, while internal and external HLoC were significantly associated with AS after adjustment for sex, age, education, BMI, heart and lung disease, depressive mood and cognitive impairment. High scores on the internal LoC scale were associated with better performance, while high scores on the chance and external LoC scale was associated with poorer performance. By considering HLoC as a personal factor [1], the results from this study complement previous reports on the importance of individual traits for physical performance. For example, studies using the big five personality traits have reported lower HGS to be associated with higher neuroticism and lower openness [43], while higher walking speed was associated with greater conscientiousness, extraversion, openness and lower neuroticism [44-46].

Several temporary and more permanent physical, environmental and psychological causes in addition to HLoC can affect WS, HGS and AS. In particular, temporary causes can be a problem when the results of these tests, as well as other physical tests, are compared with standard values or, as indicated above, are included in the assessment of health status, frailty or functional ability. Moreover, in terms of LoC, research has shown that older adults are more likely than younger individuals to acknowledge that external LoC is more important [47]. In view of the age and health status of the participants in the present study, we tried as far as possible to adjust for this fact. However, based on the aim of this study, a further hypothesis is that individuals with a predominantly internal HLoC have greater self-confidence and are more at ease in the sometimes unfamiliar clinical setting, thus performing WS, HGS and AS tests better than those with an external HLoC. Previous studies have shown that individuals whose actions in relation to health issues are characterized by an internal HLoC are more likely to exercise than those with an external HLoC, who do not take their own initiative for health promotion activities to the same extent. Thus, a likely explanation for the better results among those with an internal locus would instead be that they are fitter. However, more fit or not, it is difficult to completely disregard HLoC and whether internal or chance/external LoC is the indirect or direct explanation for physical performance.

In view of the above, one question is whether it would be worth while to try to influence individuals' HLoC as a means of making them more likely to exercise with possible long-term positive health effects and, if so, how best to do it – by improving the influence of the internal locus or reducing the influence of the external locus, or both? Overall, considering the results of the present study and earlier reported health effects associated with internal and external HLoC in older adults, an intervention aimed at improving the internal LoC would probably be best [48,49]. At the same time, HLoC does not solely predict specific health behaviours, as it also depends on how different individuals value their health and the effect of exercise [50].

Another question with reference to the present study is whether a change in internal or external HLoC would have any decisive significance for an improved result in WS, HGS or AS. Overall, even if an individual with e.g., low internal HLoC could be encouraged to rethink or change her/his attitude, thereby achieving a higher total score on the internal scale, the changes in the physical tests reported here would be small. Nevertheless, even small improvements in the physical tests can be of foremost importance. The difference in walking speed between 70-80- and 93-year-olds is approximately 0.15m/s. Individuals with the opportunity to change from a low to a high internal HLoC could improve their walking speed to a rate that corresponds to the difference between the oldest age decades. Thus, it is difficult to completely ignore the importance of HLoC and its impact on the physical tests reported in this study.

### Limitations

There are limitations in this study. We included participants at two different time points, and although the risk is small, we cannot rule out that the result was altered by a cohort effect in some of the included variables. Another limitation could be a BMI misclassification. We used standing height when calculating the BMI, which can be questioned, as lower height in older adults is often due to physiological and pathological changes, which can lead to an overestimation of BMI [26].

The attrition analysis showed that the non-participants were older and that the proportion of women lower education, underweight, heart disease, depressed mood and scoring  $\leq 24$  points on the MMSE was greater compared to the participants. A selection bias cannot be ruled out and the generalizability of the results should be done with caution. However, since it looks like non-participants are a more fragile group, e.g., greater proportion with lowered mood and heart problems, it is not unlikely that in addition to performing worse in the physical tests, the proportion with internal HLoC would also be smaller [8,9]. Thus, this would rather emphasize the overall result of this study, that more of an internal HLoC is associated with better results on the physical tests.

### Strengths

Strength of this study is the large sample and that participants were randomized from a general population in the southern part of Sweden, representing both rural and urban areas. To reduce selection bias, home visits were made, and help was offered to those who had problems reading due to impaired vision, cognitive impairment or other disabilities that could make it difficult to answer the questionnaires. All examinations and interviews were conducted by personnel who were specially trained for the study.

### Conclusion

In a clinical setting, assessment of HLoC may contribute to the understanding of physical performance in terms of WS, HGS and AS among older adults aged 60-93 years. A sense of internal control over one's own health was associated with faster walking, a stronger hand grip and a higher number of alternate steps. The findings suggest that modification of control beliefs has the potential to improve the performance of these physical tests, which could be worth considering in a medical or physiotherapeutic assessment.

### Author Statements

#### Data and Code Availability

Data analysis was performed using SPSS for Windows, version 24.0 (IBM 211 Corporation, Armonk, NY, USA). The authors do not have permission to share data.

#### Conflict of Interest Statement

The authors declare no competing interests.

#### Authors' Contributions

Data collection: HE and SE; Study design: HE, LSW and SE; Data analysis: HE, LSW and SE; Writing the manuscript: HE, LSW and SE

#### Funding

The Good Aging in Skåne project, a part of the Swedish National Study on Aging and Care ([www.snac.org](http://www.snac.org)), is supported by the Swedish Ministry of Health and Social Affairs, Region Skåne, the Medical Faculty at Lund University, the Swedish Research Council (grant 2017-01613), Konung Gustaf V och Drottning Viktorias Frimurarestiftelse and Gyllenstiernska Krapperupstiftelsen.

### References

1. Rotter JB. Generalized expectancies for internal versus external control of reinforcement. *Psychol Monogr.* 1966; 80: 1-28.
2. Rotter JB, Chance JE, Phares EJ. *Applications of a Social Learning Theory and Personality.* New York: Holt, Rinehart & Winston. 1972.
3. Wallston BS, Wallston KA, Kaplan GD, Maides SA. Development and validation of the health locus of control (HLC) scale. *J Consult Clin Psychol.* 1976; 44: 580-5.
4. Hashimoto H, Fukuhara S. The influence of locus of control on preferences for information and decision making. *Patient Educ Couns.* 2004; 55: 236-40.
5. Kostka T, Jachimowicz V. Relationship of quality of life to dispositional optimism, health locus of control and self-efficacy in older subjects living in different environments. *Qual Life Res.* 2010; 19: 351-61.
6. Kaplan GD, Cowles A. Health locus of control and health value in the prediction of smoking reduction. *Health Educ Monogr.* 1978; 6: 129-37.
7. Berglund E, Lytsy P, Westerling R. The influence of locus of control on self-rated health in context of chronic disease: a structural equation modeling approach in a cross sectional study. *BMC Public Health.* 2014; 14: 492.

8. Milte CM, Luszcz MA, Ratcliffe J, Masters S, Crotty M. Influence of health locus of control on recovery of function in recently hospitalized frail older adults. *Geriatr Gerontol Int.* 2015; 15: 341-9.
9. Aflakseir AA, Mohammad-Abadi MS. The Role of Health Locus of Control in Predicting Depression Symptoms in a Sample of Iranian Older Adults with Chronic Diseases. *Iran J Psychiatry.* 2016; 11: 82-6.
10. Stürmer T, Hasselbach P, Amelang M. Personality, lifestyle, and risk of cardiovascular disease and cancer: follow-up of population based cohort. *BMJ.* 2006; 332: 1359.
11. Zuercher-Huerlimann E, Stewart JA, Egloff N, von Känel R, Studer M, et al. Internal health locus of control as a predictor of pain reduction in multidisciplinary inpatient treatment for chronic pain: a retrospective study. *J Pain Res.* 2019; 12: 2095-2099.
12. Omeje O, Nebo C. The influence of locus control on adherence to treatment regimen among hypertensive patients. *Patient Prefer Adherence.* 2011; 5: 141-8.
13. Rongen A, Robroek SJ, Burdorf A. The importance of internal health beliefs for employees' participation in health promotion programs. *Prev Med.* 2014; 67: 330-4.
14. Brincks AM, Feaster DJ, Burns MJ, Mitrani VB. The influence of health locus of control on the patient-provider relationship. *Psychol Health Med.* 2010; 15: 720-8.
15. Sorlie T, Sexton HC. Predictors of the process of coping in surgical patients. *Pers. Individ. Differ.* 2001; 30: 947-960.
16. Binotto MA, Lenardt MH, Rodríguez-Martínez MDC. Physical frailty and gait speed in community elderly: a systematic review. *Rev Esc Enferm USP.* 2018; 52: e03392.
17. Celis-Morales CA, Welsh P, Lyall DM, Steell L, Petermann F, et al. Associations of grip strength with cardiovascular, respiratory, and cancer outcomes and all cause mortality: prospective cohort study of half a million UK Biobank participants. *BMJ.* 2018; 361: k1651.
18. Middleton A, Fritz SL, Lusardi M. Walking speed: the functional vital sign. *J Aging Phys Act.* 2015; 23: 314-22.
19. Christman SK, Fish AF, Frid DJ, Smith BA, Bryant CX. Stepping as an exercise modality for improving fitness and function. *Appl Nurs Res.* 1998; 11: 49-54.
20. Mercer VS, Freburger JK, Chang SH, Purser JL. Step Test scores are related to measures of activity and participation in the first 6 months after stroke. *Phys Ther.* 2009; 89: 1061-71.
21. Ferrucci L, Bandinelli S, Benvenuti E, Di Iorio A, Macchi C, et al. Subsystems contributing to the decline in ability to walk: bridging the gap between epidemiology and geriatric practice in the INCHIANTI study. *J Am Geriatr Soc.* 2000; 48: 1618-25.
22. Pasma JH, Stijntjes M, Ou SS, Blauw GJ, Meskers CG, et al. Walking speed in elderly outpatients depends on the assessment method. *Age (Dordr).* 2014; 36: 9736.
23. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, et al. Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci.* 2001; 56: M146-56.
24. Lagergren M, Fratiglioni L, Hallberg IR, Berglund J, Elmståhl S, et al. A longitudinal study integrating population, care and social services data. The Swedish National study on Aging and Care (SNAC). *Aging Clin Exp Res.* 2004; 16: 158-68.
25. Ekström H, Elmståhl S. Pain and fractures are independently related to lower walking speed and grip strength: results from the population study "Good Ageing in Skåne". *Acta Orthop.* 2006; 77: 902-11.
26. Gavriilidou NN, Pihlsgård M, Elmståhl S. Anthropometric reference data for elderly Swedes and its disease-related pattern. *Eur J Clin Nutr.* 2015; 69: 1066-75.
27. Bhaskaran K, Dos-Santos-Silva I, Leon DA, Douglas IJ, Smeeth L. Association of BMI with overall and cause-specific mortality: a population-based cohort study of 3•6 million adults in the UK. *Lancet Diabetes Endocrinol.* 2018; 6: 944-953.
28. Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry.* 1979; 134: 382-9.
29. Mottram P, Wilson K, Copeland J. Validation of the Hamilton Depression Rating Scale and Montgomery and Asberg Rating Scales in terms of AGE-CAT depression cases. *Int J Geriatr Psychiatry.* 2000; 15: 1113-9.
30. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975; 12: 189-98.
31. Tombaugh TN, McIntyre NJ. The mini-mental state examination: a comprehensive review. *J Am Geriatr Soc.* 1992; 40: 922-35.
32. Wrangler LS, Elmståhl S, Ekström H. Physical performance in relation to birth cohort: A comparison of 60 year old Swedish men and women born twelve years apart. *Arch Gerontol Geriatr.* 2019; 82: 22-27.
33. Kim HJ, Park I, Lee HJ, Lee O. The reliability and validity of gait speed with different walking pace and distances against general health, physical function, and chronic disease in aged adults. *J Exerc Nutrition Biochem.* 2016; 20: 46-50.
34. Jarnlo GB, Nordell E. Reliability of the modified figure of eight—a balance performance test for elderly women. *Physiother Theory Pract.* 2009; 19: 35-43.
35. Hammer A, Lindmark B. Test-retest intra-rater reliability of grip force in patients with stroke. *J Rehabil Med.* 2003; 35: 189-94.
36. Bramell-Risberg E, Jarnlo GB, Elmståhl S. Slowing of alternating forearm movements is associated with cognitive impairment in community-dwelling older people. *Dement Geriatr Cogn Disord.* 2010; 29: 457-66.
37. Bramell-Risberg E, Jarnlo GB, Elmståhl S. Separate physical tests of lower extremities and postural control are associated with cognitive impairment. Results from the general population study Good Aging in Skåne (GÅS-SNAC). *Clin Interv Aging.* 2012; 7: 195-205.
38. Tiedemann A, Lord SR, Sherrington C. The development and validation of a brief performance-based fall risk assessment tool for use in primary care. *J Gerontol A Biol Sci Med Sci.* 2010; 65: 896-903.
39. Wallston BS, Wallston KA, Kaplan GD, Maides SA. Development and validation of the health locus of control (HLC) scale. *J Consult Clin Psychol.* 1976; 44: 580-5.
40. Tabachnick BG, Fidell LS. *Using multivariate statistics* 5th edn. Pearson. 2007.
41. Kim JH. Multicollinearity and misleading statistical results. *Korean J Anesthesiol.* 2019; 72: 558-569.
42. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* 2013; 310: 2191-4.
43. Mueller S, Wagner J, Drewelies J, Duezel S, Eibich P, et al. Personality development in old age relates to physical health and



- cognitive performance: Evidence from the Berlin Aging Study II. *J Res Per.* 2012; 65: 94-108.
44. Sutin AR, Stephan Y, Terracciano A. Facets of conscientiousness and objective markers of health status. *Psychol Health.* 2018; 33: 1100-1115.
45. Tolea MI, Costa PT Jr, Terracciano A, Ferrucci L, Faulkner K, et al. Associations of openness and conscientiousness with walking speed decline: findings from the Health, Aging, and Body Composition Study. *J Gerontol B Psychol Sci Soc Sci.* 2012; 67: 705-11.
46. Stephan Y, Sutin AR, Canada B, Terracciano A. Personality and Frailty: Evidence From Four Samples. *J Res Pers.* 2017; 66: 46-53.
47. Lachman ME. Locus of control in aging research: a case for multidimensional and domain-specific assessment. *Psychol Aging.* 1986; 1: 34-40.
48. Menec VH, Chipperfield JG. Remaining active in later life. The role of locus of control in seniors' leisure activity participation, health, and life satisfaction. *J Aging Health.* 1997; 9: 105-25.
49. Slenker SE, Price JH, O'Connell JK. Health locus of control of joggers and nonexercisers. *Percept Mot Skills.* 1985; 61: 323-8.
50. Cheng C, Cheung MW, Lo BC. Relationship of health locus of control with specific health behaviours and global health appraisal: a meta-analysis and effects of moderators. *Health Psychol Rev.* 2016; 10: 460-477.