

Review Article

The Role of GSK in Ischaemic Organ Injury

Sian Mitchell, Jessica Guo, Hailin Zhou and Daqing Ma*

Department of Surgery and Cancer, Imperial College London, UK

*Corresponding author: Daqing Ma, Department of Surgery and Cancer, Anaesthetics, Pain Medicine and Intensive Care, Imperial College London, Faculty of Medicine, Chelsea and Westminster Hospital, London, UK

Received: August 30, 2014; Accepted: September 27, 2014; Published: September 30, 2014

Abstract

Glycogen synthase kinase (GSK) 3 is a serine/threonine kinase involved in many signaling pathways with phosphorylation determining its activity. Phosphorylation of the tyrosine 216 residue in the kinase domain causes activation of GSK3 while phosphorylation of the amino terminal serine residue results in GSK inactivation. Hence the dual role of GSK in apoptosis is well documented. The dysregulation of GSK has been linked to many illnesses: cancer, diabetes as well as ischaemic organ injury to name a few. Ischaemic organ injury occurs when there is not enough oxygen to meet the metabolic demands of the tissue and the involvement of GSK in brain, heart, kidney, gut and lung ischaemic organ injury is discussed further here. While GSK is a pivotal multiswitch enzyme, it is becoming an ever increasingly interesting therapeutic target with several drugs undergoing clinical trials.

Keywords: Glycogen synthase kinase; Organ injury; Molecular mechanism; Cellular signaling

Introduction

The diseases associated with depriving organs of their blood supply such as myocardial infarction, stroke or perinatal hypoxic ischaemic encephalopathy are a main cause of death in Europe [1]. Tissue ischaemia can be caused in surgical procedures such as organ transplantation and it is of major concern [2].

When cells are deprived of oxygen, irreversible damage can occur. However, reperfusing the organ, which is restoring blood flow after a period of ischaemia, also induces damage, namely reperfused injury [2], with reperfusion injury causing greater damage than simply the ischaemia alone [3]. This can lead to MODS (multiple organ dysfunction syndrome), which is fatal. Hence it is important to understand the pathophysiology behind ischaemia and ischaemia reperfusion injuries. Ischaemia is described as “inadequate oxygen delivery to cover the metabolic demands” of the tissue [2]. Ischaemia is regarded as a proinflammatory state as there is an increase in inflammatory molecules such as cytokines with a reduction in the more protective molecules [4].

There are molecular changes that occur when a tissue goes through ischaemia. Adenosinetriphosphate (ATP) is not resynthesized, but there is an increased breakdown to ADP and AMP and hypoxanthine. This change in ATP levels initiates the apoptotic cascade that results in cell death [5]. Hypoxanthine, in normal physiological settings, is oxidized by xanthine dehydrogenase to xanthine. However, in this hypoxic setting, the hypoxanthine is converted to xanthine oxidase, the O form. This form of xanthine oxidase has been shown to produce reactive oxygen species when oxygen is present again (reperfusion) [6]. These reactive oxygen species damage the tissue through lipid peroxidation [7].

The reperfusion injury that follows ischaemia increases the damage already created by ischaemia [8]. It does not just encompass local damage with reactive oxygen species, but also leads to systemic changes [2] and can increase the damage already created by ischaemia.

It is characterised by tissue oedema and increased vasculature permeability to protein [3].

The endothelial cells that line blood vessels are particularly sensitive to changes in oxygen levels and hypoxia triggers processes that lead to endothelial dysfunction. Neutrophils adhering to the surface of the endothelium cells are an essential step [3] with adherence to the vessel importance for vessel injuries. They release enzymes such as elastase [3] and block the capillaries, resulting in “the no-flow phenomenon”, which further increases the effect of the hypoxia. The no flow phenomenon describes the reduced flow of blood to an organ following ischaemia.

It also decreases the production of positive agents such as nitric oxide, prostacyclin and vasodilators whilst increasing the production of others, such as endothelin1 [1], a powerful vasoconstrictor. Endothelial swelling follows these processes with reduced capillary perfusion [9]. As ischaemia and reperfusion injuries are of significance, it is imperative to understand enzymes that play a role in such injuries in multiple organs; GSK3 is one of these enzymes.

Molecular structure of GSK

The enzyme, Glycogen Synthase Kinase (GSK) is a proline directed serine/threonine kinase involved in many signaling pathways in mammals [10]. There are 2 isoforms of GSK: GSK3 α and GSK3 β , with molecular weights of 51kda and 47kda respectively [11]. The structure of these kinases is an N terminal lobe made up of β sheets and a C terminal lobe consisting of α helices [12] with the ATP region between both these lobes. They are highly homologous [10] with the two isoforms having a sequence identity of 95% in the catalytic domain and a serine residue, which are inhibitory residues [13] near the N terminus [14]. It is the phosphorylation of the N terminus that determines the GSK enzymatic activity [15] with ser21 present for GSK3 α and Ser9 for GSK3 β [16]. In the case of GSK3 β , ser9 acts as an inhibitor by blocking the binding of substrates to the catalytic site [13].

The activating residues of GSK3 α and GSK3 β are tyrosine 279

and tyrosine 216 respectively [13]. When Tyrosine 216 is not phosphorylated, it prevents substrates from entering the catalytic site but phosphorylation of tyr216 stops this [13]. When Tyrosine 216 is phosphorylated it allows substrates to enter the catalytic site activating GSK3 β . It has however been noted that the difference between the two is due to the glycine [17] rich extension at the N terminus of GSK3 α , with the N terminal end in GSK3 α 63 residues longer than the β isoform [18]. In addition to this, a variant of GSK3 β has been identified: GSK32.

GSK belongs to the mitogen activated protein kinase family members [17] with the sequence of GSK3 β 41% similar to the sequence of cyclin dependent kinase 2 (CDK2) [18,19]. However, the substrates of GSK need a Carboxyl- terminal priming phosphate at (n+4) to be phosphorylated [19]. Nis the site of phosphorylation demonstrating its substrate specificity. This primed phosphorylation has a 100-1000 times greater efficiency than phosphorylation that does not involve priming [18] although it is not strictly necessary [14].

Only recently has the binding site for the priming substrate on GSK3 β been located and it includes 3 amino acid residues: arginine 96, Arginine 180 and lysine 205. There is also a large presence of GSK3 β in the brain [20], which has a particular importance for ischaemic and reperfusion injuries.

Function of GSK

Initially, over 20 years ago in 1980, GSK was thought to only be used in the phosphorylation of glycogen synthase, inhibiting the final step of glycogen synthesis [17]. However, since then, the physiological function of GSK3 has been discovered from genetic analysis in drosophila [21,22], dictyostelium [23] and yeast [24,25]. It is now considered a ubiquitous multiswitch enzyme and is known to phosphorylate up to 50 substrates [26] including the transcription factors c- jun, c- myb and c- myc [25]. It also phosphorylates B catenin and the APC tumour suppressor protein [27] enabling it to modulate metabolism, gene expression [28] and play a pivotal role in oxidative stress induced neuronal apoptosis [29]. With a link to many cellular processes, the dysregulation of GSK is linked to many diseases such as Cancer [30], Alzheimer's disease [31] and Diabetes [14]. Even with similar structures, the two GSK isoforms have different functions. GSK3 β null mice have an embryonic lethal phenotype [32] as it is essential for the formation of the embryonic axes [33,34].

GSK3 β is known to influence the transcription factor, Nf κ B and it is believed that this is the reason for its involvement in many of the inflammatory processes [35]. GSK3 β is of particular **interest as it** can control blood glucose levels via insulin [18] and therefore it has a role in the development of insulin resistance and type 2 diabetes. Studies have shown that GSK level is elevated in the skeletal muscle of obese patients and patients suffering from type 2 diabetes [36]. Insulin activity causes GSK inhibition via the well-known phosphoinositide 3 kinase (P13-K) signal transduction pathway [15]. P13K activates PKB (Akt), causing phosphorylation of serine 9 and 21 on the respective GSK isoforms. This prevents GSK activity towards substrates such as glycogen synthase and leads to substrates dephosphorylation and activation [17].

Cell fates during embryonic development are another role of GSK and works via the Wnt growth factor signalling pathway [19].

Wnt is essential in cellular proliferation, differentiation, motility and polarity of cells [37]. In the absence of wnt, GSK phosphorylates β catenin, downstream of Wnt and causes its ubiquitination and degradation in cells [14]. Although the precise mechanisms by which Wnt inhibits GSK activity towards β catenin remains unclear [15], it has been shown to involve the "destruction complex" consisting of GSK3, Axin/conductin, β catenin and adenomatous polyposis coli (APC) [38]. This again alludes to the importance of GSK in cancer, here through APC, a tumour suppressor gene.

Role of GSK in cell death/organ injury

GSK3 β has been shown to have opposing actions on apoptosis [28] induced cell death. This discovery was made when hepatocyte apoptosis occurred in GSK3 β knockout mice during embryonic development [32] opposing previous discoveries where expression of GSK3 β caused apoptosis. This ability is due to GSK3 β promoting the intrinsic apoptotic signaling pathway and inhibiting the extrinsic apoptotic pathway [39]. Activation of GSK3 β is caused by phosphorylation of the tyrosine 216 residue in the kinase domain and inactivation is caused by phosphorylation of the amino terminal serine residue [29].

The Bcl-2 family comprises of both pro apoptotic and anti-apoptotic proteins such as bax and bid respectively [40]. The ratio of pro apoptotic to anti apoptotic Bcl-2 family members determines whether the intrinsic pathway is activated [41]. Many stimuli can activate the apoptotic pathway such as DNA damage, stress and hypoxic ischaemia [39]. The pro-apoptotic proteins undergo conformational changes as it migrates from the cytoplasm to the mitochondria [39]. This sequesters the opening of the Mitochondrial Permeability Transition Pore (MPTP) with loss of mitochondrial membrane potential and release of cytochrome C, a component of the electron transport chain [42] from the mitochondria to the cytosol [43]. Cytochrome C forms the apoptosome, which activates caspase 9, an initiator caspase and then subsequently activates caspase 3, the most important executioner caspase [42], resulting in the degradation of chromosomal DNA, ultimately leading to cell death.

The extrinsic pathway, involves death receptors of the tumour necrosis factor family, which all share a similar extracellular domain, the death domain. This then involves the formation of the death inducing signaling complex (DISC) [44], which contains Fas, a member of the tumour necrosis factor family. Disc then activates caspase 8, which activates caspase 3 and results in cell death [45]. As GSK3 plays an important role in apoptosis, it is important to analyse the role that GSK plays in ischaemic organ injury

Brain: GSK3 β localization has occurred in some areas of the human brain [46] and is present in the central nervous system [47]. The brain is dependent on continuous blood supply and interestingly, the ATP level in the brain completely diminishes within 4 minutes [48]. Stroke, a result of brain ischaemia, is a common occurrence.

There have been numerous studies demonstrating the involvement of GSK β in brain ischaemic injury. In transient global cerebral ischaemia, induced by bilateral common carotid artery occlusion for 5 minutes with hypotension, phospho - GSK3 β , ser 9 and phospho-Akt, ser 473 were increased in the CA1 hippocampal subregion of mice brains suggesting Akt/GSK3 β signaling mediates survival of

hippocampal neurons [49]. After hypoxic ischaemia in 7-day-old neonatal rats, levels of IGF-1, pAKT and pGSK3 were measured. It was found that IGF-1 caused the activation of Akt during recovery after HI and caused inactivation of GSK3 β by phosphorylation of serine 9. During reperfusion, there was a redistribution of GSK3 β from the cytosol to the nucleus [50].

Traumatic Brain Injury (TBI) is a neurological condition linked with depressive behavior [51] activating pathways that involve apoptotic or anti-apoptotic signaling. TBI caused increased phosphorylation of serine 9 GSK3 β , the inhibitory form [51]. With this in mind and prior knowledge of the involvement of GSK3 in apoptotic pathways, it makes GSK3 β an ideal therapeutic target.

Heart: Coronary heart disease is a huge worldwide burden, with ischaemic injury a cause of myocardial infarctions [52]. Studies have demonstrated that GSK3 β plays a role in cardioprotection with GSK3 β involvement in ischaemic preconditioning [53]. Inhibitors of GSK3 β added to an *in vivo* model before ischaemia or at the beginning of reperfusion were shown to decrease necrosis. Anaesthetized Sprague Dawley rats underwent 30 minutes of myocardial ischaemia and 6 hours of reperfusion with or without the addition of the GSK3 inhibitor TDZD-8. Addition of the inhibitor decreased MI size, neutrophil infiltration, suppressed NF- κ B and p38 MAPK [52]. It has been shown that the β adrenergic receptor causes apoptosis in cardiac myocytes *in vitro* and *in vivo* [54,55] with evidence that it increases GSK3 activity. Singh et al demonstrated that extracellular ubiquitin, used for protein turnover, possesses protective properties through the inactivation of GSK3 β /JNK and the mitochondrial pathways [54].

As further evidence that GSK3 β has an effect on heart ischaemic conditions, the treatment of cardiac H9c2 cells, rat cardiomyoblast cell line, with exogenous zinc chloride prevented cardiac reperfusion by increasing the phosphorylation at serine 9 GSK3 β [56]. These studies again demonstrate the pivotal action that inhibition of GSK3 β has, on organ survival following ischaemia.

Kidneys: Acute ischaemic renal failure commonly causes death. In transient renal artery occlusion in rats, *in vivo*, GSK3 β was demonstrated to be involved in kidney dysfunction after stress or ischaemia through Bax activation, apoptosis by disturbing the balance of pro and anti-apoptotic proteins and tubular cell injury [57]. The study demonstrated that GSK3 β phosphorylates Bax at the ser163 residue [57]. Renal ischaemia also increases reactive oxygen species, which has been linked to GSK3 β activation [58].

Vasileva et al showed that the inhibition of GSK3 β is important in the protective cascade after renal ischaemia through preconditioning with lithium and insulin. Phosphorylation of the serine 9 residue of GSK3 β increases Nonspecific Mitochondrial Permeability (NMP) activation, which decreases cell death [59]. Inhibition of GSK was analysed on a mouse model of endotoxaemic acute renal failure to look at cell death, renal dysfunction and TNF α induced cytotoxicity. It was shown to decrease lipopolysaccharide-induced production of TNF α , RANTES and NF κ B [57]. As ischaemia is one of the main causes of renal failure and renal failure contributes largely to mortality in the UK, discovering ways to protect the kidney are highly important.

Gut: Gut ischaemia and reperfusion injury are greatly influenced

by the transcription factor NF κ B, which is a substrate of GSK3 β (Wang et al.). Cuzzocrea et al demonstrated that the GSK3 β inhibitor, TDZD-8 weakened apoptosis, NF κ B expression, ICAM expression and neutrophil infiltration amongst other factors when applied to an SAO animal model. This model involves ischaemia and reperfusion to splanchnic organs. It was deduced from these results that inhibiting GSK3 β stops the interaction between neutrophils and endothelial cells at the early and late of neutrophil transmigration [60]. GSK3 β is also involved in intestinal epithelial cells wound healing response. Phosphorylation of serine 9 residue of GSK3 in IEC18 cell epithelial cell monolayers and inhibition of B catenin nuclear translocation was also found, suggesting the importance of GSK3 β in wound healing.

Lungs: GSK3 action is also noted in the lungs. The presence of the GSK3/ β catenin axis plays an important role in airway smooth muscle cell proliferation as well as GSK being involved in lung injury. Cuzzocrea et al, one of the studies showing the involvement with the lungs, demonstrated that mice treated with bleomycin developed significant lung injury. However, with administration of TDZD-8, a GSK3 β inhibitor, the lung injury sustained was reduced. With evidence that GSK plays an important role in the physiology of the damage caused to many organs following ischemia, this makes it an attractive therapeutic target.

GSK as therapeutic target

GSK is a multi-tasking kinase [14] involved in the necrotic and apoptotic processes of many tissues with involvement in several pathological conditions, so it has become a common pharmacological target with increasing research to inhibit its actions. Lithium, the first GSK3 β inhibitor discovered and sodium valproate, another drug shown to inhibit GSK3, are both important in the treatment of depression and bipolar disorders [61]. Lithium, which is also a commonly used mood stabilizer [62], has been shown to protect neurons against glutamate excitotoxicity [63] and studies have implied that it can be used in the treatment of acute brain injuries. Klein and Melton first discovered that lithium could inhibit GSK3 β [64] and this was later supported by a rat brain model [65], by phosphorylation of the serine residues on both the α and β isoforms [62]. Sodium Valproate, another mood stabilizer and anti-convulsant drug, inhibits GSK3 and prevents the HIV-1 damage caused to the nervous system [66].

Alzheimer's is another disorder that is influenced by GSK-3 [30]. Thiadiazolidinones (TDZD), non-ATP competitive inhibitors of GSK-3 can prevent hyperphosphorylation [67]. The neurofibrillary tangles that are characteristic of Alzheimer's pathology are composed of a highly phosphorylated form of the microtubule associated protein, TDZD [68]. Hence, thiadiazolidinones have been implicated in the treatment of Alzheimer's [67,69,70].

Some synthetic GSK inhibitors have been developed, with two inhibitors SB-216763 and SB-415286 shown to protect against neuronal cell death [71]. However, a reported complication with such inhibitors is the build-up of β -catenin, which is linked to the development of cancers, namely colon and melanomas [72].

Another GSK3 synthetic inhibitor, AR-A014418, was found to reduce the immobility time in rats that underwent a forced swim test, which is used to analyze antidepressant efficacy [65]. As well as being a target in brain disorders, inhibition of GSK maintains

the pluripotency of embryonic stem cells [73]. 6-bromindirubin-3-oxime (BIO), the specific inhibitor of GSK3 β , activates the wnt signaling pathway, and maintains the undifferentiated phenotypes of the ESC as well as increasing the specific transcription factors such as Oct3/4 and Nanog to maintain pluripotency [73].

Along with the inhibitors ability to affect neuronal cell death, SB-216763, SB-415286 and TDZD-8 were also able to decrease nuclear factor kappa B p65 activity and therefore decrease inflammation in organs [74]. GSK3b is also involved in the pathophysiology of diabetes and therefore its inhibition could help disease progression with the discovery of many GSK3b inhibitors for this purpose. CT20026 was shown to improve oral glucose tolerance in diabetic obese rhesus monkeys with longer treatment decreasing diabetic progression [75]. 3-aminopyrazole, the water-soluble inhibitor of GSK, administered to 18 hour fasted ob/ob mice was also shown to significantly lower blood glucose levels [74]. Pyrrole-2,5-diones has also been administered to ZDF rats and its ability as an oral active GSK3b demonstrated [76]. Although the majority of the studies have been conducted in rats, these are promising results. With ischaemic and reperfusion injuries a major worry in healthcare, such advances are ever more important.

Conclusion

GSK-3 has been shown to be hugely important in many signaling pathways and it is relevant in a multitude of disease pathophysiology. Ischaemia causes a great deal of damage to organs with the involvement of GSK in this process. The phosphorylation and inhibition of GSK prevents the opening of the mitochondrial permeability transition pore (mPTP) and is known to be protective in ischaemic reperfusion injuries [77]. With these properties, it has become an interesting therapeutic target with the use of many GSK3 inhibitors currently being trialed [78]. These trials are still in the early stages, but nevertheless are an important development in scientific research.

References

- Carden DL, Granger DN. Pathophysiology of ischaemia-reperfusion injury. *J Pathol.* 2000; 190: 255-266.
- Dorweiler B, Pruefer D, Andradi TB, Maksan SM, Schmiedt W, Neufang A, et al. Ischemia-reperfusion injury. *European Journal of Trauma and Emergency Surgery.* 2007; 33: 600-612.
- Welbourn CR, Goldman G, Paterson IS, Valeri CR, Shepro D, Hechtman HB. Pathophysiology of ischaemia reperfusion injury: central role of the neutrophil. *Br J Surg.* 1991; 78: 651-655.
- Sirotković-Skerlev M, Pleština S, Bilić I, Kovac Z. [Pathophysiology of ischaemia-reperfusion injury]. *Lijec Vjesn.* 2006; 128: 87-95.
- McCully JD, Wakiyama H, Hsieh YJ, Jones M, Levitsky S. Differential contribution of necrosis and apoptosis in myocardial ischemia-reperfusion injury. *American Journal of Physiology-Heart and Circulatory Physiology.* 2004; 286: H1923-H35.
- Kaminski KA, Bonda TA, Korecki J, Musial WJ. Oxidative stress and neutrophil activation—the two keystones of ischemia/reperfusion injury. *Int J Cardiol.* 2002; 86: 41-59.
- Toyokuni S. Reactive oxygen species-induced molecular damage and its application in pathology. *Pathol Int.* 1999; 49: 91-102.
- Eltzschig HK, Collard CD. Vascular ischaemia and reperfusion injury. *Br Med Bull.* 2004; 70: 71-86.
- Seal JB, Gewertz BL. Vascular dysfunction in ischemia-reperfusion injury. *Ann Vasc Surg.* 2005; 19: 572-584.
- Hernández F, Langa E, Cuadros R, Avila J, Villanueva N. Regulation of GSK3 isoforms by phosphatases PP1 and PP2A. *Mol Cell Biochem.* 2010; 344: 211-215.
- Das S, Wong R, Rajapakse N, Murphy E, Steenbergen C. Glycogen synthase kinase 3 inhibition slows mitochondrial adenine nucleotide transport and regulates voltage-dependent anion channel phosphorylation. *Circ Res.* 2008; 103: 983-991.
- Noble ME, Endicott JA, Johnson LN. Protein kinase inhibitors: insights into drug design from structure. *Science.* 2004; 303: 1800-1805.
- Meijer L, Flajolet M, Greengard P. Pharmacological inhibitors of glycogen synthase kinase 3. *Trends Pharmacol Sci.* 2004; 25: 471-480.
- Patel S, Doble B, Woodgett JR. Glycogen synthase kinase-3 in insulin and Wnt signalling: a double-edged sword? *Biochem Soc Trans.* 2004; 32: 803-808.
- Bax B, Carter PS, Lewis C, Guy AR, Bridges A, Tanner R, et al. The structure of phosphorylated GSK-3beta complexed with a peptide, FRATtide, that inhibits beta-catenin phosphorylation. *Structure.* 2001; 9: 1143-1152.
- Bertrand JA, Thieffine S, Vulpetti A, Cristiani C, Valsasina B, Knapp S, et al. Structural characterization of the GSK-3beta active site using selective and non-selective ATP-mimetic inhibitors. *J Mol Biol.* 2003; 333: 393-407.
- Cohen P, Frame S. The renaissance of GSK3. *Nat Rev Mol Cell Biol.* 2001; 2: 769-776.
- ter Haar E, Coll JT, Austen DA, Hsiao HM, Swenson L, Jain J. Structure of GSK3beta reveals a primed phosphorylation mechanism. *Nat Struct Biol.* 2001; 8: 593-596.
- Frame S, Cohen P, Biondi RM. A common phosphate binding site explains the unique substrate specificity of GSK3 and its inactivation by phosphorylation. *Molecular cell.* 2001; 7: 1321-1327.
- Woodgett JR. Molecular cloning and expression of glycogen synthase kinase-3/factor A. *EMBO J.* 1990; 9: 2431-2438.
- Siegfried E, Perkins LA, Capaci TM, Perrimon N. Putative protein kinase product of the Drosophila segment-polarity gene zeste-white3. *Nature.* 1990; 345: 825-829.
- Ruel L, Bourouis M, Heitzler P, Pantesco V, Simpson P. Drosophila shaggy kinase and rat glycogen synthase kinase-3 have conserved activities and act downstream of Notch. *Nature.* 1993; 362: 557-560.
- Harwood AJ, Plyte SE, Woodgett J, Strutt H, Kay RR. Glycogen synthase kinase 3 regulates cell fate in Dictyostelium. *Cell.* 1995; 80: 139-148.
- Puziss JW, Hardy TA, Johnson RB, Roach PJ, Hieter P. MDS, a dosage suppressor of an mck1 mutant, encodes a putative yeast homolog of glycogen synthase kinase 3. *Mol Cell Biol.* 1994; 14: 831-839.
- Plyte SE, Feoktistova A, Burke JD, Woodgett JR, Gould KL. Schizosaccharomyces pombe skp1+ encodes a protein kinase related to mammalian glycogen synthase kinase 3 and complements a cdc14 cytokinesis mutant. *Molecular and cellular biology.* 1996; 16: 179-191.
- Jope RS, Johnson GV. The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem Sci.* 2004; 29: 95-102.
- Rubinfeld B, Albert I, Porfiri E, Fiol C, Munemitsu S, Polakis P. Binding of GSK3beta to the APC-beta-catenin complex and regulation of complex assembly. *Science.* 1996; 272: 1023-1026.
- Grimes CA, Jope RS. The multifaceted roles of glycogen synthase kinase 3beta in cellular signaling. *Prog Neurobiol.* 2001; 65: 391-426.
- Petit-Paitel A, Brau F, Cazareth J, Chabry J. Involvement of cytosolic and mitochondrial GSK-3beta in mitochondrial dysfunction and neuronal cell death of MPTP/MPP-treated neurons. *PLoS One.* 2009; 4: e5491.
- Martinez A, Castro A, Dorronsoro I, Alonso M. Glycogen synthase kinase 3 (GSK-3) inhibitors as new promising drugs for diabetes, neurodegeneration, cancer, and inflammation. *Medicinal research reviews.* 2002; 22: 373-384.
- Lovestone S, Reynolds CH, Latimer D, Davis DR, Anderton BH, Gallo JM,

- et al. Alzheimer's disease-like phosphorylation of the microtubule-associated protein tau by glycogen synthase kinase-3 in transfected mammalian cells. *Current Biology*. 1994; 4: 1077-1086.
32. Hoeflich KP, Luo J, Rubie EA, Tsao MS, Jin O, Woodgett JR. Requirement for glycogen synthase kinase-3 β in cell survival and NF- κ B activation. *Nature*. 2000; 406: 86-90.
33. Dominguez I, Itoh K, Sokol SY. Role of glycogen synthase kinase 3 beta as a negative regulator of dorsoventral axis formation in *Xenopus* embryos. *Proc Natl Acad Sci U S A*. 1995; 92: 8498-8502.
34. Nasevicius A, Hyatt T, Kim H, Guttman J, Walsh E, Sumanas S, et al. Evidence for a frizzled-mediated wnt pathway required for zebrafish dorsal mesoderm formation. *Development*. 1998; 125: 4283-4292.
35. Ali A, Hoeflich KP, Woodgett JR. Glycogen synthase kinase-3: properties, functions, and regulation. *Chem Rev*. 2001; 101: 2527-2540.
36. Henriksen EJ, Kinnick TR, Teachey MK, O'Keefe MP, Ring D, Johnson KW, et al. Modulation of muscle insulin resistance by selective inhibition of GSK-3 in Zucker diabetic fatty rats. *Am J Physiol Endocrinol Metab*. 2003; 284: E892-900.
37. Miller JR. The Wnts. *Genome Biol*. 2002; 3: REVIEWS3001.
38. Kimelman D, Xu W. beta-catenin destruction complex: insights and questions from a structural perspective. *Oncogene*. 2006; 25: 7482-7491.
39. Beurel E, Jope RS. The paradoxical pro- and anti-apoptotic actions of GSK3 in the intrinsic and extrinsic apoptosis signaling pathways. *Prog Neurobiol*. 2006; 79: 173-189.
40. Elmore S. Apoptosis: a review of programmed cell death. *Toxicol Pathol*. 2007; 35: 495-516.
41. Brunet A, Datta SR, Greenberg ME. Transcription-dependent and -independent control of neuronal survival by the PI3K-Akt signaling pathway. *Curr Opin Neurobiol*. 2001; 11: 297-305.
42. Green DR, Reed JC. Mitochondria and apoptosis. *Science*. 1998; 281: 1309-1312.
43. Yang J, Liu X, Bhalla K, Kim CN, Ibrado AM, Cai J, et al. Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science*. 1997; 275: 1129-1132.
44. Yon JH, Daniel-Johnson J, Carter LB, Jevtovic-Todorovic V. Anesthesia induces neuronal cell death in the developing rat brain via the intrinsic and extrinsic apoptotic pathways. *Neuroscience*. 2005; 135: 815-827.
45. Chinnaiyan AM, O'Rourke K, Tewari M, Dixit VM. FADD, a novel death domain-containing protein, interacts with the death domain of Fas and initiates apoptosis. *Cell*. 1995; 81: 505-512.
46. Pei JJ, Tanaka T, Tung Y-C, Braak E, Iqbal K, Grundke-Iqbal I. Distribution, levels, and activity of glycogen synthase kinase-3 in the Alzheimer disease brain. *Journal of Neuropathology & Experimental Neurology*. 1997; 56: 70-78.
47. Leroy K, Brion JP. Developmental expression and localization of glycogen synthase kinase-3 β in rat brain. *J Chem Neuroanat*. 1999; 16: 279-293.
48. Krause GS, White BC, Aust SD, Nayini NR, Kumar K. Brain cell death following ischemia and reperfusion: a proposed biochemical sequence. *Crit Care Med*. 1988; 16: 714-726.
49. Endo H, Nito C, Kamada H, Nishi T, Chan PH. Activation of the Akt/GSK3 β signaling pathway mediates survival of vulnerable hippocampal neurons after transient global cerebral ischemia in rats. *Journal of Cerebral Blood Flow & Metabolism*. 2006; 26:1479-1489.
50. Brywe KG, Mallard C, Gustavsson M, Hedtj rn M, Leverin AL, Wang X, et al. IGF-I neuroprotection in the immature brain after hypoxia-ischemia, involvement of Akt and GSK3 β ? *Eur J Neurosci*. 2005; 21: 1489-1502.
51. Shapira M, Licht A, Milman A, Pick CG, Shohami E, Eldar-Finkelman H. Role of glycogen synthase kinase-3 β in early depressive behavior induced by mild traumatic brain injury. *Mol Cell Neurosci*. 2007; 34: 571-577.
52. Gao HK, Yin Z, Zhou N, Feng XY, Gao F, Wang HC. Glycogen synthase kinase 3 inhibition protects the heart from acute ischemia-reperfusion injury via inhibition of inflammation and apoptosis. *J Cardiovasc Pharmacol*. 2008; 52: 286-292.
53. Tong H, Imahashi K, Steenbergen C, Murphy E. Phosphorylation of glycogen synthase kinase-3 β during preconditioning through a phosphatidylinositol-3-kinase-dependent pathway is cardioprotective. *Circulation research*. 2002; 90: 377-379.
54. Singh M, Roginskaya M, Dalal S, Menon B, Kaverina E, Boluyt MO, et al. Extracellular ubiquitin inhibits β -AR-stimulated apoptosis in cardiac myocytes: role of GSK-3 β and mitochondrial pathways. *Cardiovascular research*. 2010; 86: 20-28.
55. Iwai-Kanai E, Hasegawa K, Araki M, Kakita T, Morimoto T, Sasayama S. alpha- and beta-adrenergic pathways differentially regulate cell type-specific apoptosis in rat cardiac myocytes. *Circulation*. 1999; 100: 305-311.
56. Chanoit G, Lee S, Xi J, Zhu M, McIntosh RA, Mueller RA, et al. Exogenous zinc protects cardiac cells from reperfusion injury by targeting mitochondrial permeability transition pore through inactivation of glycogen synthase kinase-3 β . *American Journal of Physiology-Heart and Circulatory Physiology*. 2008; 295: H1227-H33.
57. Wang Z, Havasi A, Gall J, Bonegio R, Li Z, Mao H, et al. GSK3 β promotes apoptosis after renal ischemic injury. *J Am Soc Nephrol*. 2010; 21: 284-294.
58. Plotnikov EY, Kazachenko AV, Vysokikh MY, Vasileva AK, Tcvirkun DV, Isaev NK, et al. The role of mitochondria in oxidative and nitrosative stress during ischemia/reperfusion in the rat kidney. *Kidney Int*. 2007; 72: 1493-1502.
59. Vasileva AK, Plotnikov EY, Kazachenko AV, Kirpatovsky VI, Zorov DB. Inhibition of GSK-3 β decreases the ischemia-induced death of renal cells. *Bull Exp Biol Med*. 2010; 149: 303-307.
60. Cuzzocrea S, Mazzon E, Esposito E, Mui  C, Abdelrahman M, Di Paola R, et al. Glycogen synthase kinase-3 β inhibition attenuates the development of ischaemia/reperfusion injury of the gut. *Intensive Care Med*. 2007; 33: 880-893.
61. Rowe MK, Wiest C, Chuang DM. GSK-3 is a viable potential target for therapeutic intervention in bipolar disorder. *Neurosci Biobehav Rev*. 2007; 31: 920-931.
62. Chuang DM, Wang Z, Chiu C-T. GSK-3 as a target for lithium-induced neuroprotection against excitotoxicity in neuronal cultures and animal models of ischemic stroke. *Frontiers in molecular neuroscience*. 2011; 4.
63. Nonaka S, Hough CJ, Chuang D-M. Chronic lithium treatment robustly protects neurons in the central nervous system against excitotoxicity by inhibiting N-methyl-D-aspartate receptor-mediated calcium influx. *Proceedings of the National Academy of Sciences*. 1998; 95: 2642-2647.
64. Klein PS, Melton DA. A molecular mechanism for the effect of lithium on development. *Proc Natl Acad Sci U S A*. 1996; 93: 8455-8459.
65. Gould TD, Einat H, Bhat R, Manji HK. AR-A014418, a selective GSK-3 inhibitor, produces antidepressant-like effects in the forced swim test. *Int J Neuropsychopharmacol*. 2004; 7: 387-390.
66. Dewhurst S, Maggirwar SB, Schifitto G, Gendelman HE, Gelbard HA. Glycogen synthase kinase 3 beta (GSK-3 beta) as a therapeutic target in neuroAIDS. *J Neuroimmune Pharmacol*. 2007; 2: 93-96.
67. Martinez A, Alonso M, Castro A, P rez C, Moreno FJ. First non-ATP competitive glycogen synthase kinase 3 beta (GSK-3 β) inhibitors: thiazolidinones (TDZD) as potential drugs for the treatment of Alzheimer's disease. *J Med Chem*. 2002; 45: 1292-1299.
68. Alonso M, Martinez A. GSK-3 inhibitors: discoveries and developments. *Curr Med Chem*. 2004; 11: 755-763.
69. Medina M, Avila J. Glycogen synthase kinase-3 (GSK-3) inhibitors for the treatment of Alzheimer's disease. *Curr Pharm Des*. 2010; 16: 2790-2798.
70. Seren  L, Coma M, Rodr guez M, S nchez-Ferrer P, S nchez MB, Gich I, et al. A novel GSK-3 β inhibitor reduces Alzheimer's pathology and rescues neuronal loss in vivo. *Neurobiol Dis*. 2009; 35: 359-367.

71. Cross DA, Culbert AA, Chalmers KA, Facci L, Skaper SD, Reith AD. Selective small-molecule inhibitors of glycogen synthase kinase-3 activity protect primary neurones from death. *J Neurochem.* 2001; 77: 94-102.
72. Peifer M, Polakis P. Wnt signaling in oncogenesis and embryogenesis--a look outside the nucleus. *Science.* 2000; 287: 1606-1609.
73. Sato N, Meijer L, Skaltsounis L, Greengard P, Brivanlou AH. Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3-specific inhibitor. *Nat Med.* 2004; 10: 55-63.
74. Martinez A. Preclinical efficacy on GSK-3 inhibitors: towards a future generation of powerful drugs. *Med Res Rev.* 2008; 28: 773-796.
75. Wagman AS, Johnson KW, Bussiere DE. Discovery and development of GSK3 inhibitors for the treatment of type 2 diabetes. *Curr Pharm Des.* 2004; 10: 1105-1137.
76. Engler TA, Henry JR, Malhotra S, Cunningham B, Furness K, Brozinick J, et al. Substituted 3-imidazo[2-a]pyridin-3-yl-4-(2,3,4-tetrahydro-[4]diazepino-[6,7,1-h]indol-7-yl)pyrrole-2,5-diones as highly selective and potent inhibitors of glycogen synthase kinase-3. *J Med Chem.* 2004; 47: 3934-3937.
77. Zhai P, Sciarretta S, Galeotti J, Volpe M, Sadoshima J. Differential roles of GSK-3 β during myocardial ischemia and ischemia/reperfusion. *Circ Res.* 2011; 109: 502-511.
78. Takahashi-Yanaga F, Sasaguri T. Drug development targeting the glycogen synthase kinase-3 β (GSK-3 β)-mediated signal transduction pathway: inhibitors of the Wnt/ β -catenin signaling pathway as novel anticancer drugs. *J Pharmacol Sci.* 2009; 109: 179-183.