

Case Report

Treat PDPH (Post Dural Puncture Headache) with Sphenopalatine Ganglion Block –Transnasal Approach!

Sandeep K¹, Sravya A², Rashmi NS^{2*} and Rishardhan P^{2*}

¹Department of Anesthesiology, Associate Professor- D.Y. Patil Medical College, Kolhapur, India

²Department of Anesthesiology, Junior Resident- D.Y. Patil Medical College, Kolhapur, India

*Corresponding author: Rashmi NS, Department of Anesthesiology, Junior Resident- D.Y. Patil Medical College, Kolhapur, India

Rishardhan P, Department of Anesthesiology, Junior Resident- D.Y. Patil Medical College, Kolhapur, India

Received: February 10, 2020; Accepted: March 05, 2020; Published: March 12, 2020

Presentation of Case

Case 1

A 23 year male, ASA 1 patient after undergoing spinal anesthesia for right tibia plating came with complain of dull throbbing headache frontal- occipital region 24 hours post-operatively. Headache aggravated on sitting position and relieved on supine position. Patient also complained of one episode of vomiting and mild neck stiffness but no other signs of meningitis were present. Patient had no complains of tinnitus, hypoacusia and photophobia. Investigation, complete blood counts were also within normal range. Clinical diagnosis of PDPH was made.

Management: After the diagnosis initially conservative management was undertaken. Patient was given complete bed rest in supine position for 24 hours and IV fluids, increase oral intake of water and caffeine and abdominal binders were advised. Patient was examined again after 24 hours, the patient complained of presence persistent pain corresponding to VAS score 8.

SPG Block: Under all aseptic precautions, SPG block was given to the patient bedside. Both the nostrils were initially sprayed with 10% lox spray and the middle turbinate were packed with gauge soaked in 4% lox.

The gauges were kept in place for 20 minutes with patient in supine position. After removal of pack VAS score when reassessed and was 2 in both supine and sitting position.

Case 2

23yr old female 2nd gravida, 36 weeks of pregnancy case of PIH on tab labetalol 100 mg BD, previous LSCS 2yrs back with fetal distress underwent emergency LSCS under spinal anesthesia with 25G quincke needle. She started with headache in occipital region with neck pain 3rd postoperative day, which aggravated on sitting position and relieved in supine. No complaints fever vomiting, her BP was under control.

Management: She was treated with bedrest, NSAIDS, IV fluids

Abstract

Background and Study Objective: To treat PDPH by simple and effective means using transnasal Sphenopalatine ganglion block.

PDPH (Post Dural Puncture Headache) is one of the very well known complications. Management of PDPH is always challenging to anesthesiologist. Though the incidence of PDPH has reduced with new small spinal needles and needle tip designs, still occasional PDPH cases are observed.

her pain did not relieve with persistent VAS score of 8-9.

So we decided to manage the case with SPG block, First 10% xylocaine was sprayed in each nostril with spray nozzle.

Later each nostril was packed with gauge soaked with 4% xylocaine pack was kept for 20 mins, VAS score was assessed immediately after removal of nasal pack was 1. VAS score was reassessed after 6, 12 and 24 hours vas score was 2, 4 and 6 respectively.

After 24 hours as the VAS was 6, we decided to reblock SPG. Xylocine 10% was sprayed deep inside each nostril aiming lateral wall of nose above middle turbinate. Patient was relieved of pain within 2 mins of spray hence we decided not to pack the nose with gauge soaked in xylocaine. Patient was reassessed at 2, 6, 12 and 24 hours and her VAS score were 2, 2, 0, 0 respectively.

Case 3

42 year female patient with bilateral ureteric calculi was operated for bilateral ureteroscopy and lithoclast under spinal anesthesia using 25 G quincke spinal needle. On 2nd post-operative day, she complained of headache in occipital and frontal region which aggravated on standing and sitting position with VAS score of 7-8.

Management: She was initially treated conservatively with IV fluids, bedrest and NSAIDS and was reassessed after 48 hours. VAS was persistently 8. She was later treated with SPG block by nasal spray of 10% xylocaine and nasal packing was done with 4% xylocaine. Pack was kept for 20mins. Conservative management was also continued. Patient was reassessed after 6, 12 and 24 hours, VAS was 3, 2, 1 respectively.

Supplementary Information

PDPH is a well-known distressing complication. PDPH occurs due to the continuous loss of Cerebrospinal Fluid (CSF), higher than its production rate at the point of approach of the Dura-mater and reflex vasodilatation of the meningeal vessels [1-7]. Vasodilation is mediated by the parasympathetic activity of neurons that have synapses in the sphenopalatine ganglion, this explains relief of PDPH

by SPGB [6]. There is also consequent traction on the meninges due to loss of CSF, which causes headache.

Various methods have been used as treatment modality of PDPH. Initial treatment modality considered is conservative, which includes flat bed rest, hydration, caffeine intake and drugs like sumatriptan and gabapentin, if still persistent then Epidural Autologous Blood Patch (EBP) is performed. EBP is considered gold standard for treatment of PDPH. The clot formed by the EBP seals the dural puncture and the rapid formation of CSF allows for rapid restoration of CSF volume and resolution of the PDPH. EBP is more invasive as compared to SPG block and has its own contraindication (include those that normally apply to epidurals, but include a raised white cell count, pyrexia and technical difficulties) and complications (Immediate exacerbation of symptoms and radicular pain have been described).

The Sphenopalatine Ganglion (SPG) is a parasympathetic ganglion, located in the pterygopalatine fossa and is posterior to middle nasal turbinate. The SPG block has been used for treating headaches of varying causes. Its is said to be parasympathetic ganglion as pre ganglionic parasympathetic axons synapse within the ganglion, but few sensory branches of maxillary nerve and post-ganglionic sympathetic neurons also pass through it.

SPG block has been used to treat various chronic pain syndromes such as cluster headache, CRPS, TMJ pain, post herpetic neuralgia etc.

SPG can be approached by lateral infratemporal approach, sub zygomatic approach, trans oral or trans nasal approach. Transnasal approach is a non-invasive and a simple technique, which can be performed at the bedside with patient in supine position. The patient is given supine position with extension at neck and cotton swab soaked in local anesthetic is applied at the middle turbinate with help of an applicator or bilateral nasal packing is done with gauge soaked in local anesthetic or using nasal spray of 10% xylocaine. Though the swab doesn't come in direct contact with SPG, but the local anesthetic infiltrates around it causing the block (Figure 1 and Figure 2).

Cohen et al first published the use of SPG block for management of PDPH in 2009 [7] Patel et.al performed a retrospective study on 72 patients. The patients were divided in two groups, one group receiving Epidural Blood Patch (EBP) and the other bilateral SPG block. He found that the group that received SPG block had better pain relief after 1 hour of block being given compared to the group that received EBP but the overall pain relief after 24 hours have been found to be similar [1]. Similar observation has been seen in our 3 cases (i.e) patient had immediate pain relief with in 1 hour and was comfortable for 24 hours.

Kent et al, used SPG in 3 cases of PDPH in obstetric patients with good results, in our study we had one obstetric patient.

Preksha dubey et al, used intranasal spray for SPG block for PDPH [7]. They utilized 10% xylocaine spray instead of cotton swab soaked with 2% or 4% xylocaine to block SPG.

In our study in all three patients we initially sprayed 10% xylocaine in each nostril and then nasal packing was done. In 2nd cases when after 24 hours patient had a VAS score of 6 we just sprayed 10% xylocaine which relieved her pain immediately and this time no




Figure 1: Anatomy of sphenopalatine ganglion.



Figure 2: Clinical images.

further nasal packing was done.

No EBP was needed in any of our cases hence, bilateral SPG block holds superior compared to EBP in being a noninvasive and simple procedure with lesser complications. It can be considered as a ideal 1st line management for treatment of PDPH in future. Our case studies were limited to 3 cases, we need to carry out a larger study to further validate our results.

References

1. Nair AS, Rayani BK. Sphenopalatine ganglion block for relieving postdural puncture headache: Technique and mechanism of action of block with a narrative review of e-cacy. *Korean J Pain*. 2017; 30: 93-97.
2. Patel P, Zhao R, Cohen S, Mellender S, Shah S, et al. Sphenopalatine Ganglion Block (SPGB) versus epidural blood patch (EBP) for accidental Postdural Puncture Headache (PDPH) in obstetric patients: A retrospective observation. Poster presentation at 32nd Annual Meeting of the American Academy of Pain Medicine. Palm Springs, CA. 2016.
3. Kent S, Mehaffey G. Transnasal sphenopalatine ganglion block for the treatment of postdural puncture headache in obstetric patients. *J Clin Anesth*. 2016; 34: 194-196.
4. Cohen S, Sakr A, Katyal S, Chopra D. Sphenopalatine ganglion block for postdural puncture headache. *Anaesthesia*. 2009; 64: 574-575.

5. Turnbull DK, Shepherd DB. Post-dural puncture headache: Pathogenesis, prevention and treatment. *Br J Anaesth.* 2003; 91: 718-729.
6. Piagkou M, Demesticha T, Troupis T, Vlasis K, Skandalakis P, Makri A et al. The pterygopalatine ganglion and its role in various pain syndromes: From anatomy to clinical practice. *Pain Pract.* 2012; 12: 399-412.
7. Preksha Dubey, Prakash K. Dubey. Intranasal lignocaine spray for sphenopalatine ganglion block for postdural puncture headache. *Saudi j. anaesthesia.* 2018; 12: 364-365.