

## Mini Review

# SMG-1 as a Promising Tumor Suppressor in the Management of Cancer

Nam SW\*

Hepatobiliary Unit, Division of Gastroenterology,  
Department of Internal Medicine, The Catholic University  
of Korea, Incheon St. Mary's Hospital, South Korea

\*Corresponding author: Soon Woo Nam,  
Hepatobiliary Unit, Division of Gastroenterology,  
Department of Internal Medicine, The Catholic University  
of Korea, Incheon St. Mary's Hospital, South Korea

**Received:** March 16, 2021; **Accepted:** April 16, 2021;

**Published:** April 23, 2021

## Introduction

SMG-1 (Suppressor of Morphogenesis in Genitalia-1) is an evolutionally conserved serine/threonine-protein kinase and belongs Phosphatidylinositol-3-Kinase (PI3K) related kinases (PIKKs) that include Ataxia Telangiectasia Mutated (ATM), ATM and Rad3 Related (ATR), Mammalian Target of Rapamycin (mTOR), DNA-Dependent Protein Kinase Catalytic Subunit (DNA-PKcs) and Transformation/Transcription Domain-Associated Protein (TRRAP) [1]. The deduced 3,521-amino acid protein has a calculated molecular mass of 410kD [2]. PIKKs have diverse functions. For example, ATM, ATR and DNA-PKcs are involved in the response to DNA damage. ATM and DNA-PKcs respond to DNA Double Strand Breaks (DSBs) and ATR to DNA replication blockers or formation of long stretches of single strand DNA [3]. mTOR is a nutrient-regulated kinase that controls a wide variety of pathways involved in metabolism and cell growth [4]. TRRAP functions as part of a multiprotein co-activator complex which is involved with the transcriptional activity of c-Myc and other transcriptional factors [5]. SMG-1 is a part of the mRNA surveillance complex that regulates Nonsense-Mediated mRNA Decay (NMD) [6]. SMG-1 was firstly reported as a member of the informational suppression in *Caenorhabditis elegans* (*C. elegans*) which affected several mRNA processes in 1989 [7]. Pulak et al. reported that loss of function mutations affecting seven *C. elegans* smg genes eliminates NMD [8] and later demonstrated that smg-1 kinase activity was essential for NMD [9]. In 2001, Deming et al. reported a partial sequence of human SMG-1 as *C. elegans* SMG-1 related protein [10]. Yamashita et al. also reported the full-length sequence for human smg-1 which encoded a 3657 aa protein that was a novel PIKK and showed the involvement of SMG1 in mammalian NMD [11]. In 2004, Brumbaugh et al. reported the activation of SMG-1 by DNA damage and involvement of SMG-1 in genotoxic stress-induced phosphorylation of P53 [2]. Besides NMD, SMG-1 roles as a protective agency in genotoxic stress such as radiation, tumor proliferation, and apoptosis. I will briefly summarize the roles of SMG-1 related with NMD and others in this paper.

## SMG-1 as a Player in NMD

NMD is a cellular defense mechanism eliminating mRNA that harbors Premature Translation Termination Codons (PTCs) and

encode nonfunctional or potentially harmful polypeptides [6,12,13]. NMD pathway involves a cascade of fine-coordinated events. A SURF complex (SMG-1, SMG-8, SMG-9, UPF1 and the eukaryotic release factors 1 and 3 (eRF1 and 3) is formed on a ribosome which encounters a PTC [14,15]. Interaction of UPF1 with UPF2 and UPF3B bound to the downstream EJC triggers UPF1 phosphorylation by the SMG-1 and remodels the SURF complex to form the decay-inducing complex [16-19]. The primary structure of PIKKs contains a conserved C terminus preceded by a long stretch of helical, mostly HEAT (huntington, elongation factor 3, a subunit of PP2A and TOR1) repeats which are units of two anti-parallel  $\alpha$ -helices connected by flexible loops and large superhelical frequently twisted structures [20]. A bent arm comprising a long region of HEAT repeats at the N-terminus of SMG-1 functions as a scaffold for SMG-8 and SMG-9, and projects from C-terminal core containing PI3K domain, which modulates NMD [21,22]. Resultant NMD processes decrease the production of potentially harmful polypeptides and enhances the accuracy of gene expression. SMG-1 is one of the essential players triggering an NMD response, because it is the kinase that phosphorylates the UPF1 protein. SMG-1 functions in NMD by selective degrading PTCs mRNAs which can be generated by gene mutations, splicing or transcription errors and by phosphorylating specific serine residues in UPF1 helicase which is a crucial regulator of NMD [23]. Phosphorylated UPF1 leads the releases of eRF1 and eRF3 and recruits NMD factors such as SMG-5~7 [24]. They recognize its phosphorylation by inducing the remodeling of the mRNA surveillance complex which discriminates the PTC-containing mRNA from the normal mRNA as an essential step in NMD [11,14]. Recently, González-Huici et al. declared that SMG-1 kinase activity could be activated following DNA damage to phosphorylate specific DNA repair proteins and/or NMD inactivation may lead to aberrant mRNAs leading to synthesis of malfunctioning DNA repair proteins [25]. Consequently, SMG-1 play viral roles in the regulation of cell growth, proliferation, survival, and cellular responses to stresses as a member of PIKKs.

## SMG-1 as a Player in Tumor Suppression

SMG-1 has been suggested as a tumor suppressor, though its role as an NMD effector has been precisely documented. In the planarians study, the planarian mTORC1 signaling compounds and SMG-1 performed as a key regulator of regeneration and growth antagonistically. Rapamycin as an inhibitor of mTORC1 increased the survival rate of smg-1 animals by decreasing cell proliferation. They showed the possibility of the tumor suppressor function of SMG-1 [26]. Gubanov et al. reported that in Human Papillomavirus (HPV)-positive head and neck squamous cell carcinoma, SMG-1 was underexpressed and exhibited tumor suppressive activity [27]. They also reported that SMG-1 suppresses CKD2 and tumor growth by regulating both the p53 and Cdc25A signaling pathways in 2013. They showed that SMG-1 regulates the G 1/S checkpoint




Figure 1: SMG-1 as a tumor suppressor.

through both a p53-dependent, and a p53-independent pathway and depletion of SMG-1 increased tumor growth [28]. In 2014, Nam et al. declared that SMG-1 as a promising modulator of sorafenib resistance by using an unbiased genome-wide massive genetic screen method. They showed that the inhibition of SMG-1 reduced sorafenib sensitivity in the several hepatocellular carcinoma cell lines and could be an agent to reverse sorafenib resistance [29]. The results suggested the possibility of SMG-1 as a potent tumor suppressor (Figure 1). At a similar time, Han et al. demonstrated that expression of SMG-1 was significantly lower in the HCC tissue than that in the normal tissues and declared that SMG-1 expression was an independent prognostic marker for overall survival [30]. A novel AKT inhibitor inhibited phosphorylation of AKT downstream molecules and activated phosphorylation of mTOR and SMG-1 in the several liver cancer cell lines [31]. This report suggested the promising possibility of usage as anti-cancer regimen of SMG-1. Recently, Zhang et al. reported that SMG-1 was suppressed by miR-192 and -215 and functioned as a tumor suppressor in gastric cancer by inactive Wnt signaling and suppressing epithelial-mesenchymal transition [32].

### Conclusion

SMG-1 is involved in NMD as an active effector and tumorigenesis as a tumor suppressor. Moreover, SMG-1 may be a promising interrupter during the cancer proliferation mechanisms for development of anti-cancer agent and represent a biomarker for predicting the prognosis of cancer. SMG-1 may warrant investigation to defeat cancer in the future. Therefore, more well-coordinated studies are needed to elucidate the overall precise mechanisms of SMG-1 involving NMD and cancer development processes.

### References

1. Baretic D and Williams RL. PIKKs-the solenoid nest where partners and kinases meet. *Curr Opin Struct Biol.* 2014; 29: 134-142.
2. Brumbaugh KM, Otterness DM, Geisen C, et al. The mRNA surveillance protein hSMG-1 functions in genotoxic stress response pathways in mammalian cells. *Mol Cell.* 2004; 14: 585-598.
3. Shiloh Y. ATM and related protein kinases: safeguarding genome integrity. *Nat Rev Cancer.* 2003; 3: 155-168.

4. Wullschlegel S, Loewith R and Hall MN. TOR signaling in growth and metabolism. *Cell.* 2006; 124: 471-484.
5. McMahon SB, Wood MA and Cole MD. The essential cofactor TRRAP recruits the histone acetyltransferase hGCN5 to c-Myc. *Mol Cell Biol.* 2000; 20: 556-562.
6. Yamashita A, Kashima I and Ohno S. The role of SMG-1 in nonsense-mediated mRNA decay. *Biochim Biophys Acta.* 2005; 1754: 305-315.
7. Hodgkin J, Papp A, Pulak R, Ambros V and Anderson P. A new kind of informational suppression in the nematode *Caenorhabditis elegans*. *Genetics.* 1989; 123: 301-313.
8. Pulak R and Anderson P. mRNA surveillance by the *Caenorhabditis elegans* smg genes. *Genes Dev.* 1993; 7: 1885-1897.
9. Grimson A, O'Connor S, Newman CL and Anderson P. SMG-1 is a phosphatidylinositol kinase-related protein kinase required for nonsense-mediated mRNA Decay in *Caenorhabditis elegans*. *Mol Cell Biol.* 2004; 24: 7483-7490.
10. Denning G, Jamieson L, Maquat LE, Thompson EA and Fields AP. Cloning of a novel phosphatidylinositol kinase-related kinase: characterization of the human SMG-1 RNA surveillance protein. *J Biol Chem.* 2001; 276: 22709-22714.
11. Yamashita A, Ohnishi T, Kashima I, Taya Y and Ohno S. Human SMG-1, a novel phosphatidylinositol 3-kinase-related protein kinase, associates with components of the mRNA surveillance complex and is involved in the regulation of nonsense-mediated mRNA decay. *Genes Dev.* 2001; 15: 2215-2228.
12. Holbrook JA, Neu-Yilik G, Hentze MW and Kulozik AE. Nonsense-mediated decay approaches the clinic. *Nat Genet.* 2004; 36: 801-808.
13. Maquat LE. Nonsense-mediated mRNA decay: splicing, translation and mRNP dynamics. *Nat Rev Mol Cell Biol.* 2004; 5: 89-99.
14. Kashima I, Yamashita A, Izumi N, et al. Binding of a novel SMG-1-Upf1-eRF1-eRF3 complex (SURF) to the exon junction complex triggers Upf1 phosphorylation and nonsense-mediated mRNA decay. *Genes Dev.* 2006; 20: 355-367.
15. Yamashita A, Izumi N, Kashima I, et al. SMG-8 and SMG-9, two novel subunits of the SMG-1 complex, regulate remodeling of the mRNA surveillance complex during nonsense-mediated mRNA decay. *Genes Dev.* 2009; 23: 1091-1105.
16. Lykke-Andersen J, Shu MD and Steitz JA. Human Upf proteins target an mRNA for nonsense-mediated decay when bound downstream of a termination codon. *Cell.* 2000; 103: 1121-1131.
17. Chamieh H, Ballut L, Bonneau F and Le Hir H. NMD factors UPF2 and UPF3 bridge UPF1 to the exon junction complex and stimulate its RNA helicase activity. *Nat Struct Mol Biol.* 2008; 15: 85-93.
18. Ivanov PV, Gehring NH, Kunz JB, Hentze MW and Kulozik AE. Interactions between UPF1, eRFs, PABP and the exon junction complex suggest an integrated model for mammalian NMD pathways. *EMBO J.* 2008; 27: 736-747.
19. Melero R, Buchwald G, Castano R, et al. The cryo-EM structure of the UPF-EJC complex shows UPF1 poised toward the RNA 3' end. *Nat Struct Mol Biol.* 2012; 19: 498-505, S1-2.
20. Lempiainen H and Halazonetis TD. Emerging common themes in regulation of PIKKs and PI3Ks. *EMBO J.* 2009; 28: 3067-3073.
21. Arias-Palomo E, Yamashita A, Fernandez IS, et al. The nonsense-mediated mRNA decay SMG-1 kinase is regulated by large-scale conformational changes controlled by SMG-8. *Genes Dev.* 2011; 25: 153-164.
22. Deniaud A, Karuppasamy M, Bock T, et al. A network of SMG-8, SMG-9 and SMG-1 C-terminal insertion domain regulates UPF1 substrate recruitment and phosphorylation. *Nucleic Acids Res.* 2015; 43: 7600-7611.
23. Schweingruber C, Rufener SC, Zund D, Yamashita A and Muhlemann O. Nonsense-mediated mRNA decay - mechanisms of substrate mRNA recognition and degradation in mammalian cells. *Biochim Biophys Acta.*

- 2013; 1829: 612-623.
24. Okada-Katsuhata Y, Yamashita A, Kutsuzawa K, Izumi N, Hirahara F and Ohno S. N- and C- terminal Upf1 phosphorylations create binding platforms for SMG-6 and SMG-5: SMG-7 during NMD. *Nucleic Acids Res.* 2012; 40: 1251-1266.
  25. Gonzalez-Huici V, Wang B and Gartner A. A Role for the Nonsense-Mediated mRNA Decay Pathway in Maintaining Genome Stability in *Caenorhabditis elegans*. *Genetics.* 2017; 206: 1853-1864.
  26. Gonzalez-Estevez C, Felix DA, Smith MD, et al. SMG-1 and mTORC1 act antagonistically to regulate response to injury and growth in planarians. *PLoS Genet.* 2012; 8: e1002619.
  27. Gubanova E, Brown B, Ivanov SV, et al. Downregulation of SMG-1 in HPV-positive head and neck squamous cell carcinoma due to promoter hypermethylation correlates with improved survival. *Clin Cancer Res.* 2012; 18: 1257-1267.
  28. Gubanova E, Issaeva N, Gokturk C, Djureinovic T and Helleday T. SMG-1 suppresses CDK2 and tumor growth by regulating both the p53 and Cdc25A signaling pathways. *Cell Cycle.* 2013; 12: 3770-3780.
  29. Nam SW, Park KC, Yang KJ, Lee B and Kim SW. Genetic screen identifies suppressor of morphogenesis in genitalia-1 (SMG-1) as a modulator of sorafenib resistance in hepatocellular carcinoma cell lines. *Int J Oncol.* 2014; 45: 1450-1456.
  30. Han LL, Nan HC, Tian T, et al. Expression and significance of the novel tumor-suppressor gene SMG-1 in hepatocellular carcinoma. *Oncol Rep.* 2014; 31: 2569-2578.
  31. Zhang Y, Zheng Y, Faheem A, et al. A novel AKT inhibitor, AZD5363, inhibits phosphorylation of AKT downstream molecules, and activates phosphorylation of mTOR and SMG-1 dependent on the liver cancer cell type. *Oncol Lett.* 2016; 11: 1685-1692.
  32. Zhang X, Peng Y, Huang Y, et al. SMG-1 inhibition by miR-192/-215 causes epithelial-mesenchymal transition in gastric carcinogenesis *via* activation of Wnt signaling. *Cancer Med.* 2018; 7: 146-156.