

## Clinical Image

# Osteopetrosis Revealed by Pseudarthrosis of a Neglected Subtrochanteric Fracture in an Adult: Case Report

Yassine Ben Bouzid\*; Marouane Dinia; Rida-Allah Bassir; Monsef Boufettal; Jalal Mekkaoui; Mohamed Kharmaz; Moulay Omar Lamrani; Mohamed Saleh Ber-rada

Department of Orthopaedic and Trauma Surgery, Ibn Sina University Hospital, Rabat, Morocco

\*Corresponding author: Yassine Ben Bouzid

Department of Orthopaedic and Trauma Surgery, Ibn Sina University Hospital, Rabat, Morocco.  
Email: yassine.benbouzid2@gmail.com

Received: March 21, 2023

Accepted: April 27, 2023

Published: May 04, 2023

## Introduction

First described in 1904, Osteopetrosis (OP) is a rare genetic disorder [1] affecting osteoclast activity resulting in decreased bone resorption and bone densification [2]. There are three types of osteopetrosis depending on the world of transmission: autosomal dominant OP, autosomal recessive OP, and intermediate autosomal OP [3]. Indeed, osteopetrosis is characterized by a hard but fragile bone with a narrow or absent medullary canal [4]. Consequently, the bone is so dense and fragile that it is subject to an increased risk of fractures and complications during surgery, which is why patients with OP are treated conservatively [5]. Furthermore, due to poor blood supply, patients with osteopetrosis have a high risk of infection and osteomyelitis, in addition to the risk of pseudarthrosis [6].

## Clinical Image

A 23-year-old patient consulted for chronic intermittent pain with limited mobility of the left hip and shortening of the left lower limb. On history, the patient reported a fall on the pelvis 2 years earlier, which caused acute pain in the left hip that later, became chronic and intermittent. The patient had not initially consulted. There was no history of anemia, fractures or recurrent infection. However, the patient reported a family history of bone disease without genetic evidence. The physical examination found a left lower limb shortened by 2cm; the mobility of the left hip was painless but limited, in particular abduction

which was limited to 10°. We did not notice any pain on palpation or mobilization. Walking was possible but with a limp secondary to the shortening of the limb. The systemic examination did not reveal any other abnormalities. Biologically, there was no anemia, phosphocalcic abnormality, or alteration of renal function. Radiographically, there was a displaced left subtrochanteric fracture with an absent medullary canal and diffuse densification of the bone. The patient refused to undergo genetic testing. The diagnosis of late osteopetrosis with pseudarthrosis of a neglected subtrochanteric fracture was made. The patient underwent surgical management with open reduction and locked plate osteosynthesis.



Figure 1:

## Discussion

Osteopetrosis is a rare hereditary disease that affects the process of bone remodeling by affecting the function of osteoclasts, resulting in decreased bone resorption and increased mineral density leading to hard bones and loss of the medullary canal [7-10]. Many patients with osteopetrosis are asymptomatic and diagnosed incidentally. This disease can also be revealed after a fracture [11]. In autosomal dominant OP, a fracture can occur in any bone but with a preponderance of the hip and trochanter region [12-16]. Fractures are often of simple transverse character following minor trauma [17].

The management of fractures in patients with osteopetrosis has been discussed in small case series [14]. The current literature suggests conservative management of some long bone fractures [18]. However, intertrochanteric and subtrochanteric fractures are at risk of pseudarthrosis and coxa vara [14,18,19]. These fractures should be treated surgically [18]. Several implants have been used in the reported case series, such as locked anatomic plates [11]. Surgical intervention requires wide availability of material. Moreover, during the osteosynthesis, a recurrent change of drill bits and cooling of the bone during drilling is necessary to avoid thermal bone necrosis [15]. Bhargava et al [15] report the occurrence of a subcapital fracture during tapping for dynamic hip screw fixation. In the case of pseudarthrosis, the absence of cancellous bone during OP limits access to an autologous graft, which constitutes a particular difficulty.

## Author Statements

### Ethics Approval and Consent to Participate

Ethical approval was not sought. Written consent was obtained from the patients.

### Availability of Data and Materials

The datasets used and analysed during the study are available from the corresponding author.

### Declaration of Conflicting Interest

The authors declare that there is no conflict of interest.

### Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

### Authors Contributions

YB has conceived and developed the technique. YB has collected the data. All authors have read and approved the final manuscript.

## References

1. Stark Z, Savarirayan R. Osteopetrosis. *Orphanet J Rare Dis.* 2009; 4: 5.
2. Pang Q, Chi Y, Zhao Z, Xing X, Li M, et al. Novel mutations of CLCN7 cause autosomal dominant osteopetrosis type II (ADO-II) and intermediate autosomal recessive osteopetrosis (IARO) in Chinese patients. *Osteoporos Int.* 2016; 27: 1047–1055.
3. Wu CC, Econs MJ, DiMeglio LA, Insogna KL, Levine MA, et al. Diagnosis and Management of Osteopetrosis: consensus Guidelines From the Osteopetrosis Working Group. *J Clin Endocrinol Metab.* 2017; 102: 3111–3123.
4. Paccou J, Michou L, Kolta S, Debiais F, Cortet B, et al. High bone mass in adults. *Joint Bone Spine.* 2018; 85: 693–699.
5. Hua X, Liu Z, Wang X. Subtrochanteric Femoral Fracture in a Patient with Osteopetrosis: Treated with Internal Fixation and Complicated by Intraoperative Femoral Neck Fracture. *Int J Gen Med.* 2020; 13: 1551-1557.
6. Agrawal A, Maniar A, Maniar R. Bilateral Girdlestone Procedure as a Treatment Modality in a Case of Complicated Adult-Onset Osteopetrosis: A Case Report. *JBJS Case Connect.* 2019; 9: e0179.
7. AT Aydın, AB Yeter, Osteopetrosis. *Türkiye Klinikleri Journal of Orthopaedics & Traumatology.* 2010; 3: 82–86.
8. J Nakayama, H Fujioka, M Kurosaka, H Kitazawa, N Maesawa, et al. Surgery for clavicular and humeral fractures in an osteopetrotic patient: a case report. *Journal of Orthopaedic Surgery.* 2007; 15: 251–254.
9. RK Gupta. Long bone fractures in osteopetrosis: awareness of primary pathology and appropriate pre-operative planning necessary to avoid pitfalls in fixation. *Injury Extra.* 2005; 36: 37–41.
10. Marks Jr SC. Congenital osteopetrotic mutations as probes of the origin, structure and function of osteoclasts. *Clin Orthop Relat Res.* 1984; 189: 239–63.
11. Aslan A, Baykal YB, Uysal E, Atay T, Kirdemir V, et al. Surgical treatment of osteopetrosis-related femoral fractures: two case reports and literature review. *Case Rep Orthop.* 2014; 2014: 891963.
12. Z Stark, R Savarirayan. Osteopetrosis. *Orphanet Journal of Rare Diseases.* 2009; 4: 5.
13. LL Ihde, DM Forrester, CJ Gottsegen, Masih S, Patel DB, et al. Sclerosing bone dysplasias: review and differentiation from other causes of osteosclerosis. *Radiographics.* 2011; 31: 1865–1882.
14. P Birmingham, KA Mchale. Case reports: treatment of subtrochanteric and ipsilateral femoral neck fractures in an adult with osteopetrosis. *Clinical Orthopaedics and Related Research.* 2008; 466: 2002–2008.
15. A Bhargava, M Vagela, CME Lennox. Challenges in the management of fractures in osteopetrosis! Review of literature and technical tips learned from long-term management of seven patients. *Injury.* 2009; 40: 1167–1171.
16. Rafiq A Kapoor, DJC Burton, JF Haines. A new modality of treatment for non-united fracture of the humerus in a patient with osteopetrosis: a case report. *Journal of Medical Case Reports.* 2009; 3: 15.
17. Dahl N, Holmgren G, Holmberg S, Ersmark H. Fracture patterns in malignant osteopetrosis (Albers-Schonberg disease). *Arch Orthop Trauma Surg.* 1992; 111: 121–3.
18. Armstrong DG, Newfield HJT, Gillespie R. Orthopaedic management of osteopetrosis: results of a survey and review of the literature. *J Pediatr Orthop.* 1999; 19: 122–32.
19. H Bombacı, I Esenkaya, M Gorgec, S Kus. Femoral neck fractures in osteopetrosis. *Eklemler Hastalıkları ve Cerrahisi.* 1998; 9: 59–62.