

Mini Review

Lactic Acid Bacteria: Their Applications in Foods

Bintsis T*

Department of Agricultural Technology, TEI of West Macedonia, Kozani, Greece

***Corresponding author:** Bintsis T, Department of Agricultural Technology, TEI of West Macedonia, Kozani, Greece

Received: February 27, 2018; **Accepted:** March 29, 2018; **Published:** April 05, 2018

Abstract

Lactic acid bacteria (LAB) are a heterogeneous group of bacteria which plays a significant role in a variety of fermentation processes. They ferment food carbohydrates and produce lactic acid as the main product of fermentation.

The main application of LAB is as starter cultures, with an enormous variety of fermented dairy, meat, fish, fruit, vegetable and cereal products. Besides, LAB contribute to the flavor, texture and nutritional value of the fermented foods, and such applications as adjunct, bio-protective, probiotic cultures and various applications in food industry are discussed.

Keywords: Lactic Acid Bacteria; Applications; Fermented Foods

Introduction

Lactic acid bacteria (LAB) play an important role in food, agricultural, and clinical applications. The general description of the bacteria included in the group is gram-positive, nonsporing, nonrespiring cocci or rods, which produce lactic acid as the major end product during the fermentation of carbohydrates [1]. The common agreement is that there is a core group consisting of four genera; *Lactobacillus*, *Leuconostoc*, *Pediococcus* and *Streptococcus*. Recent taxonomic revisions have proposed several new genera and the remaining group now comprises the following: *Aerococcus*, *Alloiococcus*, *Carnobacterium*, *Dolosigranulum*, *Enterococcus*, *Globicatella*, *Lactococcus*, *Oenococcus*, *Tetragenococcus*, *Vagococcus*, and *Weissella* [2]. Their importance is associated mainly with their safe metabolic activity while growing in foods utilizing available sugar for the production of organic acids and other metabolites. Their common occurrence in foods along with their long-lived uses contributes to their natural acceptance as GRAS (Generally Recognized as Safe) for human consumption [3]. The EFSA's 'Panel on Biological Hazards (BIOHAZ)' has concluded that the fermenting bacteria associated with food, whether resistant to antibiotics or not with the possible exception of enterococci do not pose a clinical problem [4]. However, they can act as a reservoir for transferable resistance genes. Strains with genes transferable in such a way could enter the food chain and increase the probability of a transfer to food associated intestinal pathogenic organisms.

The three main pathways which are involved in the manufacture and development of flavour in fermented food products are 1) glycolysis (fermentation of sugars), 2) lipolysis (degradation of fat) and 3) proteolysis (degradation of proteins) [1,5-9]. Lactate is the main product generated from the metabolism of carbohydrates and a fraction of the intermediate pyruvate can alternatively be converted to diacetyl, acetoin, acetaldehyde or acetic acid (some of which can be important for typical yogurt flavours). The contribution of LAB to lipolysis is relatively little, but proteolysis is the key biochemical pathway for the development of flavour in fermented foods [10,11]. Degradation of such components can be further converted to various alcohols, aldehydes, acids, esters and sulphur compounds for specific flavour development in fermented food products [10,11].

The genetics of the LAB have been reviewed [12-18] and complete genome sequences of a great number of LAB have been published [19] since 2001, when the first genome of LAB (*Lactococcus lactis* ssp. *lactis* IL1403) was sequenced and published [20].

Applications of LAB

Starter cultures for fermented foods

Fermented foods are produced through fermentation of certain sugars by LAB and the origins of them are lost in antiquity. The most commonly LAB used, nowadays, as starter cultures in food fermentations are shown in Table 1. For a detailed classification of starter cultures see [21,41,42].

Adjunct cultures

Secondary cultures, or adjunct cultures or adjuncts, are defined as any cultures that are deliberately added at some point of the manufacture of fermented foods, but whose primary role is not acid production. Adjunct cultures are used in cheese manufacture to give back to the cheese, some of the biodiversity removed by pasteurisation, improved hygiene and the addition of defined-strain starter culture [22,43]. These are mainly non-starter LAB which they can have a significant impact on flavor and accelerate the maturation process [22-43].

Extracellular polysaccharides (EPSs) are produced by a variety of bacteria and are present as capsular polysaccharides bound to the cell surface, or are released into the growth medium [44]. These polymers play a major role in the production of yogurt, cheese, fermented cream and milk-based desserts [45] where they contribute to texture, mouth-feel, taste perception and stability of the final products. In addition, it has been suggested that these EPSs or fermented milks containing these EPSs are active as prebiotics [46], cholesterol-lowering [47] and immunomodulators [48]. EPS-producing strains of *Streptococcus thermophilus* and *Lactobacillus delbrueckii* ssp. *bulgaricus* have been shown to enhance the texture and viscosity of yogurt and to reduce syneresis [49].

For the production of wine, LAB are involved in the malolactic fermentation, that is a secondary fermentation, which involves the conversion of L-malate to L-lactate and CO₂ via malate decarboxylase, also known as the malolactic enzyme, resulting in a reduction of wine

Table 1: Lactic acid bacteria used as starter cultures in the production of some fermented food products.

Product	Genera of LAB ¹	References
	Dairy products	
Cheese (Mesophilic starter)	<i>Lc. lactis</i> ssp. <i>lactis</i>	[22]
	<i>Lc. lactis</i> ssp. <i>cremoris</i>	
	<i>Lc. lactis</i> ssp. <i>lactis</i> var. <i>diacetylactis</i>	
	<i>Leuc. mesenteroides</i> ssp. <i>cremoris</i>	
Cheese (Thermophilic starter)	<i>S. thermophilus</i>	[22]
	<i>Lb. delbrueckii</i> ssp. <i>bulgaricus</i>	
	<i>Lb. helveticus</i>	
	<i>Lb. delbrueckii</i> ssp. <i>lactis</i>	
Cheese (Mixed starter)	<i>Lc. lactis</i> ssp. <i>lactis</i>	[22]
	<i>Lc. lactis</i> ssp. <i>cremoris</i>	
	<i>S. thermophilus</i>	
Yogurt	<i>Lb. delbrueckii</i> ssp. <i>bulgaricus</i> , <i>S. thermophilus</i>	[22]
Fermented milks	<i>Lb. delbrueckii</i> ssp. <i>bulgaricus</i> , <i>S. thermophilus</i> , <i>Lb. casei</i> , <i>Lb. acidophilus</i> , <i>Lb. rhamnosus</i> , <i>Lb. johnsonii</i>	[22]
Yakult	<i>Lb. casei</i> ssp. <i>casei</i>	[2]
Acidophilus milk	<i>Lb. Acidophilus</i>	[22]
Butter and buttermilk	<i>Lc. lactis</i> ssp. <i>lactis</i> , <i>Lc. lactis</i> ssp. <i>lactis</i> var. <i>diacetylactis</i> , <i>Lc. lactis</i> ssp. <i>cremoris</i> , <i>Leuc. mesenteroides</i> ssp. <i>cremoris</i>	[22]
Kefir	<i>Lb. kefir</i> , <i>Lb. kefiranofacies</i> , <i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. paracasei</i> ssp. <i>paracasei</i> , <i>Lc. lactis</i> ssp. <i>lactis</i> , <i>Leuc. mesenteroides</i>	[23]
Trahanas	<i>Lc. lactis</i> ssp. <i>lactis</i> , <i>Lc. lactis</i> ssp. <i>lactis</i> var. <i>diacetylactis</i> , <i>Leuc. mesenteroides</i> ssp. <i>cremoris</i> , <i>Lb. delbrueckii</i> ssp. <i>lactis</i> , <i>Lb. casei</i> , <i>Lb. delbrueckii</i> ssp. <i>bulgaricus</i> and <i>Lb. Acidophilus</i>	[24]
Fermented meat products		
Dry sausages	<i>Lb. sakei</i> , <i>Lb. curvatus</i> , <i>Lb. plantarum</i> , <i>Lb. pentosus</i> , <i>Lb. casei</i> , <i>P. pentosaceus</i> , <i>P. acidilactici</i>	[25,26]
Salami Milano	<i>Lb. sakei</i> , <i>Lb. plantarum</i>	[26]
Salame Piacentino	<i>Lb. acidophilus</i> , <i>Lb. helveticus</i> , <i>Lb. sakei</i> , <i>Lb. antri</i> , <i>Lb. oris</i> , <i>Lb. vaginalis</i> , <i>Lb. brevis</i> , <i>Lb. panis</i> , <i>Lb. versmoldensis</i> , <i>Lb. zeae</i> , <i>Lb. curvatus</i> , <i>Lb. paralimentarius</i> , <i>Lb. frumenti</i> , <i>Lb. plantarum</i> , <i>Lb. graminis</i> , <i>Lb. reuteri</i>	[27]
Greek dry fermented sausages	<i>Lb. sakei</i> , <i>Lb. plantarum</i> , <i>Lb. curvatus</i> , <i>Lb. pentosus</i> , <i>Lc. lactis</i> ssp. <i>lactis</i> , <i>W. hellenica</i> , <i>W. paramesenteroides</i> , <i>W. viridescens</i> , <i>W. minor</i>	[27]
Chorizo	<i>Lb. brevis</i> , <i>Lb. curvatus</i> , <i>Lb. sakei</i> , <i>Lc. lactis</i> , <i>P. acidilactici</i> , <i>P. pentosaceus</i> , <i>Leuc. mesenteroides</i>	[27]
Fermented fish products		
Thai fish	<i>Lb. plantarum</i> , <i>Lb. reuteri</i>	[28]
Pickled fruits and vegetables		
Cabbage (Sauerkraut)	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. fermentum</i>	[29]
Cucumber	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. pentosus</i> , <i>Lb. acidophilus</i> , <i>Lb. fermentum</i> , <i>Leuc. Mesenteroides</i>	[30,31]
Olives	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. pentosus</i>	[32,33]
Fermented cereal products		
Sourdough	<i>Lb. brevis</i> , <i>Lb. hilgardii</i>	[34,35]
	<i>Lb. sanfransicensis</i> , <i>Lb. farciminis</i> , <i>Lb. fermentum</i> , <i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. amylovorus</i> , <i>Lb. reuteri</i> , <i>Lb. pontis</i> , <i>Lb. panis</i> , <i>Lb. alimentarius</i> , <i>W. cibaria</i>	
Kimchi	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i> , <i>W. kimchii</i> sp. nov., <i>Lb. kimchi</i> , <i>Lb. sakei</i> , <i>W. koreensis</i>	[36,37,38]
Bushera	<i>Lb. plantarum</i> , <i>Lb. paracasei</i> ssp. <i>paracasei</i> , <i>Lb. fermentum</i> , <i>Lb. brevis</i> , <i>Lb. delbrueckii</i> ssp. <i>delbrueckii</i> , <i>S. thermophilus</i>	[39]
Pozol	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i> , <i>Lb. confusus</i> , <i>Lc. lactis</i> , <i>Lc. raffinolactis</i>	[40]

¹*Lc. Lactococcus*, *Lb. Lactobacillus*, *Leuc. Leuconostoc*, *P. Pediococcus*, *S. Streptococcus*, *W. Weissella*

acidity, providing microbiological stabilization and modifications of wine aroma [50,51].

Bio-protective cultures

Certain LAB has been found to produce bacteriocins, namely, polypeptides synthesized ribosomally by bacteria that can have a bacteriocidal or bacteriostatic effect on other bacteria [52,53].

In general, bacteriocins lead to cell death by inhibiting cell wall biosynthesis or by disrupting the membrane through pore formation [54]. Bacteriocins are therefore important in food fermentations where they can prevent food spoilage or the inhibition of food pathogens. The best known bacteriocin is nisin, which has gained widespread application in the food industry and is used as a food additive in at least 50 countries, particularly in processed cheese, dairy products

and canned foods [55]. Examples of useful bacteriocins produced by LAB are lactacin 3147 [56-59] from lactococci, macedovicin from *Streptococcus macedonicus* ACA-DC 198 [60,61], reuterin from *Lactobacillus reuteri* [62], sakacin M from *Lactobacillus sake* 148 [63] curvacin A, curvaticin L442 and lactocin AL705 from *Lactobacillus curvatus* LTH1174 [64], pediocin PA-1/AcH from *Pediococcus acidilactici* [65], plantaricins (A, EF and JK) from *Lactobacillus plantarum* [65]. The above bacteriocins have proved effective in many food systems for the control of food spoilage or pathogenic bacteria.

Antifungal activities of LAB have been reported [66-68]. In addition, LAB strains also have the ability to reduce fungal mycotoxins, either by producing anti-mycotoxinogenic metabolites, or by absorbing them [68].

For LAB to be used as bio-protective starter cultures, they must possess a range of physical and biochemical characteristics, and most importantly, the ability to achieve growth and sufficient production of antimicrobial metabolites, which must be demonstrated in the specific food environment.

Probiotic cultures

LAB is considered as a major group of probiotic bacteria [69,70]; probiotic has been defined by Fuller [71] as “a live microbial feed supplement which beneficially affects the host animal by improving its intestinal microbial balance”. Salminen et al. [72] proposed that probiotics are microbial cell preparations or components of microbial cells that have a beneficial effect on the health and well-being of the host. *Lactobacillus acidophilus*, *Lactobacillus casei*, *Lb. reuteri*, *Lactobacillus rhamnosus* and *Lb. plantarum* are the most used LAB in functional foods containing probiotics [73,74]. In addition, LAB, as part of gut microbiota ferments various substrates such as biogenic amines and allergenic compounds into short-chain fatty acids and other organic acids and gases [75].

Conclusion

LABS are the most commonly used microorganisms for the fermentation and preservation of foods. Bacterial cultures with specific traits have been developed during the last 17 years, since the discovery of the complete genome sequence of *Lc. lactis* ssp. *lactis* IL1403 and a variety of commercial starter, functional, bio-protective and probiotic cultures with desirable properties have marketed. Advances in the genetics, molecular biology, physiology, and biochemistry of LAB have provided new insights and applications for these bacteria. Food industry is now capable of producing standardized, safe and nutritious products with different flavours, sometimes with special health-promoting properties.

References

- Hayek SA, Ibrahim SA. Current Limitations and Challenges with Lactic Acid Bacteria: A Review. *Food and Nutrition Sciences*. 2013; 4: 73-87.
- Khalid K. An overview of lactic acid bacteria. *Intern J Biosci*. 2011; 1: 1-13.
- Bourdichon F, Berger B, Casaregola S, et al. A Safety Assessment of Microbial Food Cultures with History of Use in Fermented Dairy Products. *Bullet IDF*. 2012; 455: 2-12.
- EFSA. Scientific opinion of the panel on biological hazards on the request from EFSA on the maintenance of the list of QPS microorganisms intentionally added to food or feed. *EFSA J*. 2008; 928: 1-48.
- Wedajo B. Lactic Acid Bacteria: Benefits, Selection Criteria and Probiotic Potential in Fermented Food. *J Prob Health*. 2015; 3: 129.
- Grattepanche F, Miescher-Schwenninger S, Meile L, et al. Recent developments in cheese cultures with protective and probiotic functionalities. *Dairy Sci Technol*. 2008; 88: 421-444.
- Law BA. Cheese ripening and cheese flavour technology. In: Law BA (ed), *Technology of Cheese making*, Sheffield Academic Press Ltd., Sheffield. UK. 2010.
- Smit G, Smit BA, Engels WJ. Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. *FEMS Microbiology Reviews*. 2005; 29: 591-610.
- Tamime AY, Robinson RK. (eds) *Yoghurt Science and Technology*. 2nd Ed, Woodhead Publishing Ltd. Cambridge. 1999.
- Ammor MS, Mayo B. Selection criteria for lactic acid bacteria to be used as functional starter cultures in dry sausage production: An update. *Meat Science*. 2007; 76: 138-146.
- Souza MJ, Ardo Y, Mc Sweeney PLH. Advances in the study of proteolysis in cheese. *Int Dairy J*. 2001; 11: 327-345.
- Broadbent JR, McMahon, DJ, Welker DL, et al. Biochemistry, genetics, and applications of exopolysaccharide production in *Streptococcus thermophilus*: A review. *J Dairy Sci*. 2003; 86: 407-423.
- Callanan MJ, Ross RP. Starter Cultures: Genetics. In: Fox PF, McSweeney PLH, Cogan TM, et al. *Cheese: Chemistry, Physics and Microbiology*, Vol 1, 4th Ed, Elsevier Academic Press. London. UK. 2004; 149-161.
- Klaenhammer T, Altermann E, Arigoni F, et al. Discovering lactic acid bacteria by genomics. *Antonie van Leeuwenhoek*. 2002; 82: 29-58.
- Morelli L, Vogensen FK, von Wright A. Genetics of Lactic Acid Bacteria. In: Salminen S, von Wright A, Ouwehand A, *Lactic Acid Bacteria - Microbiological and Functional Aspects*, 3rd Ed, Marcel Dekker Inc., New York. 2011; 249-293.
- Mills S, O'Sullivan O, Hill C, et al. The changing face of dairy starter culture research: From genomics to economics. *Intern J Dairy Tech*. 2010; 63: 149-170.
- Castillo Martinez FA, Balciunas EM, Salgado JM, et al. Lactic acid properties, applications and production: A review. *Trends Food Sci & Tech*. 2013; 30: 70-83.
- Zhu Y, Zhang Y, Li Y. Understanding the industrial application potential of lactic acid bacteria through genomics. *Appl Microbiol Biotechnol*. 2009; 83: 597-610.
- NCBI Data collected from Genbank. Accessed. 2018
- Bolotin A, Wincker P, Mauger S, et al. The complete genome sequence of the lactic acid bacterium *Lactococcus lactis* ssp. *lactis* IL1403. *Genome Res*. 2001; 11: 731-753.
- Bintsis T, Athanasoulas A. Dairy Starter Cultures. In: Papademas P (ed). *Dairy Microbiology, a Practical Approach*, CRC Press. Boca Raton. FL. 2015; 114-154.
- Rattanachaiyapong P, Phumkhachorn P. Lactic acid bacteria: Their antimicrobial compounds and their uses in food production. *Annals of Biol Res*. 2010; 1: 218-228.
- Kivanç M, Yapıcı E. Kefir as a Probiotic Dairy Beverage: Determination Lactic Acid Bacteria and Yeast. *Intern J Food Engin*. 2015; 1: 55-60.
- Lazos ES, Aggelousis G, Bratakos M. The fermentation of trahanas: A milk-wheat flour combination. *Plant Foods Human Nutr*. 1993; 44: 45-62.
- Hammes WP, Hertel C. New developments in meat starter cultures. *Meat Sci*. 1998; 49: 25-38.
- Hugas M, Monfort JM. Bacterial starter cultures for meat fermentation. *Food Chem*. 1997; 59: 547-554.
- Aquilanti L, Garofalo C, Osimani A, et al. Ecology of lactic acid bacteria and coagulase negative cocci in fermented dry sausages manufactured in Italy and other Mediterranean countries: An overview. *Int Food Res J*. 2016; 23:

- 429-445.
28. Saithong P, Panthavee W, Boonyaratanakornkit M et al. Use of a starter culture of lactic acid bacteria in plaasom, a Thai fermented fish. *J Biosc Bioengin.* 2010; 110: 553-557.
29. Pundir RK, Jain P. Change in Microflora of Sauerkraut during Fermentation and Storage. *World J Dairy & Food Sci.* 2010; 5: 221-225.
30. Wakil SM, Lada SA, Fasika SA. Isolation and identification of antimicrobial-producing lactic acid bacteria from fermented cucumber. *African J Biotech.* 2014; 13: 2556-2564.
31. Nilchian Z, Sharifan A, Rahimi E, et al. Improvement of Fermented Cucumber Characteristics by Starter Culture of *Lactobacillus plantarum*, *L. bulgaricus* and *S. thermophiles*. *J Food Biosc Techn.* 2016; 6: 31-40.
32. Con AH, Karasli N. Determination of Antagonistic Starter Cultures for Pickle and Olive Fermentation Processes. *Czech J Food Sci.* 2009; 27: 185-193.
33. Hurtado A, Reguant C, Bordons A, et al. Lactic acid bacteria from fermented table olives. *Food Microb.* 2012; 31: 1-8.
34. Vollmor A, Meuser F. Influence of Starter Cultures Consisting of Lactic Acid Bacteria and Yeasts on the Performance of a Continuous Sourdough Fermenter. *Cereal Chem.* 1992; 69: 20-27
35. Palla M, Cristani C, Giovanneti M et al. Identification and characterization of lactic acid bacteria and yeasts of PDO Tuscan bread sourdough by culture dependent and independent methods. *Intern J Food Microb.* 2017; 250: 19-26.
36. Mheen TI, Kwon TW. Effect of temperature and salt concentration on kimchi fermentation. *Kor J Food Sci Technol.* 1984; 16: 443-450.
37. Choi HJ, Cheigh CI, Kim SB, et al. *Weissella kimchii* sp. nov., a novel lactic acid bacterium from kimchi. *Intern J of System and Evolut Microb.* 2002; 52: 507-511.
38. Lee KB, Kim HJ, Lee EJ. Mixed cultures of Kimchi lactic acid bacteria show increased cell density and lactate productivity. *African J Biotech.* 2013; 12: 4000-4005.
39. Muyenja CMBK, Narvhus JA, Treimo J, et al. Isolation, characterisation and identification of lactic acid bacteria from bushera: A ugandan traditional fermented beverage. *Intern J Food Microb.* 2003; 80: 201-210.
40. Nuraida L, Wacher MC, Owens JD. Microbiology of pozol, a Mexican fermented maize dough. *World J Microb & Biotech.* 1995; 11: 567-571.
41. Tamime AY. Microbiology of Starter Cultures. In: Robinson RK (ed). *Dairy Microbiology Handbook*, 3rd Ed, John Wiley & Sons Inc. New York. 2002; 261-366.
42. Parente E, Cogan TM. Starter Cultures: General Aspects. In: Fox PF, McSweeney PLH, Cogan TM et al. (eds), *Cheese: Chemistry, Physics and Microbiology*, Vol 1, 4th Ed, Elsevier Academic Press, London, UK. 2004: 123-147.
43. Cogan TM, Beresford TP, Steele J, et al. Invited review: Advances in starter cultures and cultured foods. *J Dairy Scien.* 2007; 90: 4005-4021.
44. Jolly L, Vincent SJ, Duboc P, et al. Exploiting exopolysaccharides from lactic acid bacteria. *Antonie van Leeuwenhoek.* 2002; 82: 367-374.
45. Gibson GR, Robertfroid MB. Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutr.* 1995; 125: 1401-1412.
46. Nakajima H, Suzuki Y, Kaizu H et al. Cholesterol-lowering activity of ropy fermented milk. *J Food Scien.* 1992; 57: 1327-1329.
47. Hosono A, Lee J, Ametani A, et al. Characterization of a water-soluble polysaccharide fraction with immunopotentiating activity from *Bifidobacterium adolescentis* M101-4. *Biosc Biotechn Biochem.* 1997; 61: 312-316.
48. Hassan AN, Ipsen R, Janzen T, et al. Microstructure and rheology of yogurt made with cultures differing only in their ability to produce exopolysaccharides. *J Dairy Scien.* 2003; 86: 1632-1638.
49. Doleyres Y, Schaub L, Lacroix C. Comparison of the functionality of exopolysaccharides produced in situ or added as bioingredients on yogurt properties. *J Dairy Scien.* 2005; 88: 4146-4156.
50. Liu SQ. A review: Malolactic fermentation in wine beyond deacidification. *J. Appl. Microbiol.* 2002; 92: 589-601.
51. Cappelo MS, Zapparoli G, Logrieco A, et al. Linking wine lactic acid bacteria diversity with wine aroma and flavour. *Int J Food Microb.* 2017; 243: 16-27.
52. McAuliffe O, Ross RP, Hill C. Lantibiotics: Structure, biosynthesis and mode of action. *FEMS Microbiol Rev.* 2001; 25: 285-308.
53. Ross RP, Morgan S, Hill C. Preservation and fermentation: Past, present and future. *Intern J Food Microb.* 2002; 79: 3-16.
54. Twomey D, Ross RP, Ryan M, et al. Lantibiotics produced by lactic acid bacteria: Structure, function and applications. *Antonie Van Leeuwenhoek.* 2002; 82: 165-185.
55. Delves-Broughton J, Blackburn P, Evans RJ, et al. Applications of the bacteriocin, nisin. *Antonie Van Leeuwenhoek.* 1996; 69: 193-202.
56. Ryan MP, Rea MC, Hill C, et al. An application in cheddar cheese manufacture for a strain of *Lactococcus lactis* producing a novel broad-spectrum bacteriocin, lacticin 3147. *Applied Environ Microb.* 1996; 62: 612-619.
57. Coakley M, Fitzgerald G, Ross RP. Application and evaluation of the phage resistance- and bacteriocin encoding plasmid pMRC01 for the improvement of dairy starter cultures. *Applied Environ Microb.* 1997; 63: 1434-1440.
58. Morgan SM, Galvin M, Ross RP, et al. Evaluation of a spray-dried lacticin 3147 powder for the control of *Listeria monocytogenes* and *Bacillus cereus* in a range of food systems. *Letters Applied Microb.* 2001; 33: 387-391.
59. O'Sullivan L, O'Connor EB, Ross RP, et al. Evaluation of live-culture-producing lacticin 3147 as a treatment for the control of *Listeria monocytogenes* on the surface of smear-ripened cheese. *J Applied Microb.* 2006; 100: 135-143.
60. Georgalaki MD, Van Den Berghe E, Kritikos D, et al. Macedocin, a food-grade lantibiotic produced by *Streptococcus macedonicus* ACA-DC 198. *Appl Environ Microbiol.* 2002; 68: 5891-5903.
61. Georgalaki MD, Papadimitriou K, Anastasiou R, et al. Macedovicin, the second food-grade lantibiotic produced by *Streptococcus macedonicus* ACA-DC 198. *Food Microb.* 2013; 33: 124-130.
62. Rodríguez E, Arqués JL, Rodríguez R, et al. Reuterin production by lactobacilli isolated from pig faeces and evaluation of probiotic traits. *Lett Appl Microbiol.* 2003; 37: 259-263.
63. Parada JL, Caron CR, Medeiros ABP. Bacteriocins from Lactic Acid Bacteria: Purification, Properties and use as Biopreservatives. *Brazilian Arch Biology Techn.* 2007; 50: 521-542.
64. Castellano P, Vignolo G. Inhibition of *Listeria innocua* and *Brochothrix thermosphacta* in vacuum-packaged meat by addition of bacteriocinogenic *Lactobacillus curvatus* CRL705 and its bacteriocins. *Letters in Applied Microb.* 2006; 43: 194-199.
65. Castellano P, Belfiore C, Fadda S, et al. A review of bacteriocinogenic lactic acid bacteria used as bioprotective cultures in fresh meat produced in Argentina. *Meat Scien.* 2008; 79: 483-499.
66. Cheong EYL, Sandhu A, Jayabalan J, et al. Isolation of lactic acid bacteria with antifungal activity against the common cheese spoilage mould *Penicillium commune* and their potential as biopreservatives in cheese. *Food Control.* 2014; 46: 91-97.
67. Crowley S, Mahony J, van Sinderen D. Current perspectives on antifungal lactic acid bacteria as natural bio-preservatives. *Trends Food Scien & Techn.* 2013; 33: 93-109.
68. Dalié DKD, Deschamps AM, Richard-Forget F. Lactic acid bacteria - Potential for control of mould growth and mycotoxins: A review. *Food Control.* 2010; 21: 370-380.
69. Collins JK, Thornton G, Sullivan GO. Selection of probiotic strains for human applications. *Int Dairy J.* 1998; 8: 487-490.
70. Tannock GW. *Probiotics: A Critical Review*, Horizon Scientific Press, Wymondham. U.K. 1999.

71. Fuller R. A review: Probiotics in man and animals. *J Appl Bact.* 1989; 66: 365-378.
72. Salminen S, Ouwehand A, Benno Y, et al. Probiotics: How should they be defined? *Trends Food Sci Techn.* 1999; 10: 107-110.
73. Champagne C, Gardner N, Roy D. Challenges in the addition of probiotic cultures to foods. *Crit Rev Food Sci Nutr.* 2005; 45: 61-84.
74. Champagne CP, Møllgaard H. Production of Probiotic Cultures and Their Addition in Fermented Foods. In: Farnworth ER (ed), *Handbook of Fermented Functional Foods*, 2nd ed, CRC Press, Boca Raton, FL. US. 2008; 513-532.
75. Gibson GR, Fuller R. Aspects of *in vitro* and *in vivo* research approaches directed toward identifying probiotics and prebiotics for human use. *J Nutr.* 2000; 130: 391-395.