

Research Article

Antimicrobial Effect of *Cyperus rotundus* Tuber Extract on the Microorganisms of the Urinary Tract Infection

Dadooka M¹, Mehrabianb S¹ and Irianc S^{2*}¹Department of Biological Sciences, Islamic Azad University, Iran²Department of Cell and Molecular Biology, Kharazmi University, Iran***Corresponding author:** Saeed Irianc, Department of Cell and Molecular Biology, Faculty of Biological Sciences, Kharazmi University, Tehran, Iran**Received:** August 21, 2019; **Accepted:** September 20, 2019; **Published:** September 27, 2019**Abstract****Background:** Medicinal use of plants dates to long back in the history of the human beings and has recently gained a great deal of importance in treating different diseases including the urinary tract infection.**Objectives:** In the present study the effects of *Cyperus rotundus* tuber extract on the microorganisms of the urinary tract infection was investigated.**Materials and Methods:** Ethanol extracts of *Cyperus rotundus* tuber was prepared by maceration. Antimicrobial effect of these extracts on the isolated strains was determined by disk diffusion and broth microdilution methods.**Results:** Results revealed a growth inhibitory concentration greater than 0.5 mg/ml of the ethanol extract on all the examined microorganisms of the urinary tract infection.**Conclusions:** It was thus concluded that *Cyperus rotundus* has a significant antimicrobial property with a potentially important role in the treatment of the urinary tract infection.**Keywords:** *Cyperus rotundus*; Antibactria; Urinary Tract Infection (UTI)**Introduction**

Urine is normally sterile; however, it can be subjected to infection by bacteria, viruses, parasites and fungi [1]. Urinary Tract Infection (UTI) is one of the most common infections in all age groups, and when left untreated may result in liver conditions, urinary apparatus disturbances, uremia, and premature birth in pregnant women. UTI is more common in children, pregnant women and diabetic patients [2-5]. Bacterial infection of the urine is one of the most important causes of kidney stone formation [6]. Gram negative bacteria, in particular *E. coli*, are the most common causes of urinary tract infection [7,8].

Plants have served as medicine since the ancient time [9]. Similar to antibiotics, the antimicrobial effects of medicinal plants have gained a great deal of attention [10,11]. Use of medicinal plants in the past several decades has increasingly turned popular such that a great number of the available medicines are of plant origin [12,13], and medicinal plants are currently being used for the treatment of many diseases including fungal infections [14].

Cyperus rotundus also known as nut grass as a member of the Cyperaceae family is distributed in humid, marshy, warm and moderate environments including Northern Iran with a local name of Teplagh. It is a perennial plant with a 20-40 cm height, long roots with rhizomes and a black shelled tuber with a white and odorous interior. Leaves are numerous, small, linear, dark green and spikelets are reddish brown [15-17]. The rhizome contains essential oil, Saponins, Vitamine C, Polyphenol and Flavonol glycoside [18,19]. *Cyperus rotundus* has had medicinal use in gastrointestinal bloating, stomach burning, stomach ulcers, menstrual periods, kidney problems, headaches and liquid mouthwash [20-22].

Due to the increasing importance of plants in herbal medicine

and their low rates of side effects, this study aimed at studying the antimicrobial effect of *Cyperus rotundus* extract on urinary tract infecting microorganisms.

Materials and Methods**Pathogenic microorganisms**

Pathogenic microorganisms were isolated from urine samples of urinary tract infected patients [23] and identified by biochemical tests [23,24].

***Cyperus rotundus* tuber extract preparation**

Cyperus rotundus tubers were collected from the agricultural lands of Aznava-Behnamir in Mazandaran province of Iran. Extracts were prepared by maceration or wetting by ethanol solution. Tubers were dried in a closed environment with air conditioning, and powdered. A total of 200 grams of the powder was added to 100 ml of ethanol and incubated for 48 h, and then filtered through a sterile paper filter. This filtered extract was then dried by incubation at 40°C.

Antimicrobial activity of the tuber extract

Strain sensitivity was determined by both disc diffusion and broth microdilution methods. Initially, suspensions of 0.5 McFarland (1.5×10^8 CFU/ml) were prepared [26].

In the disc diffusion method, using a sterile swap, suspensions were grown on Muller Hinton agar medium, while *Candida albicans* was grown on Saboro dextrose agar medium. Sterile blank discs were placed in plates containing different concentrations of the tuber extract and then placed on agar plates containing microorganisms. These plates were incubated at 37°C for 24 h, and then subjected to measurements of the zone of growth inhibition [17].

Table 1: Results of disc diffusion test.

Microorganism	Diameter of zone of growth inhibition (mm) at 25mg/ml of the extract
<i>Staphylococcus aureus</i>	15
<i>Staphylococcus epidemidis</i>	12
<i>Staphylococcus saprophyticus</i>	15
<i>Micrococcus</i>	12
<i>Escherichia coli</i>	14
<i>Enterococcus faecalis</i>	14
<i>Proteus</i>	15
<i>Klebsiella pneumoniae</i>	12
<i>Pseudomonas aeruginosa</i>	15
<i>Citrobacter</i>	15
<i>Candida albicans</i>	11

In the broth microdilution method, tubes containing Muller Hinton broth medium and Saboro Dextrose medium containing 0.5-20 mg/ml of tuber extract to a total volume of 1 ml were prepared and inoculated with 100 µl of 0.5 McFarland suspension. A solution of 0.05 mg/ml of the extract was also included as negative control. MIC values were determined after 24 h of incubation at 37°C [28].

Results

A total of five Gram positive bacteria including *Staphylococcus aureus*, *Staphylococcus epidemidis*, *Staphylococcus saprophyticus*, *Enterococcus faecalis*, *Micrococcus* and six different strains of Gram negative bacteria including *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus*, *Pseudomonas aeruginosa*, *Citrobacter* and one yeast strain the *Candida albicans* were isolated from urinary tract infected samples, and their sensitivity to tuber extract was determined.

Strain sensitivity to *Cyperus rotundus* tuber extract

In the disc diffusion method, zone of growth inhibition appeared at a minimum inhibitory concentration of 25 mg/ml for all UTI strains (Table 1).

In the broth microdilution method, the MIC value for all bacterial strains was at a concentration of 0.2 mg/ml, while that of *Candida albicans* was determined at 0.5 mg/ml. A greater sensitivity was observed for *Staphylococcus aureus*, *Staphylococcus saprophyticus*, *Pseudomonas aeruginosa* and *Citrobacter* with a MIC value at 0.5 mg/ml (Table 2).

Discussion

In the present study, the antimicrobial effect of *Cyperus rotundus* tuber ethanolic extract on urinary tract infecting microorganisms was investigated. Our collective results revealed that the extract did possess an antimicrobial property against all the examined UTI microorganisms. The MIC value for all bacterial strains was greater than 0.2 mg/ml, and *Candida albicans* had the greatest MIC value at 0.5 mg/ml. *Staphylococcus saprophyticus*, *Pseudomonas aeruginosa* and *Citrobacter* were the most sensitive strains.

Cyperus rotundus tuber extract has been shown to possess antimicrobial activity [29], while its inhibitory effect against *Streptococcus pyogenes* growth was demonstrated by Mehta et al. [30].

Table 2: MIC values of the ethanolic extract of *Cyperus rotundus* for microorganisms.

Microorganism	MIC value (mg/ml)
<i>Staphylococcus aureus</i>	0.1
<i>Staphylococcus epidemidis</i>	0.2
<i>Staphylococcus saprophyticus</i>	0.1
<i>Micrococcus</i>	0.2
<i>Escherichia coli</i>	0.2
<i>Enterococcus faecalis</i>	0.2
<i>Proteus</i>	0.2
<i>Klebsiella pneumoniae</i>	0.2
<i>Pseudomonas aeruginosa</i>	0.1
<i>Citrobacter</i>	0.1
<i>Candida albicans</i>	0.5

On the other hand, *Cyperus rotundus* whole plant extract has also been shown to be ineffective against strains of *E. coli*, *Pseudomonas aeruginosa* and *Salmonella typhi* [31]. Eltayebi et al. determined a MIC value of 12.5 mg/ml for their bacterial strains [17], while the maximum MIC value in our study was determined to be 0.5 mg/ml.

In conclusion, *Cyperus rotundus* extract possesses antimicrobial effect and thus inhibits the growth of urinary tract infected microorganisms and thus has a great potential in treating patients with UTI.

Acknowledgment

MD contributed in performing laboratory experiments. SM conceived the idea. SI prepared the manuscript.

References

- Sorkhi H, Jabarian A and Asqarian A. "E. Coli and Its Drug Resistance in Childhood Urinary Tract Infectious," Guilan University of Medical Sciences. 2005; 14: 8-23.
- Kliegman RM, Behrman RE, Jenson HB, Stanton BF. Nelson text book of pediatrics. 18th ed. Philadelphia: W.B. Saunders. 2007
- Behrman RE, Kliegman RM, Jenson HB. Urologic Disorders in Infants and Children. In: Elder JS, Nelson Textbook of Pediatrics Saunders. 2004; 1785.
- Mims CA, Dockrell HM, Goering RV. Medical Microbiology. 3rd ed. London: Mosby Co. 2004.
- Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. Dis Mon. 2003; 49: 53-70.
- Jan H, Akbar I, Kamran H, Khan J. Frequency of renal stone disease in patients with urinary tract infection. J Ayub Med Coll Abbottabad. 2008; 20: 60-62.
- Ejraes K. Urinary tract infection; study: Recurrent UTI caused by *E. coli* strain causing preceding UTI. Women's Health Weekly. Atlanta. 2004; 25: 166.
- Mims C, Dockrell H, Goering R, Roitt I, Wakelin D, Zuckerman M. Urinary tract infection. Medical microbiology. 3rd ed. New York: Mosby. 2004; 241.
- Cowan MM. Plant products as antimicrobial agents. Clin Microbiol Rev. 1999; 12: 564-582.
- Alviano DS, Alviano CS. Plant extracts: search for new alternatives to treat microbial diseases. Curr Pharm Biotechnol. 2009; 10: 106-121.
- Kumar P, Ansari SH, Javed A. Herbal remedies for the treatment of periodontal diseases. Recent Pat Drug Deliv Formul. 2009; 3: 221-228.

12. Grag SC, Denger SL. Antifungal activity of the essential oil of *Myrtus communis* var. *microphylla*. *Herba Hungarica*. 1998; 27: 123-124.
13. Sri Ranjani S. Medicinal uses and Pharmacological activities of *Cyperus rotundus* Linn—A Review. *International Journal of Scientific and Research Publications*. 2013; 3: 1-8.
14. Falahati M, Omid Tabrizi N, Jahanian F. Anti Dermatophyte Activity of *Eucalyptus camadulensis* in comparison with Griseofulvin. *Iranian J Pharm & Ther*. 2005; 4: 80-83.
15. Burkill HM. *The Useful Plants of West Tropical Africa*, Vol. 1. 2nd ed. Kew (UK): Royal Botanical Gardens. 1985.
16. Gunasekera TGLG and Fernando DNS. Agricultural importance, biology, control and utilisation of *Cyperus rotundus*. *Planter (Malaysia)*. 1994; 70: 537-544.
17. Eltayeib AA, Ismaeel HU. Extraction of *Cyperus rotundus* Rhizomes Oil, Identification of Chemical Constituents and Evaluation of Antimicrobial Activity of the Oil in North Kordofan State. *International Journal of Advanced Research in Chemical Science (IJARCS)*. 2014; 1: 18-29.
18. Adeniyi TA, Adeonipekun PA, Omotayo AE. Investigating the phytochemicals and Antimicrobial properties of three sedge (cyperaceae) species. *International journal of tropical medicine*. 2013; 8: 92-98.
19. Sivapalan SR. Physico-Chemical and Phyto-Chemical Study of Rhizome of *Cyperus rotundus* Linn. *International Journal of Pharmacology and Pharmaceutical Technology (IJPPT)*. 2012; 1: 42-46.
20. Raut NA, Gaikwad NJ. Antidiabetic activity of hydro-ethanolic extract of *Cyperus rotundus* in alloxan induced diabetes in rats. *Fitoterapia*. 2006; 77: 585-588.
21. Uddin SJ, Mondal K, Shilpi JA, Rahnan MT. Antidiarrhoeal activity of *Cyperus rotundus*. *Fitoterapia*. 2006; 77: 134-136.
22. Muthu K, Hema M, Nagaraj S, Rengasamy R. *In-vitro* antibacterial potential, Phytochemical characterization of *Cyperus rotundus* flower extract. *International Journal of Natural Products Research*. 2014; 4: 6-8.
23. Ohkawa M, Tokunaga S, Nakashima T, Yamaguchi K, Orito M, Hisazumi H. Composition of urinary calculi related to urinary tract infection. *J Urol*. 1992; 148: 995-997.
24. Griffith DP. Infection-induced renal calculi. *Kidney Int*. 1982; 21: 422-30.
25. Collee JG, Marinon BP, Fraser AG, Simmons A (eds.). Mackie and McCartney, *Practical Medical Microbiology*, 14th ed. Church-hill Livingstone. 1996; 131-151.
26. Chessbrough M. *District laboratory practice in tropical countries*. United Kingdom: Cambridge Univ Press. 2000; 222.
27. Harris DC. *Quantitative Chemical Analysis*, Eighth Edition. W.H. Freeman and Co., New York, NY. 2010; 750.
28. Sharma SK, Singh AP. Antimicrobial investigations on rhizomes of *Cyperus rotundus* Linn. *Der Pharmacia Lettre*. 2011; 3: 427-431.
29. Bisht A, Bisht GRS, Singh M, Gupta R and Singh V. Chemical composition and antimicrobial activity of essential oil of tubers of *Cyperus rotundus* Linn. collected from Dehradun (Uttarakhand). *International Journal of Research in Pharmaceutical and Biomedical Sciences*. 2011; 2: 661-665.
30. Mehta M, Bharmuche A and Bhatkala A. Investigation of the Anti Microbial and Anti-Inflammatory Effect of *Cyperus rotundus* on Tonsillitis. *International Journal of Current Engineering and Technology*. 2013; 27: 135-138.
31. Parekh J, Chanda S. *In-vitro* Antimicrobial Activities of Extracts of *Launaea procumbens* roxb. (Labiatae), *Vitis vinifera* L. (Vitaceae) and *Cyperus rotundus* L. (Cyperaceae). *African Journal of Biomedical Research*. 2006; 9: 89-93.