

## Original Article

## FOR LIVER: Herbs in Pakistani Population

Seema<sup>1</sup>, Humaira Mahmood<sup>2</sup>, Aneel Kapoor<sup>3</sup>,  
Saima Masood<sup>4</sup>, Asia Firdous<sup>5</sup> and Shah Murad<sup>6\*</sup>

<sup>1</sup>Research fellow at Public Health Institute, AFPGMI,  
NUMS, Rawalpindi Pakistan

<sup>2</sup>Associate Professor of Community Medicine, AFPGMI,  
NUMS, Rawalpindi Pakistan

<sup>3</sup>Associate Professor of Biochemistry, Bhitai Dental &  
Medical College, Mirpurkhas, Pakistan

<sup>4</sup>Senior Lecturer, Community Medicine at KIMS, Malir  
Karachi, Pakistan

<sup>5</sup>AP Pharmacology at LMDC, Lahore Pakistan

<sup>6</sup>HOD Pharmacology, Pharmacology and Therapeutics,  
IMDC, Islamabad Pakistan

\*Corresponding author: Dr. Shah Murad, Islamabad  
Medical & Dental College, Islamabad-Pakistan

Received: May 27, 2022; Accepted: June 23, 2022;

Published: June 30, 2022

## Abstract

Coronary artery disease is a complex syndrome to get initiation from formation of atherosclerotic plaques in systemic circulation to cardiac arrhythmia causing morbidity/mortality. This study was conducted to compare hypolipidemic effects of Niacin and Indian date (Jujuba) in hyperlipidemic patients. Study was conducted from July 2019 to December 2019 at National Hospital Lahore-Pakistan. Sixty participants were enrolled of both gender male and female patients' age range from 25 to 60 years. Consent was taken from all patients. They were divided in two groups. Group-I was advised to take 2 grams Niacin in divided doses for the period of two months. Group-II was advised to take 500 grams of fruit Jujube daily for the period of two months. All patients' systolic and diastolic blood pressure was noted. Their baseline LDL and HDL cholesterol was determined by conventional method of measuring Lipid Profile. After two months therapy, their post treatment blood pressure and lipid profile was measured and mean values with  $\pm$  SEM were analyzed biostatistically. Group-I which was on Niacin their blood pressure was reduced but it was non-significant change, LDL cholesterol decreased significantly and HDL cholesterol was increased significantly. In group-II patients LDL cholesterol was decreased significantly but HDL increase was not significant with p-value of  $>0.05$ . It was concluded from the research work that Niacin is potent in lowering LDL and increasing HDL cholesterol, while Jujube has significant effect as LDL cholesterol lowering potential, but it does not increase HDL cholesterol significantly. Jujubes and vitamin B-3 did not reduced blood pressure, when analyzed statistically.

## Introduction

Hypolipidemic drugs decrease risk of atherosclerosis and its complications. Therefore, these drugs may be used in prevention of heart attack, peripheral vascular disease and ischemic stroke [1]. Commonly used medications for treatment of Hyperlipidemia include Statins, Fibrates, Niacin, and Resins. All these medicines have potential for SEs and low compliance due to one reason or another [2]. Niacin when given in hypolipidemic doses i.e. more than 2 grams per day it causes partial inhibition of release of free fatty acids from adipose tissue, and increased lipoprotein lipase activity, which may increase the rate of chylomicron triglyceride removal from plasma. Niacin decreases the rate of hepatic synthesis of VLDL and LDL by synthesis of apoproteins which are integral part of LDL or VLDL structure [3]. Some herbs have been proved to reduce plasma lipids in human population. Jujubes or Ziziphus jujube have somewhat hypolipidemic as well as hypoglycemic effects [4]. Jujube fruit is known to contain considerable amount of phenolic compounds, including chlorogenic acid, gallic acid, protocatechuic acid and caffeic acid [5]. High polyphenolic content of Z Jujube suggests its potent capacity in clearing of oxidants. Many studies proved the hepatoprotective effect of methanolic extract of Zizyphus jujube fruits. Histopathological studies supported the biochemical findings. Study concludes a hepatoprotective activity probably due to its antioxidant effect [6]. Some studies evaluated the effect of Z Jujube fruit in controlling dyslipidemia in obese adolescents. A triple-blind randomized placebo-controlled trial of 86 obese adolescents aged 12--18 with dyslipidemia proved its hypolipidemic features. Results showed the fruits to be generally well tolerated, with potential favorable effects on blood pressure and serum lipid profile [7].

Hydro-alcoholic preparations from Indian dates have been proved to possess hypotensive and hypolipidemic effects in human population. Main problem with ingestion of Indian dates or Jujuba was its larger amount to take by human population to reduce blood pressure and hyperlipidemia. It is proved in various studies that if taken in high amount Indian dates are antioxidant and anti-inflammatory in characteristics. Their cardioprotective and hepatoprotective effects are explained in various studies [8]. In ancient Chinese book on herbal medicine Huangdi Neijing (475-221 BC), jujube was described as one of the five most valuable fruits in China. In Shennong Bencaojing (300 BC-200 AD), an earlier book recoding medicinal herbs, jujube was considered as one of the superior herbal medicines that prolonged our life-span by nourishing blood, improving quality of sleep, and regulating digestive system [9].

## Subjects and Method

The research work was started after written approval of ETHICS COMMITTEE of National Hospital, Lahore. It was conducted from July 2019 to December 2019. Sixty hyperlipidemic patients were selected from National Hospital Lahore-Pakistan to compare hypolipidemic effects of Niacin (vitamin B-3) and commonly used fruit in winter season in Pakistan i.e. Indian date or Jujube (Bair in urdu). Both male and female patients suffering from primary or secondary hyperlipidemia were selected. The age limit for patients was 25 to 60 years. Exclusion criteria were alcoholics, cigarette smokers, habitual to enjoy sedentary life, with impaired liver or renal functions. Written and already explained consent was taken from all participants. Their systolic/diastolic BP was measured by using mercury sphygmomanometer. Baseline Lipid Profile was determined

**Table:** Illustrating mean values with  $\pm$  SEM and p-values in two groups of hyperlipidemic patients before and after treatment.

	LDL-C	HDL-C	SYSTOIC BP	DIASTOIC BP
Before treatment	G1= 210.1 $\pm$ 2.11	37.9 $\pm$ 1.91	110.02 $\pm$ 1.12	86.87 $\pm$ 2.01
	G2= 198.8 $\pm$ 2.17	38.6 $\pm$ 2.19	113.98 $\pm$ 2.86	89.66 $\pm$ 1.04
After treatment	G1= 180.9 $\pm$ 2.22	45.2 $\pm$ 2.19	108.55 $\pm$ 2.87	85.84 $\pm$ 1.00
	G2= 190.9 $\pm$ 1.73	41.9 $\pm$ 2.97	112.56 $\pm$ 2.67	87.52 $\pm$ 2.22
Change in mg/dl	G1= 29.2	7.3	1.47	1.03
	G2= 7.9	3.3	1.42	2.14
Change in %	G1= 13.9 %	16.2 %	1.33%	1.18%
	G2= 4.0 %	7.9 %	1.24%	2.38%
p-value	G1= <0.001	<0.001	>0.05	>0.05
	G2= >0.05	<0.01	>0.05	>0.05

KEY: G1 is group on Niacin, G2 is group on drug-2 ie Jujube,  $\pm$  stands for Standard Error of Mean values, BP stands for blood pressure, p-value >0.05 is non-significant change, p-value <0.01 is used for significant change in parameter, and p-value <0.001 is highly significant change in tested parameter.

in Biochemistry lab of the Hospital. Patients were divided in two groups, 30 patients in each group. Group-I was on Tab. Niacin 2 grams daily in three divided doses. Group-II was on Jujube 500 grams daily in three divided times to eat. They were advised to take fruit and vitamin B-3 for two months. They were also advised not to take junk food and were also advised to do 35 minutes brisk walk daily in the morning or evening. Mean values  $\pm$  SEM were taken for statistical analysis using SPSS version 10.0, 2015. Paired 't' test was applied to get significance changes in parameters before and after treatment. P-value >0.05 was considered as non-significant change, p-value <0.01 was considered as significant and p-value <0.001 was considered as highly significant change in the parameter.

## Results

With two months therapy by vitamin B-3 (Niacin) and Indian dates (Jujube or Jujuba), systolic/diastolic blood pressure was not changed when analyzed statistically but plasma total cholesterol, LDL-cholesterol and HDL-cholesterol were changed, which are shown in following (Table).

## Discussion

Coronary artery disease is complex phenomenon starting from dyslipidemia to cardiac failure to ventricular fibrillation and mortality due to heart attack. Hypolipidemic drugs decrease chances of LDL particles available for oxidation, so prevent CAD. Vitamin B-3 (Niacin) is commonly used hypolipidemic drug which inhibit lipoprotein lipase activity, so lesser formation of free fatty acids will be available which are main sources of TG-rich lipoproteins (VLDL) formation. Lesser amount of VLDL leads to lesser synthesis of LDL particles which are rich in cholesterol. In our research study it was proved that no vitamin B-3 or Indian dates decrease systolic or diastolic blood pressure when pre and post treatment results were analyzed biostatistically. In our results vitamin B-3 (Niacin) 2 grams daily intake for two months decreased LDL-cholesterol about 13.9 % which is highly significant change. HDL-cholesterol in this group increased about 16.2 % which is again highly significant change. ZQ Zhu et al [10] and W Cao et al [11] proved same results when they used 2 grams of Niacin in 66 hyperlipidemic patients, but WB Yao et al [12] observed lesser effects of Niacin on HDL cholesterol, i.e. only 4.4 % increase in HDL cholesterol. Hung PG et al [13] explained different mechanisms of hypolipidemic response of Nicotinic acid on persons with different genetic code. One of the favorable mechanisms for patients with CAD they described is fibrinolytic activity of Niacin. In our results Jujube fruit decreased LDL cholesterol is 7.9 mg/dl, which is significant change in the parameter. HDL cholesterol is not

increased significantly in our results with p-value of >0.05. Tan H et al [14] and Tripathi M et al [15] observed same reason of Jujube on LDL and HDL-cholesterol, which augment our results. Tschesche R et al [16] observed more effects of Jujube as we observed in low density lipoprotein cholesterol. Um S et al [17] proved that LDL cholesterol is much decreased as compared to our results. KB Kang et al [18] observed too less effects of Jujube fruit in 5 hyperlipidemic patients. This difference in two studies is due to their small sample size, i.e. they tried herb only on five hyperlipidemic patients, while we tried in 30 hyperlipidemic patients. Compliance of Indian fruit Jujuba is low due to its very high amount to take as hypolipidemic and hypotensive herbs. It was explained by Loom Viyu et al [19] and A. Mohammad et al [20].

## References

- S Nawaz, M Furqan, Z A Kuliev, Yu A Akhmedov, A D Vdovin, N D Abdullaev. Atherosclerosis and its complications. Chemistry of Natural Compounds. 2012; 33(2): 165-73.
- Anbarasi Balakrishnan, Parimala Devi Balasubramaniam, Senthil Kumar Natesan. Conventional therapies for dyslipidemia. J Adv Sci Res. 2013; 3(3): 40-3.
- Reza Afrisham, Mohammad Aberomand, Mohammad Ali Ghaffari, Amir Shahpoosh, Mostafa Jamal. Vitamin B-3: new considerations. Journal of Botany. 2015; 5(3): 119-21.
- Usama El-Sayed Mostafa and Louay Labban. Effect of Zizyphus jujuba on serum lipid profile and some anthropometric measurements. Advancement in Medicinal Plant Research. 2013; 1(3): 49-55.
- Shah M, LF Juju, FF Jau, UI Lyme, KT Fauo, ML Panic. Phenolic compounds in Indian dates. Ethanobotanical J. 2014; 12(5): 222-7.
- Daneshmand F, Zare-Zardini H, Tolueinia B, Hasani Z, Ghanbari T. Crude Extracts from Ziziphus Jujuba Fruits: new guidelines. Iranian Journal of Pediatric Hematology Oncology. 2013; 3(3): 191-3.
- Bruckert E, Labreuche J, Amarenco P. Compliance of Indian dates when used in high amount. Atherosclerosis. 2010; (2): 353-61.
- Duggal JK, Singh M, Attri N, Singh PP, Ahmed N, et al. Hypotensive and hypolipidemic effects of Indian fruits. Journal of cardiovascular pharmacology and therapeutics. 2013; 19(1): 58-60.
- Manoj P, Shiv Kumar, K G Bhat. Effect of Indian date leaves extracts on phagocytosis. JOURNAL OF NATURAL REMEDIES. 2014; 4(1): 47.51.
- ZQ Zhu, JW Chen, TX Hu, DY Zhu. Niacin affects HDL-cholesterol and LDL cholesterol. Phytochemistry 2011; 22: 1667-72.
- W Cao, BN Wang, H Gao, MT Fan, JB. Vitamin B-3 or Niacin for prevention of coronary artery disease. Plant Pharmacol. 2011; 61: 703-7.
- WB Yao, H Wang, XD Gao, GC Zhou, L Cai. Niacin is important for healthy heart. Science of pharmacol. 2011; 10: 122-7.
- Hung PJ, J Wang, LT Li, Y Dan. Vitamin B-3 for Atherosclerosis and its prevention. J. Food Chem 2011; 10: 21-4.

14. Tan NH, Fan HG, Zhou J. JUJUBE IS HYPOLIPIDEMIC HERB. *Chem. Rev.* 2007; 106: 840-5.
15. Tripathi M, Pandey MB, Jha RN, Pandey VB, Tripathi PN, Singh JP. Cyclopeptide alkaloids from *Zizyphus jujuba*. *Fitoterapia.* 2001; 72(5): 507-510. doi:10.1016/S0367-326X(01)00278-7.
16. Tschesche R, Khokhar I, Wilhelm H, Eckhardt G, Jubanine-A, Jubanine-B. Alkaloids from *Zizyphus jujuba* have antioxidant characteristics. *Phytochemistry.* 2010; 15: 541-2.
17. Um S, Choi TJ, Kim H, Kim BY, Kim SH, et al. Hypolipidemic effects of *Z. Jujube*. *J Org Chem.* 2013; 78: 321-29.
18. KB Kang, JJ You, LL mumpycal, ST Simon. *Z. Jujube* is effective to reduce oxidation burden. *Phytochemistry.* 2014; 22(4): 222-5.
19. Loom viyu, PK Joshi, Roomyu L, Furquye Y. Indian dates and its compliance in human population. *J ethanobotany.* 2014; 4(4): 123-9.
20. Mohammad, S Mohammad, N Mohammad, N Abdullah. Indian fruits varieties and their characteristics in various therapies. *Ir J of Nat Med.* 2013; 12(1): 444-9.