

Editorial

The Importance of Recent Advances in Liquid Chromatography Techniques to the Biomedical Field

Florence CC Tan¹, Tzi Bun Ng² and Evandro Fei Fang^{3*}

¹Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

²School of Biomedical Sciences, Faculty of Medicine, Chinese University of Hong Kong, Hong Kong

³Laboratory of Molecular Gerontology, National Institute on Aging, USA

*Corresponding author: Evandro Fei Fang,

Laboratory of Molecular Gerontology, National Institute on Aging, NIH, USA, Tel: 410 558 8217; Email: evandro.fang@nih.gov or evandrofeifang@yahoo.com

Received: November 19, 2014; Accepted: November 20, 2014; Published: November 21, 2014

Editorial

The versatility and applicability of Liquid Chromatography (LC) techniques have long been recognized and consequently used for numerous analytical problems in fields ranging from bioanalysis, environmental analysis to materials analysis. Recent advances in the field of chromatography including innovative supports and instrumentation lead to a higher separation efficiency and faster analysis time, which have further expanded its applicability [1]. Especially for pharmaceuticals and biomedical research, the multitude of chromatographic techniques plays an increasingly important role.

Since natural compounds with medicinal applications can be separated based on many different properties, an array of different separation techniques will allow highly purified analytes, which are useful for detecting and characterizing proteins and small organic molecules, which has not been previously described in quest of new therapeutics [2-5]. One example is the purification of the Blue Tiger King Lectin (BTKL) from the *Phaseolus vulgaris* plant using a multistep purification protocol involving ion exchange chromatography, affinity chromatography and gel filtration [6]. For subsequent characterization, a Fast Protein Liquid Chromatography (FPLC) gel-filtration method was applied to determine the molecular mass under native conditions, and the N-terminal amino acid sequence was analyzed with HP 1000A Edman degradation and HP 1000 HPLC system. BTKL showed hemagglutinating antiproliferative activity against Hep G2 cells, a perpetual cell line derived from well-differentiated hepatocellular carcinoma tissue, and yet no toxicity is found against the normal human cell line WRL 68. Thus, it may provide a new therapeutic candidate against hepatocellular carcinoma, a globally common cancer with currently limited options of treatment and poor prognosis.

A widely used technique is High-Performance Liquid Chromatography (HPLC), which utilizes pumps to drive a pressurized liquid sample through the column. Introduction of sub 2 μ m particles offers simultaneously higher resolution, better sensitivity and faster

analysis time due to the increased surface area and the ensuing requirement of higher backpressure. The latter has given rise to the name Ultra-High Performance Liquid Chromatography (UHPLC). Another innovation of this UHPLC is the fused core consisting of a solid core, most commonly made of silica, coated by a porous material, which determines the mode of interaction with the sample components. This shortens the diffusion path, leading to decreased resistance to mass transfer and thereby minimizing peak broadening [1]. The advantages of UHPLC to conventional HPLC enables analyses of much more complex samples with a growing number of analytes, which have implicated its use in fields such as toxicology, forensics and doping control.

The analysis of Traditional Chinese Medicines (TCM) is one interesting application [7]. These compounds have been used for centuries in China and are typically composed of a cocktail of herbal ingredients with many active constituents believed to act synergistically. The concentrations of the active substances are dependent on breed, growing conditions and processing techniques inter alia, making the individual samples highly complex and difficult to perform quality control on. UHPLC with a variation of detecting methods including Diode Array Detector (DAD), Mass Spectrometry (MS) and Quadrupole Time-of-Flight MS (QTOF-MS) has allowed many quantitative studies determining the active ingredients in a multitude of TCM concoctions, providing an analysis time of 2-8 min and separation of some eighty constituents. Purification and characterization of these active ingredients in the so-called fingerprint analyses allow better controlled use of TCM and have been accepted by the WHO for the assessment of herbal medicines. Furthermore, it may lead to the discovery of substances that may be introduced as new therapeutics in western medicine.

The higher separation efficiency of UHPLC compared to conventional HPLC is also utilized in many emerging omic fields, such as genomics, proteomics and metabolomics. Such studies involve very complex samples and they are driving the demand for increasingly more efficient and high-throughput separation techniques. Being fast and accurate UHPLC-MS has become the predominant analytical technology in global metabolite profiling. This holistic approach holds especially great promise for identifying new biomarkers for early disease detection as well as disease progression. Drug efficacy prediction is another useful application [8]. Much of the metabolomics research has focused on finding biomarkers for cancer [9]. Attaining better diagnostic tools is expected to have a significant impact on the prognoses in particular for those cancer forms in which the majority of cases have already progressed to advanced stages at the time of diagnosis. The greatest challenge for untargeted metabolite profiling continues to be, however, whether instrumentation can provide a high enough resolution and sensitivity [10]. It will be interesting to follow the development and new applications for UHPLC and other

chromatographic techniques in the biomedical field as improvements in methods and instrumentation are constantly being achieved.

Disclaimer

This editorial was written in a personal capacity (EFF) and does not represent the opinions of the National Institute on Aging, the US Department of Health and Human Services, or the US Federal Government.

Acknowledgement

EFF is supported by the Intramural Research Program of the NIH, National Institute on Aging. We thank Drs. Joseph K. Hsu and Prabhat Khadka for critical reading of the manuscript.

References

1. Guillaume D, Ruta J, Rudaz S, Veuthey JL. New trends in fast and high-resolution liquid chromatography: a critical comparison of existing approaches. *Anal Bioanal Chem.* 2010; 397: 1069-1082.
2. Pavel M, Rubinsztein DC. Autophagy upregulation as a therapeutic strategy for neurodegenerative diseases. In: EF Fang, TB Ng, editors. *Antitumor potential and other emerging medicinal properties of natural compounds.* Springer, Dordrecht. 2013; 227-238.
3. Fang EF, Ng TB. Bitter gourd (*Momordica charantia*) is a cornucopia of health: a review of its credited antidiabetic, anti-HIV, and antitumor properties. *Curr Mol Med.* 2011; 11: 417-436.
4. Fang EF, Ng TB, Shaw PC, Wong RN. Recent progress in medicinal investigations on trichosanthin and other ribosome inactivating proteins from the plant genus *Trichosanthes*. *Current Med Chem.* 2011; 18: 4410-4417.
5. Fang EF, Ng TB. Ribonucleases of different origins with a wide spectrum of medicinal applications. *Biochim Biophys Acta.* 2011; 1815: 65-74.
6. Fang EF, Pan WL, Wong JH, Chan YS, Ye XJ, Ng TB. A new *Phaseolus vulgaris* lectin induces selective toxicity on human liver carcinoma Hep G2 cells. *Arch Toxicol.* 2011; 85: 1551-1563.
7. Huang H, Liu M, Chen P. Recent Advances in Ultra-High Performance Liquid Chromatography for the Analysis of Traditional Chinese Medicine. *Anal Lett.* 2014; 47: 1835-1851.
8. Gika HG, Theodoridis GA, Plumb RS, Wilson ID. Current practice of liquid chromatography-mass spectrometry in metabolomics and metabonomics. *J Pharm Biomed Anal.* 2014; 87: 12-25.
9. Armitage EG, Barbas C. Metabolomics in cancer biomarker discovery: current trends and future perspectives. *J Pharm Biomed Anal.* 2014; 87: 1-11.
10. Theodoridis GA, Gika HG, Want EJ, Wilson ID. Liquid chromatography-mass spectrometry based global metabolite profiling: a review. *Anal Chim Acta.* 2012; 711: 7-16.