

Case Report

A 68-Year-Old Man Presenting to the ER with Sudden Dyspnea, A Rare Case of Snowstorm View in the Right Ventricle during Echocardiography

Arman Hakemi^{1*}; Elham Angouraj Taghavi²¹Department of Clinical Sciences, Shahroud University of Medical Sciences, Iran²Student Research Committee, School of Medicine, Shahroud University of Medical Sciences, Shahroud, Iran***Corresponding author: Arman Hakemi**

Department of Clinical Sciences, Shahroud University of Medical Sciences, Iran

Received: February 20, 2023

Accepted: March 28, 2023

Published: April 04, 2023

Introduction

Pulmonary Thromboembolism (PTE) is the third leading cause of cardiovascular death worldwide [1]. Smoking cigarettes, malignancies, obesity, age, heredity, prolonged comorbidities, and surgical history are known risk factors for PTE [2,3]. PTE can present with common symptoms that can be misdiagnosed as other critical or non-critical diseases [1,4]. Some of the symptoms include respiratory distress, chest pain, presyncope or syncope, and bloody sputum [2,5,6]. Echocardiography in patients with PTE may represent different views; however, a snowstorm view in the ventricle is rare to report since yet.

Herein, we report a rare case of PTE with a snowstorm view in the right ventricle during echocardiography.

Case Presentation

A 68 year-old Caucasian man without a history of cardiovascular or pulmonary diseases presented to the Emergency Room (ER) with sudden onset of dyspnea. He was a known case of bone cancer with metastasis to the peritoneum and was undergoing chemotherapeutic treatment. Physical examination revealed edema in the right lower extremity. His Heart Rate (HR) was 125 beats/min, with Blood Pressure (BP) 80/60 mmHg, oxygen saturation (SpO₂) 65% and had no fever. Laboratory examinations, including complete blood cell count, diff, creatinine, blood urea nitrogen and electrolytes were within the normal range. Except a two plus positive C-reactive protein was reported. D-dimer was requested, but wasn't reported until the patient was in the ER. Bedside echocardiography revealed Right Ventricle (RV) enlargement and a snowstorm view in the

Abstract

Pulmonary Thromboembolism (PTE) is a fatal condition and the third leading cause of hospital-related deaths. We present a case of acute PTE in a 68 year-old man with dyspnea. Patient had a history of bone malignancy. Bedside echocardiography revealed snowstorm view in the right ventricle. Patient was treated with 10 units reteplase intravenous bolus. Echocardiography was repeated and no signs of snowstorm view were detected. Finally, patient died of cardiogenic shock and massive PTE. In this case we report snowstorm view during echocardiography which is rare and should be considered in clinical assessment of PTE.

RV and inferior vena cava (Figure 1). The patient was subsequently treated with 10 units of a reteplase intravenous bolus over 2 min. Echocardiography was repeated 30 minutes after injection and no snowstorm view was detected again. However, no change was observed in RV size. The patient was intubated because of worsening dyspnea. For hypotension, 15mcg/min norepinephrine and 500cc normal saline serum were administered. Nevertheless, BP did not improve. Finally, after 3.5 hours patient died in the ER due to the massive PTE.




Figure 1: Bedside echocardiography. Before (A) and after (B) thrombolytic therapy.

References

1. Alirezaei T, Mousavi S, Hesami H. A rare presentation of pulmonary thromboembolism as seizure. *Archive of Clinical Cases*. 2022; 9: 136-139.
2. Ebrahimi M, Arab MM, Moghadam HZ, Yazdi MJ, Foroughian M. Risk Stratification of Pulmonary Thromboembolism using Brain Natriuretic Peptide and Troponin I; a Brief Report. *Archives of Academic Emergency Medicine*. 2022; 10: e8.
3. Gjonbrataj E, Kim JN, Gjonbrataj J, Jung HI, Kim HJ, et al. Risk factors associated with provoked pulmonary embolism. *The Korean journal of internal medicine*. 2017; 32: 95-101.
4. Alirezaei T, Sattari H, Irilouzadian R. Significant decrease in plas-mad-dimer levels and mean platelet volume after a 3-month treatment with rosuvastatin in patients with venous thrombo-embolism. *Clinical Cardiology*. 2022; 45: 717-22.
5. Miniati M, Prediletto R, Formichi B, Marini C, Di Ricco G, et al. Accuracy of clinical assessment in the diagnosis of pulmonary embolism. *American journal of respiratory and critical care medicine*. 1999; 159: 864-71.
6. Wells PS, Ginsberg JS, Anderson DR, Kearon C, Gent M, et al. Use of a clinical model for safe management of patients with suspected pulmonary embolism. *Annals of internal medicine*. 1998; 129: 997-1005.
7. Konstantinides SV, Meyer G, Becattini C, Bueno H, Geersing G-J, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS) The Task Force for the diagnosis and management of acute pulmonary embolism of the European Society of Cardiology (ESC). *European heart journal*. 2020; 41: 543-603.
8. Bagot CN, Arya R. Virchow and his triad: a question of attribution. *British journal of haematology*. 2008; 143: 180-90.
9. Kucher N. Deep-vein thrombosis of the upper extremities. *New England Journal of Medicine*. 2011; 364: 861-9.
10. Adhikari S, Vaidya N, Poudel P, Pathak S. Successful thrombolysis with low dose thrombolytic agent in a patient with acute life-threatening massive pulmonary thromboembolism: A case report. *Annals of Medicine and Surgery*. 2022; 82: 104742.
11. Urikura A, Yoshida T, Endo M, Asakura K, Sato R, et al. Computed tomographic pulmonary angiography: Three cases of low-tube-voltage acquisition with a slow injection of contrast medium. *Acta Radiologica Open*. 2022; 11: 20584601221131476.