

## Clinical Image

# Congenital Unilateral Lamellar Cataract

Habiba T\*, Afaf E-R, Zahra HF, Elhassan A and

Amina B

Ophthalmology B, Ibn-Sina University Hospital, Rabat, Morocco

\*Corresponding author: Tazi Habiba, Ophthalmology B, Ibn-Sina University Hospital, Rabat, Morocco

Received: December 03, 2021; Accepted: December 29, 2021; Published: January 05, 2022

## Clinical Image

Lamellar or zonular cataract is a hereditary cataract that has a variable phenotype. It is transmitted in an autosomal dominant mode [1,2]. The crystalline opacities are fine and dispersed and are located at the level of the primary fibers in the embryonic nucleus [2]. This cataract is usually bilateral and asymmetrical. Sometimes the whole nucleus can be affected causing a significant decrease in visual acuity [1].

We report the case of a 7-year-old patient who was consulting for a decrease in visual acuity. The patient had no notable pathological history. His visual acuity at Snellen chart was 4/10 in the right eye and 10/10 in the left eye. The slit lamp revealed a lamellar cataract on the right eye (Figure 1). The fundus had no particularity. The patient underwent cataract surgery with an intraocular implant. The operative follow-up was uneventful and the visual acuity has improved to 10/10.




Figure 1: Congenital Unilateral Lamellar Cataract.

## References

1. Bu L, Jin Y, Shi Y, Chu R, Ban A, et al. Mutant DNA-binding domain of HSF4 is associated with autosomal dominant lamellar and Marnier cataract. *Nat Genet.* 2002; 31: 276-278.
2. Patil B, Sharma R, Nayak B, Sinha G, Khokhar S. Pediatric cataract. *Delhi J Ophthalmol.* 2015; 25: 160-165.