

Letter to the Editor

Ultrasound Applications in Nasal Bone Fractures

Heidari SF*

Department of Emergency Medicine, Emam Khomeini Hospital, Medical Faculty, Ilam University of Medical Sciences, Ilam, Iran

***Corresponding author:** Seyed Farshad Heidari, Department of Emergency Medicine, Emam Khomeini Hospital, Medical Faculty, Ilam University of Medical Sciences, Ilam, Iran**Received:** August 16, 2018; **Accepted:** August 28, 2018; **Published:** September 04, 2018

Letter to the Editor

Nasal bone fracture is one of the most usual fractures in patients with facial trauma [1]. It constitutes one third of the cases of facial bones fractures and is more common in the males than the females [2]. Generally, younger ones are more susceptible to displaced fractures and the old people develop comminuted fractures. Almost three quarters of cases of nasal bone fractures occur between the middle third and the distal third of the nasal bone [3].

Although physical examination is the gold standard manner for the diagnosis of nasal bone fracture, but it cannot determine the convolution of the fracture [4]. Because of the high prevalence of nasal bone fracture in facial trauma, it is essential for an accurate imaging method to diagnosis such injuries. Missed diagnosis or inadequate reduction can lead to nasal deformity or other complications. Therefore, quick diagnosis of nasal bone fracture and confirmation of proper reduction is very important to prevent possible complications [5].

Prior studies show that the plain radiography was negative in one quarter of cases of patients with nasal bone fractures who needed surgical treatment [6]. Also, facial computed tomography (CT) evaluation of these patients imposes a high risk of radiation

to the patient. When comparing ultrasonography (US) with CT-scan and plain radiography, US is inexpensive, available and simple to perform, and has no risk of ionizing radiation to the patient. The diagnostic value of ultrasound as a diagnostic modality in comparison with another evaluation method for diagnosis of nasal fracture lines has been shown in previous studies [4,7]. The other advantage of ultrasound over plain radiograph is that, it can show trauma of the cartilaginous part of the nose more accurately [4]. Also, it is useful for intra-operative reduction of the fractured nasal bone and in decreasing unnecessary manipulation of the fractured site for patients undergoing local anesthesia reduction [8,9].

References

1. Hwang K, You SH, Kim SG, et al. Analysis of nasal bone fractures; a six-year study of 503 patients. *J Craniofac Surg*. 2006; 17: 261-264.
2. Fraioli RE, Branstetter BF, Deleyiannis FW. Facial fractures: beyond Le Fort. *Otolaryngol Clin North Am*. 2008; 41: 51-76.
3. Piliija V, Buljick M, Mihalj M, et al. Classification and qualification of nose injuries—clinical and forensic aspects. *Med Pregl*. 2005; 58: 33-36.
4. Hong HS, Cha JG, Paik SH, et al. High-resolution sonography for nasal fracture in children. *AJR Am J Roentgenol*. 2007; 188: 86-92.
5. Park CH, Joung HH, Lee JH, et al. Usefulness of ultrasonography in the treatment of nasal bone fractures. *J Trauma*. 2009; 67: 1323-1326.
6. Damman F. Imaging of paranasal sinuses today. *Radiologe*. 2007; 7: 576, 578-583.
7. Thiede O, Krömer JH, Rudack C, et al. Comparison of ultrasonography and conventional radiography in the diagnosis of nasal fractures. *Arch Otolaryngol Head Neck Surg*. 2005; 131: 434-439.
8. Green KM. Reduction of nasal fractures under local anesthetic. *Rhinology*. 2001; 39: 43-46.
9. Park CH, Joung HH, Lee JH, et al. Usefulness of ultrasonography in the treatment of nasal bone fractures. *J Trauma*. 2009; 67: 1323-1326.