

Research Article

Association between cSNPs of BMP2 Gene and Degenerative Lumbar Scoliosis in Korean Population

Hyung-Ki Kim¹, Hwayoung Lee¹, Jun-Tack Kwon¹,
Ki Tack Kim², Hak-Jae Kim^{1,3*}

¹Department of Clinical Pharmacology, Soonchunhyang University, Republic of Korea

²Department of Orthopedic Surgery, Kyung Hee University East West Neomedical Center, Republic of Korea

³Soonchunhyang Medical Research Institute, Soonchunhyang University, Cheonan, Republic of Korea

*Corresponding author: Hak-Jae Kim, Department of Clinical Pharmacology, College of Medicine, Soonchunhyang University, Republic of Korea, Tel: +82 41 570 2478; Fax: +82 41 575 2406; Email: hak3962@sch.ac.kr

Received: July 31, 2014; Accepted: October 28, 2014;

Published: October 29, 2014

Abstract

Background: Bone morphogenetic proteins (BMPs) are involved in cartilage and bone formation. However, the association between single nucleotide polymorphisms (SNPs) of *BMP* genes and degenerative lumbar scoliosis (DLS) has not been investigated yet. The aim of this study was to determine whether coding SNPs (cSNPs) of *BMP* genes (*BMP2*, 3, 4, 5, 6, and 10) are associated with DLS in Korean population.

Methods: Seven cSNPs in *BMP* genes were selected and genotyped for 66 patients with DLS and 127 healthy controls using direct sequencing.

Results: Of the SNPs examined, two cSNPs of *BMP2* gene showed weak associations with DLS in the codominant (rs235768, Arg190Ser, missense, OR = 2.28, 95% CI = 1.19 – 4.36, $p = 0.03$) and dominant models (rs235768, OR = 1.99, 95% CI = 1.07 – 3.72, $p = 0.03$; rs1049007, Ser87Ser, synonymous, OR = 1.93, 95% CI = 1.039 – 3.62, $p = 0.04$).

Conclusion: *BMP2* may be a susceptible gene for DLS in Korean population.

Introduction

Lumbar scoliosis is a three-dimensional deformity of spine associated with structural alterations of vertebral bodies. Lumbar scoliosis in adult stage could be from deformity developed at growing age or de novo degenerative deformity after skeletal maturity [1-6]. Degenerative lumbar scoliosis (DLS), “de novo” lumbar scoliosis, usually occurs after the fourth or fifth decade in patients without history of scoliosis. The curve is composed of a few vertebral bodies with its apex in the intervertebral space, most frequently at the L2 - L3 or the L3 - L4 level [7-9]. The causes of DLS development are various. Degeneration of the spinal column is the most common cause. Neuromuscular disorders, metabolic abnormalities (osteoporosis), leg length discrepancy, long-standing pelvic obliquity, and outcomes of prior surgical interventions are other causes [10]. Several genetic variations have been linked to lumbar spine degeneration or osteoporosis [11, 12]. Intra genic polymorphisms of vitamin D receptor gene were reported to be associated with lumbar spine degeneration and bone density [11]. Gómez et al. (2007) reported that estrogen receptor alpha gene polymorphisms were associated with osteoporosis [12].

Bone morphogenetic proteins (BMPs) are phylogenetically conserved signaling growth factors belonging to the transforming growth factor beta super family [13-15]. BMPs also play important roles in the pathophysiology of several diseases, including osteoporosis [16], arthritis [17], pulmonary hypertension [17, 18], and kidney diseases [19]. Previous studies have reported that *BMPs* (*BMP2*, *BMP6*, *BMP7*, and *BMP15*) are associated with several diseases such as ossification of the posterior longitudinal ligament (OPLL), a vascular necrosis, and ovarian failure [20-26]. Wang et al. (2008) reported a positive association between *BMP2* polymorphisms [Ser37Ala (T/G) and Ser87Ser (A/G)] and OPLL of the cervical spine.

Despite the potentially important role of BMPs in the development of DLS, the association between genetic variations of *BMPs* and DLS has not been reported. Therefore, the objective of this study was to determine whether coding single nucleotide polymorphisms (cSNPs) of *BMP* family genes (*BMP2*, 3, 4, 5, 6, and 10) are associated with DLS in Korean population.

Methods

Subjects

All patients with DLS were from Kyung Hee University East-West Neo-Medical Center, Seoul, Republic of Korea and National Medical Center, Seoul, Republic of Korea. The DLS group included 66 patients with mean age of 69.1 ± 7.7 years (7 male, 65.4 ± 8.1 years; 59 female, 69.6 ± 7.6 years). The control group was recruited after it was confirmed in a general health check-up program that they had no clinical evidence of DLS or any other disorders. A total of 127 healthy controls with mean age of 68.1 ± 8.6 years (17 male, 71.6 ± 8.3 years; 110 female, 67.5 ± 8.6 years) were recruited. All case-control subjects used in this study were surveyed through the same center. Each patient was diagnosed by a special spine surgeon. All patients fulfilled physical examination and radiographic criteria (Cobb's angle over 10 degrees) [27]. Informed consent was obtained from all individuals according to the Declaration of Helsinki guidelines [28]. The study was approved by the ethics review committee of the Medical Research Institute, Kyung Hee University Medical Center, Seoul, Republic of Korea.

SNP Genotyping

First, we searched for cSNPs of *BMP* genes originally known as “*BMP*” (*BMP2/3/4/5/6/8/10/15*) in GenBank database (<http://www.ncbi.nlm.nih.gov/gene>). The SNPs of procollagen C-proteinase (*PCP*, same as *BMP1*), osteogenic protein 1 (*OPI*, same as *BMP7*), and

Table 1: Sequences of primers.

SNP/gene		Sequence(5'→3')	Product Size (bp)	Annealing temperature (°C)
rs235768	Forward	TTATCACCTCAGCAGAGCTTCA	375	58
BMP2	Reverse	GGCCAAAAGTTACTAGCAATGG		
rs1049007	Forward	GACGAGGTCCTGAGCGAGTTCCG	339	58
BMP2	Reverse	TACAGAAGCAAGAGTGGAAACG		
rs3733549	Forward	AGTTGTCCAGTGTCTGGAGGAT	351	58
BMP3	Reverse	TCCCTGTAAGCTTGATACCACA		
rs17563	Forward	CAGTAGGTTTCCCTGCATAAG	361	58
BMP4	Reverse	TCCAGTAGTCGTGTGATGAGGT		
rs3734444	Forward	GAGGATGTTGTGCTCAGAAATG	399	58
BMP5	Reverse	AGCATAAAGAGAGGTGCAGAGG		
rs17557	Forward	GAAAGGAGTGCTTTGATTCTGC	433	58
BMP6	Reverse	ACAAGGCTTTCTGGAAACACAG		
rs2231344	Forward	GGTGCTGGGGAGATATATGGA	326	58
BMP10	Reverse	CAGGTCCACTGGAAAAGCTATC		

Table 2: Genotype and allele frequencies of the *BMP* gene SNPs in degenerative lumbar scoliosis (DLS) and controls.

gene	SNP	Genotype	DLS		Control		Model	OR (95% CI)	p
			Freq	%	Freq	%			
BMP2	rs235768 (Arg190Ser)	T/T	36	54.5	90	70.9	codominant	2.28 (1.19-4.36)	0.03
		T/A	28	42.4	30	23.6	dominant	1.99 (1.07-3.72)	0.03
		A/A	2	3.0	7	5.5	recessive	0.52 (0.10-2.63)	0.40
BMP2	rs1049007 (Ser87Ser)	G/G	37	56.1	91	71.7	codominant	1.57 (0.92-2.67)	0.10
		G/A	27	40.9	31	24.4	dominant	1.93 (1.03-3.62)	0.04
		A/A	2	3.0	5	3.9	recessive	0.78 (0.15-4.17)	0.77
BMP3	rs3733549 (Arg192Gln)	G/G	44	66.7	94	74.0	codominant	1.40 (0.76-2.58)	0.29
		G/A	21	31.8	32	25.2	dominant	1.40 (0.73-2.70)	0.31
		A/A	1	1.5	1	0.8	recessive	1.95 (0.12-32.05)	0.64
BMP4	rs17563 (Val152Ala)	T/T	34	51.5	78	61.4	codominant	1.40 (0.86-2.28)	0.18
		T/C	27	40.9	43	33.9	dominant	1.46 (0.80-2.68)	0.22
		C/C	5	7.6	6	4.7	recessive	1.74 (0.51-6.00)	0.39
BMP5	rs3734444 (Ser37Ser)	T/T	49	74.2	86	67.7	codominant	0.77 (0.42-1.42)	0.40
		T/C	16	24.2	38	29.9	dominant	0.76 (0.39-1.49)	0.42
		C/C	1	1.5	3	2.4	recessive	0.61 (0.06-6.01)	0.66
BMP6	rs17557 (Val368Val)	C/C	37	62.7	77	61.1	codominant	0.83 (0.48-1.45)	0.51
		C/G	21	35.6	42	33.3	dominant	0.90 (0.48-1.72)	0.76
		G/G	1	1.7	7	5.6	recessive	0.31 (0.04-2.64)	0.23
BMP10	rs2231344 (Asp242Asp)	T/T	51	77.3	102	80.3	codominant	1.15 (0.57-2.31)	0.60
		T/C	15	22.7	24	18.9	dominant	1.20 (0.58-2.48)	0.62
		C/C	0	0	1	0.8	recessive	0.00 (0.00-NA)	0.41

Freq, frequency; OR, odds ratio; CI, confidence intervals; NA, not applicable; p, p value. Bold characters represent statistically significant values and its rs number of SNP.

GDF2/5/6/7/11 (same as *BMP9/14/13/12/11*) genes were excluded from this study. Related information of cSNP sequences was obtained from the SNP database (dbSNP #130) of the National Center for Biotechnology Information (NCBI). The SNPs with unknown heterozygosity and minor allele frequency (below 5%) were excluded. Finally, rs235768 (*BMP2*), rs1049007 (*BMP2*), rs3733549 (*BMP3*),

rs17563 (*BMP4*), rs3734444 (*BMP5*), rs17557 (*BMP6*), and rs2231344 (*BMP10*) were selected.

Genomic DNA was extracted from blood samples collected in EDTA using a commercially available Qiagen DNA Extraction kit (Qiagen, Tokyo, Japan). Genomic DNA was amplified using the gene-specific primers for each SNP (Table 1). PCR products were sequenced

Table 3: The haplotype frequencies of the *BMP* gene SNPs in degenerative lumbar scoliosis (DLS) patients and healthy controls.

Haplotype	Type	DLS		Freq	Control		Freq	Models	OR(95% CI)	p
		Freq	%		Freq	%				
TGGTTCT	HH	5	7.6	0.24	12	9.4	0.26	codominant	0.90(0.57-1.43)	0.67
	H-	22	33.3		43	33.9		dominant	0.91(0.50-1.66)	0.75
	--	39	59.1		72	56.7		recessive	0.79(0.26-2.33)	0.66
TGGTTGT	HH	0	0.0	0.11	2	1.6	0.15	codominant	0.68(0.35-1.34)	0.26
	H-	14	21.2		33	26.0		dominant	0.71(0.35-1.44)	0.34
	--	52	78.8		92	72.4		recessive	0.00(0.00--)	0.99
AAGTTCT	HH	1	1.5	0.15	0	0.0	0.11	codominant	1.57(0.81-3.02)	0.18
	H-	18	27.3		27	21.3		dominant	1.50(0.76-2.96)	0.25
	--	47	71.2		100	78.7		recessive	NA	0.99
TGGCTCT	HH	2	3.0	0.17	2	1.6	0.09	codominant	1.93(1.04-3.59)	0.04
	H-	18	27.3		19	15.0		dominant	2.19(1.09-4.43)	0.03
	--	46	69.7		106	83.5		recessive	1.95(0.27-14.19)	0.51
TGATTCT	HH	0	0.0	0.08	0	0.0	0.06	codominant	1.39(0.60-3.19)	0.44
	H-	11	16.7		16	12.6		dominant	1.39(0.60-3.19)	0.44
	--	55	83.3		111	87.4		recessive	NA	NA
TGGTTCC	HH	0	0.0	0.03	1	0.8	0.07	codominant	0.42(0.14-1.26)	0.12
	H-	4	6.1		16	12.6		dominant	0.42(0.13-1.30)	0.13
	--	62	93.9		110	86.6		recessive	NA	0.99

Freq, frequency; OR, odds ratio; CI, confidence intervals; NA, not applicable; p, p value. Bold characters represent statistically significant values and its structure of haplotype.

using ABI PRISM 3730XL analyzer (PE Applied Biosystems, Foster City, California, USA). Sequence data were analyzed using SeqManII software (DNASTAR Inc., Madison, WI, USA).

Statistical analysis

Hardy-Weinberg equilibrium (HWE) was assessed by SNP stats in both controls and cases [29]. For linkage disequilibrium (LD) block, Haploview version 3.32 was used [30]. Haplotypes and their frequencies was inferred using expectation-maximization (EM) algorithm [31]. Multiple logistic regression model was used to calculate odds ratios (OR), 95% confidence interval, and corresponding p values. While controlling age and gender as co variables, SNP stats, Hap Analyzer version 1.0 [32], and Helixtree (Golden Helix Inc., MT, USA) were used to analyze association of SNPs and haplotypes with DLS.

Results

Of the seven SNPs of *BMP* genes examined, all SNPs were polymorphic. The genotype distributions of all seven SNPs (rs235768, rs1049007, rs3733549, rs17563, rs3734444, rs17557, and rs2231344) were in HWE ($p > 0.05$). Of the SNPs examined, two SNPs (rs235768 and rs1049007) of *BMP2* gene were weakly associated with DLS in the co dominant (rs235768, Arg190Ser, missense, OR = 2.28, 95% CI = 1.19 - 4.36, $p = 0.03$) and dominant model (rs235768, OR = 1.99, 95% CI = 1.07 - 3.72, $p = 0.03$; rs1049007, Ser87Ser, synonymous, OR = 1.93, 95% CI = 1.039 - 3.62, $p = 0.04$).

A haplotype-based association analysis was performed for different combinations of SNPs within *BMP* genes between the DLS group and the control group. The haplotypes were constructed

with cSNPs of *BMP* genes selected for this study [rs235768 (*BMP2*, A/G), rs1049007 (*BMP2*, A/G), rs3733549 (*BMP3*, A/G), rs17563 (*BMP4*, A/G), rs3734444 (*BMP5*, A/G), rs17557 (*BMP6*, G/A), and rs2231344 (*BMP10*, T/C)]. The order of SNPs in the haplotypes was: rs235768, rs1049007, rs3733549, rs17563, rs3734444, rs17557, and rs2231344. Haplotype (TGGCTCT) showed weak association with DLS. The haplotype was associated with DLS in the co dominant (OR = 1.93, 95% CI = 1.04 - 3.59, $p = 0.04$, Table 2) and dominant model (OR = 2.19, 95% CI = 1.09 - 4.43, $p = 0.03$, Table 3). Therefore, the TGGCTCT of *BMP* genes may be a susceptible factor of DLS.

To compare our genotypic results with different ethnic populations, we searched the human SNP database (www.ncbi.nlm.nih.gov/SNP, dbSNP Build 130). Database representing genotype frequencies for the SNPs analyzed in this manuscript is shown in Table 4. The genotype distributions of the control group of all SNPs analyzed in our study are similar to those of Asian population, especially Japanese population (Table 4).

Discussion

Several researches have suggested genetic associations between *BMP* genes and some diseases [26-32]. The association between several SNPs of *BMP2* and *BMP4* genes and genetic hemochromatosis was investigated in the study. A SNP of *BMP2* gene observed in this study (rs235756) was reported to be associated with hemochromatosis penetrance [26]. Wang et al. (2008) reported that SNP of *BMP2* gene (rs1049007 analyzed in our study) was significantly associated with the occurrence of OPLL in the cervical spine [27]. The association of a single SNP of *BMP6* gene (rs3812163) was suggested to have a potential role of *BMP6* in the development of a vascular necrosis in

Table 4: Genotype frequencies of the SNPs of *BMP* genes in each population.

Gene	SNP	Genotype	DLS	Control	Europe	China	Japan	Sub-Saharan African
BMP2	rs235768 (Arg190Ser)	T/T	0.55	0.71	0.57	0.50	0.64	1.00
		T/A	0.42	0.24	0.28	0.48	0.32	0.00
		A/A	0.03	0.06	0.15	0.02	0.04	0.00
BMP2	rs1049007 (Ser87Ser)	G/G	0.56	0.72	No data	No data	No data	No data
		G/A	0.41	0.24				
		A/A	0.03	0.04				
BMP3	rs3733549 (Arg192Gln)	G/G	0.67	0.74	0.90	0.67	0.73	0.88
		G/A	0.32	0.25	0.17	0.33	0.23	0.12
		A/A	0.02	0.01	0.00	0.00	0.04	0.00
BMP4	rs17563 (Val152Ala)	T/T	0.52	0.61	0.15	0.44	0.67	0.65
		T/C	0.41	0.34	0.48	0.47	0.32	0.32
		C/C	0.08	0.05	0.37	0.09	0.01	0.03
BMP5	rs3734444 (Ser37Ser)	T/T	0.74	0.68	0.40	0.71	0.76	0.07
		T/C	0.24	0.30	0.38	0.29	0.22	0.48
		C/C	0.02	0.02	0.22	0.00	0.02	0.45
BMP6	rs17557 (Val368Val)	C/C	0.63	0.61	0.15	0.47	0.71	0.05
		C/G	0.36	0.33	0.45	0.47	0.22	0.35
		G/G	0.01	0.06	0.40	0.06	0.07	0.60
BMP10	rs2231344 (Asp242Asp)	T/T	0.77	0.80	0.82	0.78	0.87	0.47
		T/C	0.23	0.19	0.18	0.20	0.13	0.35
		C/C	0.00	0.01	0.00	0.02	0.00	0.18

From database (<http://www.ncbi.nlm.nih.gov/sites/entrez?db=snp>).

sickle cell disease patients [30]. These genetic studies on the association of SNP of *BMP* genes with many different diseases containing spine disease provided important information for our case-control study with DLS patients. We hypothesized that *BMP* gene polymorphisms might affect the susceptibility of spine diseases such as DLS. We first investigated the genetic association between cSNPs of *BMP* genes and DLS. Given the important biological and genetic functions of *BMP* genes during developmental process, we investigated whether *BMP* gene variations acted as risk factors for DLS in Korean sample. Our results suggested that *BMP* genes may have no involvement in the pathogenesis of DLS. Of all the SNPs and haplotypes analyzed, only two SNPs (rs235768 and rs1049007) showed weak associations with DLS. In addition, haplotype analysis between DLS and control subjects revealed that TGGCTCT of *BMP* genes might be a susceptible factor of DLS, indicating that there is a genetic association between cSNPs of *BMP* genes and DLS.

We compared our genotype frequencies with the human SNP database to show ethnic similarities and differences. The genotype frequencies of our study sample resembled those of Japanese and Chinese Hapmap populations (Table 4). Further studies are needed to elucidate: (i) whether another case-control study with different populations will reveal the association between SNP of *BMP* and DLS; and (ii) whether cSNPs can affect the expression of *BMP* genes. To confirm negative association between *BMP* genes and DLS, replication studies with adequate sample size or an association study with different SNPs not analyzed in this study may be required.

This study has several limitations. First, the sample size may not have been sufficient to detect the associations of smaller effects for DLS. Second, in this study, when we designed the experiment, only cSNPs of *BMP* genes were included for the association study. As a result, our selection of SNPs provided incomplete coverage of currently known common variation in *BMP* genes. Therefore, additional studies with different SNPs, especially promoter SNPs, are needed.

Conclusion

In conclusion, we investigated possible association between SNPs of *BMP* genes and DLS. Our results suggested that cSNPs of *BMP* genes may have influence on the development of DLS in Korean population. However, additional genetic studies are needed to help us understand the precise mechanisms underlying the pathogenesis of DLS.

References

- McPhee IB, Swanson CE . The surgical management of degenerative lumbar scoliosis. Posterior instrumentation alone versus two stage surgery. *Bull Hosp Jt Dis.* 1998; 57: 16-22.
- Grubb SA, Lipscomb HJ, Coonrad RW . Degenerative adult onset scoliosis. *Spine (Phila Pa 1976).* 1988; 13: 241-245.
- Hanley EN Jr . The indications for lumbar spinal fusion with and without instrumentation. *Spine (Phila Pa 1976).* 1995; 20: 143S-153S.
- Sapkas G, Efsthathiou P, Badekas AT, Antoniadis A, Kyrtzoulis J, Meleteas E . Radiological parameters associated with the evolution of degenerative scoliosis. *Bull Hosp Jt Dis.* 1996; 55: 40-45.

5. Tribus CB . Degenerative lumbar scoliosis: evaluation and management. *J Am Acad Orthop Surg.* 2003; 11: 174-183.
6. Bradford DS, Tay BK, Hu SS . Adult scoliosis: surgical indications, operative management, complications, and outcomes. *Spine (Phila Pa 1976).* 1999; 24: 2617-2629.
7. Benner B, Ehni G . Degenerative lumbar scoliosis. *Spine (Phila Pa 1976).* 1979; 4: 548-552.
8. Daffner SD, Vaccaro AR . Adult degenerative lumbar scoliosis. *Am J Orthop (Belle Mead NJ).* 2003; 32: 77-82.
9. Pritchett JW, Bortel DT . Degenerative symptomatic lumbar scoliosis. *Spine (Phila Pa 1976).* 1993; 18: 700-703.
10. Oskoui RJ Jr, Shaffrey CI . Degenerative lumbar scoliosis. *Neurosurg Clin N Am.* 2006; 17: 299-315, vii.
11. Videman T, Gibbons LE, Battié MC, Maravilla K, Vanninen E, Leppävuori J, et al . The relative roles of intragenic polymorphisms of the vitamin d receptor gene in lumbar spine degeneration and bone density. *Spine (Phila Pa 1976).* 2001; 26: E7-7E12.
12. Gómez R, Magaña JJ, Cisneros B, Pérez-Salazar E, Faugeron S, Véliz D, et al . Association of the estrogen receptor alpha gene polymorphisms with osteoporosis in the Mexican population. *Clin Genet.* 2007; 72: 574-581.
13. Canalis E, Economides AN, Gazzerro E . Bone morphogenetic proteins, their antagonists, and the skeleton. *Endocr Rev.* 2003; 24: 218-235.
14. Reddi AH . Interplay between bone morphogenetic proteins and cognate binding proteins in bone and cartilage development: noggin, chordin and DAN. *Arthritis Res.* 2001; 3: 1-5.
15. Attisano L, Wrana JL . Signal transduction by members of the transforming growth factor-beta superfamily. *Cytokine Growth Factor Rev.* 1996; 7: 327-339.
16. Wang EA . Bone morphogenetic proteins (BMPs): therapeutic potential in healing bony defects. *Trends Biotechnol.* 1993; 11: 379-383.
17. Miyazono K, Kusanagi K, Inoue H . Divergence and convergence of TGF-beta/BMP signaling. *J Cell Physiol.* 2001; 187: 265-276.
18. Morse JH, Deng Z, Knowles JA . Genetic aspects of pulmonary arterial hypertension. *Ann Med.* 2001; 33: 596-603.
19. Yanagita M . Modulator of bone morphogenetic protein activity in the progression of kidney diseases. *Kidney Int.* 2006; 70: 989-993.
20. Milet J, Dehais V, Bourgain C, Jouanolle AM, Mosser A, Perrin M, et al . Common variants in the BMP2, BMP4, and HJV genes of the hepcidin regulation pathway modulate HFE hemochromatosis penetrance. *Am J Hum Genet.* 2007; 81: 799-807.
21. Wang H, Yang ZH, Liu DM, Wang L, Meng XL, Tian BP. Association between two polymorphisms of the bone morpho-genetic protein-2 gene with genetic susceptibility to ossification of the posterior longitudinal ligament of the cervical spine and its severity. *Chin Med J (Engl).* 2008; 121:1806-1810
22. Lin JY, Chen YJ, Huang YL, Tang GP, Zhang L, Deng B, et al . Association of bone morphogenetic protein 4 gene polymorphisms with nonsyndromic cleft lip with or without cleft palate in Chinese children. *DNA Cell Biol.* 2008; 27: 601-605.
23. Devaney JM, Tosi LL, Fritz DT, Gordish-Dressman HA, Jiang S, Orkunoglu-Suer FE, et al . Differences in fat and muscle mass associated with a functional human polymorphism in a post-transcriptional BMP2 gene regulatory element. *J Cell Biochem.* 2009; 107: 1073-1082.
24. Ulug P, Vasavda N, Awogbade M, Cunningham J, Menzel S, Thein SL . Association of sickle avascular necrosis with bone morphogenetic protein 6. *Ann Hematol.* 2009; 88: 803-805.
25. Freedman BI, Bowden DW, Ziegler JT, Langefeld CD, Lehtinen AB, Rudock ME, et al . Bone morphogenetic protein 7 (BMP7) gene polymorphisms are associated with inverse relationships between vascular calcification and BMD: the Diabetes Heart Study. *J Bone Miner Res.* 2009; 24: 1719-1727.
26. Zhang P, Shi YH, Wang LC, Chen ZJ . Sequence variants in exons of the BMP-15 gene in Chinese patients with premature ovarian failure. *Acta Obstet Gynecol Scand.* 2007; 86: 585-589.
27. Aebi M . The adult scoliosis. *Eur Spine J.* 2005; 14: 925-948.
28. Rickham, PP. Human experimentation. Code of ethics of the world medical association. Declaration of helsinki. *Br Med J.* 1964; 2: 177.
29. Solé X, Guinó E, Valls J, Iniesta R, Moreno V . SNPStats: a web tool for the analysis of association studies. *Bioinformatics.* 2006; 22: 1928-1929.
30. Barrett JC, Fry B, Maller J, Daly MJ . Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics.* 2005; 21: 263-265.
31. Stephens M, Smith NJ, Donnelly P . A new statistical method for haplotype reconstruction from population data. *Am J Hum Genet.* 2001; 68: 978-989.
32. Jung HY, Park JS, Park YJ, Kim YK, Kimm K, Koh IS. HapAnalyzer: Minimum Haplotype Analysis System for Association Studies. *Genomics & Informatics.* 2004; 2: 107-109.