

## Short Communication

# Is PDL1 Assessment a Prime Time Predictor for Esophageal Cancer Management?

**Paluri RK\***

Assistant Professor of Medicine, University of Alabama at Birmingham, USA

**\*Corresponding author:** Ravi Kumar Paluri, Assistant Professor of Medicine, University of Alabama at Birmingham, USA**Received:** November 05, 2018; **Accepted:** November 26, 2018; **Published:** December 03, 2018

## Short Communication

Programmed Death-Ligand 1 (PD-L1) expression has been emerged as one of the key predictive marker of response to immunotherapy [1]. However, the timing of usage of Checkpoint Inhibitors (CPI) is not in global consensus. In United States, CPI is approved in third line setting, while in Japan it is approved upon progression from initial chemotherapy with no regard to biomarker expression [2]. The recent trial [3] comparing pembrolizumab with chemotherapy (paclitaxel monotherapy) for PDL1 CPS scores >1 did not significantly prolong overall survival (median 9.1 *versus* 8.3 months, HR 0.82, 95% CI 0.66-1.03), and the Objective Response Rate (ORR) was similar (16 *versus* 14 percent). However, the adverse event profile was favorable with pembrolizumab. Therefore, the combination of Ramucirumab and paclitaxel remains the preferred strategy for most patients with advanced oesophageal cancers who have progressed on first line fluoropyrimidine or platinum based regimen [4].

## References

1. Muro K, Chung HC, Shankaran V, Geva R, Catenacci D, Gupta S, et al. Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial. *Lancet Oncol.* 2016; 17: 717-726.
2. Kang YK, Boku N, Satoh T, Ryu MH, Chao Y, Kato K, et al. Nivolumab in patients with advanced gastric or gastro-oesophageal junction cancer refractory to, or intolerant of, at least two previous chemotherapy regimens (ONO-4538-12, ATTRACTION-2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet.* 2017; 390: 2461-2471.
3. Shitara K, Özgüroğlu M, Bang YJ, Di Bartolomeo M, Mandalà M, Ryu MH, et al. Pembrolizumab *versus* paclitaxel for previously treated, advanced gastric or gastro-oesophageal junction cancer (KEYNOTE-061): a randomized, open-label, controlled, phase 3 trial. *Lancet.* 2018; 392: 123-133.
4. Wilke H, Muro K, Van Cutsem E, Oh SC, Bodoky G, Shimada Y, et al. Ramucirumab plus paclitaxel *versus* placebo plus paclitaxel in patients with previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (RAINBOW): a double-blind, randomised phase 3 trial. *Lancet Oncol.* 2014; 15: 1224-1235.