

## Mini Review

# Treating Mitochondrial Myopathy: Can we Boost Oxidative Phosphorylation Activities by Enhancing Mitochondrial Biogenesis?

De Paepe B\*

Department of Pediatrics, Division of Pediatric Neurology and Metabolism & Neuromuscular Reference Centre, Ghent University Hospital, Belgium

\*Corresponding author: Boel De Paepe, Department of Pediatrics, Division of Pediatric Neurology and Metabolism, Ghent University Hospital, Lab Van Coster, UZ 3K5, De Pintelaan 185, 9000 Ghent, Belgium

Received: August 20, 2014; Accepted: October 20, 2014; Published: October 24, 2014

**Abstract**

In clinical practice today, only symptomatic therapeutic approaches are available to patients with mitochondrial myopathies. A possible innovative strategy to ameliorate oxidative phosphorylation deficiency would be to increase the mitochondrial load of the patient's muscle tissue. Some of the most promising stimulators of mitochondrial biogenesis: resveratrol, quercetin, catechins and fibrates, are evaluated in this review. Results so far are promising, but there is need to determine if sufficient mitochondrial proliferation can be achieved to normalize OXPHOS function in patients with mitochondrial myopathy.

**Keywords:** Catechins; Fibrates; Mitochondrial biogenesis; Mitochondrial myopathy; Oxidative phosphorylation deficiency; Quercetin; Resveratrol

**Abbreviations**

EGCG: Epigallocatechin-3-Gallate; MELAS: Mitochondrial Encephalomyopathy, Lactic Acidosis and Stroke-Like Episodes; OXPHOS: Oxidative Phosphorylation; PGC-1 $\alpha$ : Peroxisome Proliferator-Activated Receptor- $\gamma$  Coactivator  $\alpha$ ; PPAR: Peroxisome Proliferation Activated Receptor; SIRT: Sirtuin

**Introduction**

In humans, skeletal muscle is one of the tissues with the highest energetic demand. The majority of the necessary cellular energy is supplied by the mitochondria through Oxidative Phosphorylation (OXPHOS). The OXPHOS system consists of five multiprotein complexes embedded within the inner mitochondrial membrane: complex I (NADH:ubiquinone oxidoreductase), complex II (succinate:ubiquinone oxidoreductase), complex III (ubiquinol:cytochrome c oxidoreductase), complex IV (cytochrome c oxidase) and complex V (ATP synthase). Defects in the OXPHOS system are associated with a broad spectrum of clinical manifestations, not surprisingly often with prominent muscular problems [1]. Muscular complaints range from relatively benign myalgias to severe disabilities. The estimated incidence of OXPHOS defects in humans is approximately 1 in 5,000 live births. OXPHOS defects are complex due to the dual origin of genes involved and the specific nature of the mitochondrial genome. The mitochondrial DNA is present in multiple copies in individual cells, and co-existence of mutated and wild type variants, termed heteroplasmy, can occur. Due to this genetic complexity, pinpointing the origin of the defect is often a time-consuming process that requires thorough biochemical and molecular characterization of patient tissues. Diagnostic testing for OXPHOS deficiencies includes spectrophotometry and blue native-polyacrylamide gel electrophoresis, evaluating the activities of the individual OXPHOS complexes I-V [2].

Under physiological conditions, mitochondrial biogenesis is synchronized with changing energetic demands. The process of


mitochondrial proliferation necessitates the coordinated expression of mitochondrial and nuclear genes and is governed by a complex system of transcription factors and co-activators, of which the peroxisome proliferator-activated receptor- $\gamma$  coactivator  $\alpha$  (PGC-1 $\alpha$ ) seems the master regulator [3,4]. Stimulation of PGC-1 $\alpha$  activity results in increased mitochondrial mass and function. PGC-1 $\alpha$  expression is controlled by the peroxisome Proliferation Activated Receptor (PPAR) family [5], and by deacetylation mediated by NAD<sup>+</sup>-dependent protein deacetylase Sirtuin (SIRT) 1 [6,7].

So far, the development of effective therapies for OXPHOS deficiencies has been extremely limited [8]. A proposed therapeutic approach for OXPHOS patients would be to increase the number of mitochondria per cell, which could result in greater capacities to produce ATP (Figure), higher mitochondrial DNA content and increased mitogenesis has recently been suggested to protect mutation carriers from developing Leber's hereditary optic neuropathy [9]. In cybrids harbouring the classical m.3243A>G mitochondrial encephalomyopathy, lactic acidosis and stroke-like episodes (MELAS) mutation, PGC-1 $\alpha$  overexpression improved OXPHOS complex III and complex IV deficiency [10].

This mini review will focus on some of the most promising stimulators of mitochondrial biogenesis, in relation to their potential to treat OXPHOS deficiencies.

**Exercise**

Physical activity results in increases in both mitochondrial content and function in skeletal muscle tissue [11], typically doubling the amount of muscle mitochondria. In elderly subjects, 8 weeks of high-intensity exercise training increased expression of PPAR $\gamma$ -1 $\alpha$  and various OXPHOS components, and resulted in increased exercise endurance [12]. Significantly increased citrate synthase and OXPHOS complex I, III and IV activities were described in vastus lateralis of young and elderly healthy subjects [13]. A study monitoring patients with heteroplasmic mitochondrial DNA defects showed that, in



**Figure: Rationale for treating oxidative phosphorylation (OXPHOS) deficiencies by stimulating mitochondrial biogenesis.** Therapeutic measures stimulating mitochondrial proliferation include exercise and administering mitogenetic compounds, resveratrol, quercetin, catechins and fibrates. In theory, they enhance the OXPHOS capacities, but this needs to be proven in a therapeutic setting.

response to exercise, levels of the mitochondrial marker citrate synthase increased 50%, while OXPHOS activities went up 20% on average [14], resulting in improved muscle symptoms [15]. In a cohort of patients with chronic progressive external ophthalmoplegia due to single, large-scale mtDNA deletions, 12 weeks of resistance training led to increased muscle strength and a 2-fold significant increase in satellite cell numbers [16]. These data offer promising, yet somewhat unequivocal findings, as different types of exercise were performed and the number of patients enrolled often was low. Although exercise training is regarded a highly efficient means of increasing muscle function, there is an important drawback. Many patients have difficulties maintaining a regular exercise program, which makes other strategies involving nutritional supplements and drugs an attractive alternative.

### Caloric restriction

Myriad animal studies have taught us that caloric restriction has the potential to prolong lifespan and diminish aging-associated changes. In the mechanisms responsible for these beneficial effects, increased mitochondrial biogenesis and activity are important players [17]. Caloric restriction increased PGC-1 $\alpha$  expression significantly in the gastrocnemius of rats [18]. In a cohort of healthy overweight patients, 25% caloric deficit was shown to increase mitochondrial content and associated markers of biogenesis in vastus lateralis, however, without changing OXPHOS complex IV activity [19]. It needs said that in human due to poor compliance, implementing caloric restriction is not a reliable therapeutic approach.

### Resveratrol

Resveratrol (3,5,4'-trihydroxystilbene) is a natural polyphenol present in the skin of red grapes presumed to boost mitogenesis by activating PGC-1 $\alpha$  via SIRT1 [20,21]. Significantly increased mitochondrial biogenesis was shown in resveratrol fed mice [21], an effect comparable to the one achieved in PGC-1 $\alpha$  overexpressing mice [22]. An *in vitro* study assaying the effect of resveratrol in cultured skin fibroblasts from patients with OXPHOS complex I and complex IV deficiency showed stimulated mitochondrial biogenesis and increased OXPHOS protein levels, resulting in corrected deficiency (measured through oxygen uptake rates) in 6 out of 16 patients' cell lines [23]. In our own *in vitro* study however, resveratrol was unable to normalize mitochondrial enzyme activities in a series of fibroblast cell lines from patients with isolated complex II and complex IV deficiencies, and seemed to necessitate a threshold of residual OXPHOS activity to generate a response [24]. At this time, the first results are beginning to surface addressing the effects of resveratrol in patients. A study evaluating the benefits of exercise in elderly healthy subjects showed no additional gain of adding resveratrol to the study protocol [12].

### Quercetin

Quercetin (2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4H-chromen-4-one) is a widely distributed plant pigment that gets its name from the oak tree (*quercus*). It is present in many foods, including red onions, apples and berries [25]. Seven days of quercetin feedings significantly increased a set of mitochondrial markers in murine soleus muscle, among which PGC-1 $\alpha$  and SIRT1 expression, and resulted in increased maximal endurance capacity and voluntary wheel-running activity [26]. In humans four-week quercetin treatment achieved a small but significant performance effect in young healthy adults [27]. Combined with additional antioxidants and caffeine, a six-weeks supplement has been shown to improve endurance in a bicycle test [28].

### Catechins

Epigallocatechin-3-Gallate (EGCG) is known for its anti-cancer and mitogenetic properties. It is present in large quantities in tea leaves and protects mitochondria from oxidative stress. The reduced OXPHOS complex I and complex V activities observed in EBV-immortalized lymphoblasts and fetal skin fibroblasts from subjects with Down's syndrome, were restored to normal levels by treating the cells *in vitro* with 20 $\mu$ M EGCG for 24h [29].

### Fibrates

Fibrates are amphipathic carboxylic acids that act as PPAR agonists. They are routinely used for treating diabetes mellitus and metabolic syndrome. Bezafibrate was reported to boost residual OXPHOS enzyme activity that lead to normal oxygen consumption rates in the treated cells [30]. Four-day bezafibrate treatment of cybrids carrying homoplasmic mitochondrial tRNA gene mutations resulted in increased PGC-1 $\alpha$  mRNA expression and ATP synthesis. It also increased activities of citrate synthase and OXPHOS complex IV [31].

### Conclusion

Based on the literature reviewed, one can conclude that stimulators of mitochondrial biogenesis could be of benefit to patients with

inborn OXPHOS deficiencies. However, this remains to be tested thoroughly in humans. Also, development of potent derivatives of the known mitogenetic compounds could increase therapeutic responses further.

## References

- Zeviani M, Di Donato S. Mitochondrial disorders. *Brain*. 2004; 127: 2153-2172.
- Smet J, Seneca S, De Paepe B, Meulemans A, Verhelst H, Leroy J, et al. Subcomplexes of mitochondrial complex V reveal mutations in mitochondrial DNA. *Electrophoresis*. 2009; 30: 3565-3572.
- Rodgers JT, Lerin C, Haas W, Gygi SP, Spiegelman BM, Puigserver P. Nutrient control of glucose homeostasis through a complex of PGC-1alpha and SIRT1. *Nature*. 2005; 434: 113-118.
- Scarpulla RC. Metabolic control of mitochondrial biogenesis through the PGC-1 family regulatory network. *Biochim Biophys Acta*. 2011; 1813: 1269-1278.
- Wenz T, Diaz F, Spiegelman BM, Moraes CT. Activation of the PPAR/PGC-1alpha pathway prevents a bioenergetic deficit and effectively improves a mitochondrial myopathy phenotype. *Cell Metab*. 2008; 8: 249-256.
- Rodgers JT, Lerin C, Gerhart-Hines Z, Puigserver P. Metabolic adaptations through the PGC-1 alpha and SIRT1 pathways. *FEBS Lett*. 2008; 582: 46-53.
- Gerhart-Hines Z, Rodgers JT, Bare O, Lerin C, Kim SH, Mostoslavsky R, et al. Metabolic control of muscle mitochondrial function and fatty acid oxidation through SIRT1/PGC-1alpha. *EMBO J*. 2007; 26: 1913-1923.
- Pfeffer G, Majamaa K, Turnbull DM, Thorburn D, Chinnery PF. Treatment for mitochondrial disorders. *Cochrane Database Syst Rev*. 2012; 4: CD004426.
- Giordano C, Iommarini L, Giordano L, Maresca A, Pisano A, Valentino ML, et al. Efficient mitochondrial biogenesis drives incomplete penetrance in Leber's hereditary optic neuropathy. *Brain*. 2014; 137: 335-353.
- Srivastava S, Diaz F, Iommarini L, Aure K, Lombes A, Moraes CT. PGC-1 alpha/beta induced expression partially compensates for respiratory chain defects in cells from patients with mitochondrial disorders. *Hum Molec Genet*. 2009; 18: 1805-1812.
- Jacobs RA, Lundby C. Mitochondria express enhanced quality as well as quantity in association with aerobic fitness across recreationally active individuals up to elite athletes. *J Appl Physiol (1985)*. 2013; 114: 344-350.
- Olesen J, Gliemann L, Biensø R, Schmidt J, Hellsten Y, Pilegaard H. Exercise training, but not resveratrol, improves metabolic and inflammatory status in skeletal muscle of aged men. *J Physiol*. 2014; 592: 1873-1886.
- Andersen SP, Thøgersen F, Sveen ML, Ormgreen MC, Kragh T, Vissing J, et al. Effect of aerobic training and detraining on mitochondrial function in aging. *J Neuromusc Dis*. 2014; 1: S94.
- Taivassalo T, Shoubridge EA, Chen J, Kennaway NG, DiMauro S, Arnold DL, et al. Aerobic conditioning in patients with mitochondrial myopathies: physiological, biochemical, and genetic effects. *Ann Neurol*. 2001; 50: 133-141.
- Taivassalo T, Haller RG. Exercise and training in mitochondrial myopathies. *Med Sci Sports Exerc*. 2005; 37: 2094-2101.
- Murphy JL, Blakely EL, Schaefer AM, He L, Wyrick P, Haller RG, et al. Resistance training in patients with single, large-scale deletions of mitochondrial DNA. *Brain*. 2008; 131: 2832-2840.
- López-Lluch G, Hunt N, Jones B, Zhu M, Jamieson H, Hilmer S, et al. Calorie restriction induces mitochondrial biogenesis and bioenergetic efficiency. *Proc Natl Acad Sci U S A*. 2006; 103: 1768-1773.
- Baker DJ, Betik AC, Krause DJ, Hepple RT. No decline in skeletal muscle oxidative capacity with aging in long-term calorically restricted rats: effects are independent of mitochondrial DNA integrity. *J Gerontol A Biol Sci Med Sci*. 2006; 61: 675-684.
- Civitarese AE, Carling S, Heilbronn LK, Hulver MH, Ukropcova B, Deutsch WA, et al. Calorie restriction increases muscle mitochondrial biogenesis in healthy humans. *PLoS Med*. 2007; 4: e76.
- Biala A, Tauriainen E, Siltanen A, Shi J, Merasto S, Louhelainen M, et al. Resveratrol induces mitochondrial biogenesis and ameliorates Ang II-induced cardiac remodelling in transgenic rats harbouring human renin and angiotensinogen genes. *Blood Pressure*. 2010; 19: 196-205.
- Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C, Daussin F, et al. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell*. 2006; 127: 1109-1122.
- Viscomi C, Bottani E, Civiletto G, Cerutti R, Moggio M, Fagioli G, et al. *In vivo* correction of COX deficiency by activation of the AMPK/PGC-1α axis. *Cell Metab*. 2011; 14: 80-90.
- Lopes Costa A, Le Bachelier C, Mathieu L, Rotig A, Boneh A, De Lonlay P, et al. Beneficial effects of resveratrol on respiratory chain defects in patients' fibroblasts involve estrogen receptor and estrogen-related receptor alpha signaling. *Hum Molec Genet*. 2014; 23: 2106-2119.
- De Paepe B, Vandemeulebroecke K, Smet J, Vanlander A, Seneca S, Lissens W, et al. Effect of resveratrol on cultured skin fibroblasts from patients with oxidative phosphorylation defects. *Phytother Res*. 2014; 28: 312-316.
- Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of *in vivo* toxicity, including lack of genotoxic/carcinogenic properties. *Food Chem Toxicol*. 2007; 45: 2179-2205.
- Davis JM, Murphy EA, Carmichael MD, Davis B. Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am J Physiol Regul Integr Comp Physiol*. 2009; 296: R1071-1077.
- Nieman DC, Williams AS, Shanely RA, Jin F, McNulty SR, Triplett NT, et al. Quercetin's influence on exercise performance and muscle mitochondrial biogenesis. *Med Sci Sports Exerc*. 2010; 42: 338-345.
- MacRae HS, Mefferd KM. Dietary antioxidant supplementation combined with quercetin improves cycling time trial performance. *Int J Sport Nutr Exerc Metab*. 2006; 16: 405-419.
- Valenti D, De Rasmio D, Signorile A, Rossi L, de Bari L, Scala I, et al. Epigallocatechin-3-gallate prevents oxidative phosphorylation deficit and promotes mitochondrial biogenesis in human cells from subjects with Down's syndrome. *Biochim Biophys Acta*. 2013; 1832: 542-552.
- Bastin J, Aubey F, Rotig A, Munnich A, Djouadi F. Activation of peroxisome proliferator-activated receptor pathway stimulates the mitochondrial respiratory chain and can correct deficiencies in patient's cells lacking its components. *J Clin Endocrinol Metab*. 2008; 93: 1433-1441.
- Wenz T, Wang X, Marini M, Moraes CT. A metabolic shift induced by a PPAR panagonist markedly reduces the effects of pathogenic mitochondrial tRNA mutations. *J Cell Mol Med*. 2011; 15: 2317-2325.