

Special Article - Antioxidants in Foods

Effects of Agronomic and Oil Processing Conditions on Natural Antioxidative Phenolics in Olive (*Olea europaea* L.)

Ye JH^{1*}, Wijesundera C² and Shi M¹¹Department of Tea Science, Zhejiang University, China²CSIRO Animal, Food and Health Sciences, Australia***Corresponding author:** Jian-Hui Ye, Zhejiang University Tea Research Institute, Hangzhou 310058, China, Email: jx0515@163.com**Received:** November 05, 2014; **Accepted:** November 17, 2014; **Published:** November 18, 2014**Abstract**

Olive fruits contain appreciable amounts of phenolic compounds which are valuable antioxidants and bioactives. Secoiridoids, such as oleuropein, demethyloleuropein, oleuropein aglycone and ligstroside are the predominant classes of phenolic compounds in intact olive fruits, while 4-Hydroxyphenylethanol (4-HPEA), 3,4-Dihydroxyphenylethanol (3,4-DHPEA), the dialdehydic form of decarboxymethyl elenolic acid linked to 3,4-DHPEA (3,4-DHPEA-EDA) and oleuropein aglycone (3,4-DHPEA-EA) are the principal phenolic constituents of Virgin Olive Oil (VOO). The phenolic compounds originally present in the olive fruit are substantially changed by the conditions of oil processing (crushing, malaxation and phase separation) and distributed among the final product (VOO) and by-products in various ways. They may undergo further changes during storage. This paper reviews the effects of the processing steps on the transfer of phenolic compounds from intact fruit to the final product (VOO), as well as the effects of agronomic factors (cultivar, maturity and irrigation) on the phenolic constituents of olive fruits at harvest.

Keywords: Olives; Olive oil; Phenols; Milling; Extraction/separation**Introduction**

The olive tree (*Olea europaea* L.) is a small tree that grows natively in the tropical and subtropical regions of the world, such as the eastern Mediterranean Basin, the coastal areas of south Eastern Europe, northern Africa and Australia. As a member of the family *Oleaceae*, the olive tree is best known for its fruit which is the primary source of olive oil. The olive tree has a long history of nutritional and medicinal uses in the Mediterranean region, and olive leaves and olive-leaf extracts in particular have been claimed to have antioxidant, antimicrobial and cardio protective properties which have been attributed to their phenolic constituents [1,2].

It has been widely acknowledged that consumption of olive oil exerts benefits on human health, including reducing the risk coronary heart disease, preventing several types of cancers, as well as enhancing immune and inflammatory responses [3-5]. This is evidenced by a reduced incidence of cardiovascular diseases and certain cancers (e.g. breast cancer and colorectal cancer) in the traditional "Mediterranean diet" countries in which olive oil is the major source of dietary fat, in comparison with other regions [4,6]. The nutritional and medicinal properties of olive oil have been attributed to the high content of monounsaturated fatty acids and the presence of functional bioactives such as oleic acid, tocopherols, carotenoids, phospholipids and phenolic compounds [7,8]. Recently, it has become increasingly evident that Virgin Olive Oil (VOO), which is the main source of fat in the Mediterranean diet, is more than a monounsaturated-rich source, and its phenolics could account for much of its nutrigenomic protective effects of the Mediterranean diet [3,5,9]. Numerous studies testify the physiological functions of olive phenolic isolates *in vivo* and *in vitro* [10-12]. In addition to the beneficial role in human


health, the olive phenolics impart unique sensory characteristics to the oils and enhance oil stability [13].

The total concentration of phenolic compounds in Virgin Olive Oil (VOO) generally ranges from 40 to 1000 mg/kg [14], with various concentrations of individual phenolic compounds being present. These phenolic compounds in VOO originate from those originally present in the olive fruits used for oil extraction but are substantially modified depending on the conditions used for oil extraction and subsequent storage [15]. The phenolic composition of olive fruit at harvest is influenced by agronomic factors [16]. Only a small fraction (approximately 2%) of the phenolic compounds present in the olive fruit is transferred to the extracted oil, the rest (98%) is either retained in the pulp or washed off with the wastewater [17,18]. Yet, the phenolic composition of the olive fruit at harvest is probably the most important variable involved in the phenolic composition of the final VOO [19].

As the type and concentration of phenolic compounds present in VOO are key determinants of both the bioactivity and sensory characteristics of the oil, it is important to understand how these compounds are impacted by agronomic conditions used to grow the fruit as well as the processing conditions used for oil extraction. Whilst, several papers have been published reviewing the effects of some of these variables often on the total phenolic concentration of VOO [19], few papers are available on the effect of each variable on individual phenolic components. Additionally, the transformations involving individual phenolics during oil extraction and their transfer routes to the final product (VOO) are still unclear. This review aims to fulfil this gap by discussing the transfer of phenolics from olive fruit to VOO by dividing the transfer path into two segments:

Table 1: The main classes of phenolic compounds of olive fruit.

Class	Phenolic compounds	Reference
Phenolic acids	Chlorogenic acid, Caffeic acid, 4-Hydroxybenzoic acid, Protocatechuic acid, Vanilic acid, Syringic acid, 4-Coumaric acid, 2-Coumaric acid, Ferulic acid, Sinapic acid, Benzoic acid, Cinnamic acid, Gallic acid, Caffeoylquinic acid, 3,4-Dihydroxyphenyl-propionic acid	[14,21,61]
Phenolic alcohols	3,4-Dihydroxyphenylethanol (3,4-DHPEA), 4-Hydroxyphenylethanol (4-HPEA), Homovanillyl alcohol (the metabolite of 3,4-DHPEA)	[22,45,55]
Flavonoids or the glucosides	Quercetin, Luteolin, Apigenin, Chrysoeriol, Rutin, Quercetin-3-O-rutinoside, Luteolin-7-O-glucoside, Luteolin-5-O-glucoside, Apigenin-7-O-glucoside, Apigenin-7-O-rutinoside, Apigenin-6,8-glucoside, Chrysoeriol-7-O-glucoside, Luteolin-4-glucoside	[21,55,61]
Secoiridoids or the hydrolytic derivatives	Oleuropein, Ligstroside, Nüzhenide, Elenolic acid, Demethyloleuropein, Oleuropein aglycone (3,4-DHPEA-EA), the dialdehydic form of decarboxymethyl elenolic acid linked to 3,4-DHPEA (3,4-DHPEA-EDA), Ligstroside aglycone (4-HPEA-EA), the dialdehydic form of decarboxymethyl elenolic acid linked to 4-HPEA (4-HPEA-EDA)	[14,22,24]
Anthocyanins	Cyanidin-3-O-glucoside, Cyanidin-3-O-rutinoside, Cyanidin-3-caffeylglucoside, Cyanidin-3-caffeylaltinoside, Delphinidin 3-rhamosylglucoside-7-xyloside	[14,22,24]
Hydroxycinnamic acid derivatives	Verbascoside	[70]




preharvest period of olive fruit and postharvest oil processing. In this approach, to the extent possible, the effects of the agronomic factors (olive cultivar, maturity and irrigation) and oil processing variables (crushing, malaxation, phase separation and storage conditions) on individual phenolic components are reviewed.

Phenolics in olive fruit and VOO

Olive fruit generally contains 50% water, 1.6% protein, 22% oil, 19.1% carbohydrate, 5.8% cellulose, 1.5% inorganic substances and 1-3% phenolic compounds [20]. Phenolic compounds in olive fruits are primarily classified into phenolic acids, phenolic alcohols, flavonoids and the glucosides, secoiridoids and the hydrolytic derivatives, anthocyanins as well as hydroxycinnamic acid derivatives (Table 1).

Secoiridoids are the predominant phenolic compounds in olive fruits which are exclusive to the plants of the *Oleaceae* family (Figure 1) [14]. Secoiridoids are characterised by the presence in their molecules of elenolic acid (EA, Figure 1 *a*) or its derivatives linked to 3, 4-dihydroxyphenylethanol (3,4-DHPEA/hydroxytyrosol, Figure 1 *b*) or 4-hydroxyphenylethanol (4-HPEA/tyrosol, Figure 1 *c*). Oleuropein (Figure 1 *d*), the 3,4-DHPEA ester of β -glucosylated elenolic acid, is an important secoiridoid mainly responsible for the bitterness of olive fruits [21]. Other major olive phenolics include demethyloleuropein (Figure 1 *e*), oleuropein aglycone (3,4-DHPEA-EA, Figure 1 *f*), ligstroside (Figure 1 *g*), ligstroside aglycon (4-HPEA-EA, Figure 1 *h*), the dialdehydic form of decarboxymethyl elenolic acid linked to 3,4-DHPEA (3,4-DHPEA-EDA/oleacein, Figure 1 *i*), and the related 4-HPEA-EDA (Figure 1 *j*) and nüzhenide



(Figure 1 k) [22-24]. The most abundant secoiridoids in intact olives are oleuropein, demethyloleuropein, ligstroside, nüzhenide [25]. Oleuropein and demethyloleuropein are found mainly in the pericarp, whereas nüzhenide is found exclusively in the seed [25]. While, oleuropein is always present in the drupes of all the olive cultivars, demethyloleuropein and verbascoside can be cultivar-dependent [26].

Phenolic compounds are transferred to oil during processing and contribute to the sensory characteristics (bitter, astringent and pungent taste), stability and health benefits of olive oil. VOO contain higher levels of phenolics than refined olive oil due to losses occurring during oil refining [27]. The derivatives of the major secoiridoid constituent of olive fruit, oleuropein, form the majority of the phenolic fraction in the oil: 3,4-DHPEA, 3,4-DHPEA-EDA and 3,4-DHPEA-EA [2,29,30]. 4-HPEA, 3,4-DHPEA and 3,4-DHPEA-EA have been reported to constitute approximately 90% of VOO phenolics [27,31]. Notably, oleuropein and demethyloleuropein are absent in VOO, and there are two plausible explanations for this: (a) hydrolysis of the corresponding glucosides induced by endogenous β -glucosidases during the crushing and malaxation processes [32]; (b) the low solubility of these glucosidic substances in the oil phase [26].

Phenolic acids, such as caffeic, vanillic, syringic, 4-coumaric, 2-coumaric, protocatechuic, sinapic and 4-hydroxybenzoic acids, and flavonoids, such as luteolin, apigenin and taxifolin, have been also detected in VOO [29,33]. Luteolin and apigenin are the two main flavonoids present in VOO, which originate from their corresponding glucosides present in the olive fruit [34]. A new class of phenolic compounds 3,4-DHPEA-isochromans has been reported to occur in VOO [35], but it is still unclear whether they are natural compounds present in olive fruits or products formed during the technological “kneader” process or storage of oil.

Effects of agronomic factors on the phenolic composition of olive fruit

Olive cultivar: There can be qualitative and quantitative differences in the phenolic composition of olive fruit depending upon the cultivar and where they are grown (country / region) [36]. The concentration of oleuropein has been observed to vary with the type of olive cultivar [37-39]. The Greek and Italian cultivars contain greater amounts of oleuropein in comparison with the Spanish and Portuguese cultivars [40], and slow-ripening cultivars contain more oleuropein and its aglycone than fast-ripening cultivars [23]. In two cultivars grown in the south of Tunisia, oleuropein was abundant in the *Gemri-Dhokar* cultivar (61.04 mg/100 g of fresh olive) but occurred at relatively much lower concentrations in the *Dhokar* cultivar (0.25 mg/100 g of fresh olive) according to HPLC analysis

[41]. This explained the distinctive sweet taste of the *Dhokar* cultivar. A similar result was reported by Jemaiet *al.* where *Dhokar* cultivar contained much lower concentrations of phenolics compared to *Chemlali* cultivar, being 0.16 g Gallic Acid (GA) equivalents/kg and 3.75g GA equivalents/kg of fresh olive respectively at harvest [42]. *Koroneiki* cultivar, an indigenous Greek olive variety, is thought to be resistant to verticillium wilt by involvement of phenolic compounds in the defense mechanism compared with the susceptible *Amfissis* cultivar [43].

3,4-DHPEA and oleuropein were the most abundant phenolic compounds identified in 18 different olive cultivars from north and central Portugal, where rutin and luteolin 7-O-glucoside were the two main flavonoids [44]. 3,4-DHPEA has been reported as the major phenolic compound in the olive cultivars *Ayvalık*, *Domat* and *Gemlik* where its concentration generally exceeds that of oleuropein [37]. Demethyloleuropein only occurs in the cultivars of *Coratina* and *Leccino* and not other Italian cultivars. Indeed, this compound has been recommended as a potential marker for these particular cultivars [45]. Large differences in the concentration of verbascoside have been observed in the olive cultivars of *Picual* (Spain), *Frantoio* (Italy), *Salonenque* (France) and *Galega* (Portugal), raising the prospect of using this compound as a marker for differentiating these cultivars harvested at the same maturity [36]. Similarly, the ratio of anthocyanins to anthocyanins derivatives has been proposed as a parameter to differentiate five Italian cultivars, namely *Frantoio*, *Rossellino*, *Cilegino*, *Cuoricino*, and *Grossolana* [22]. A new phenolic compound 3-O-galloyl quinic acid butyl ester has been reported from the fruit pulp of a Chinese olive variety, but it is still uncertain whether this compound is unique to this variety [46].

Maturity of olives

The maturation stages of olive fruits represented by green, spotted, purple and black fruit colours correspond to the concentration of anthocyanins [47]. Differences in the phenolic composition between mature and immature fruits have been attributed to chemical and enzymatic reactions occurring during the ripening process.

Oleuropein and ligstroside which are the major secoiridoids constituents of unripe olive fruit have been reported to decrease during ripening and is almost undetectable in dark olive fruits [48,49]. This is accompanied by increases in the concentrations of 3,4-DHPEA, 4-HPEA, and verbascoside [50], as well as that of the nonphenolic compound elenolic acid glucoside [51]. The degradation of oleuropein and ligstroside as shown in Figure 2 could be due to the increased activity of hydrolytic enzymes and esterase during ripening [42,50], producing oleuropein aglycone and glucose by β -glucosidase enzyme hydrolysis followed by degradation into elenolic acid and 3,4-DHPEA by esterase [42]. Since oleuropein and ligstroside degrade into aglycones during ripening, it has been suggested that elenolic acid glucoside and 3,4-DHPEA might serve as potential indicators of olive maturity [45]. However, in *Chétoui* cultivar, oleuropein and 3,4-DHPEA in olive fruit exhibited the same trends during maturation, and the concentration of oleuropein was not inversely correlated with the concentration of 3,4-DHPEA [52]. Polymerization of oleuropein might explain the decrease in oleuropein concentration with the formation of phenolic oligomers [21], as evidenced by the detection of oleuropein trimers in *Chétoui* olive extracts at harvest

[51]. Similarly, changes in the concentration of phenolic acids, such as *trans*-cinnamic acid, sinapic acid, 4-coumaric acid and syringic acid, exhibit cultivar based differences during ripening [53].

An apparent decrease in the concentrations of sugars and phenolics during maturation of olive fruit has been reported [39], which might be attributed to a reaction between the two [54]. Flavonoid glucosides, mainly luteolin 7-*O*-glucoside, initially increased during green maturation and later declined, accompanied by an increase in the concentration of the flavonoid aglycones at the last stage of ripening [55]. L-Phenylalanine Ammonia Lyase (PAL) is the key enzyme in phenolic biosynthesis. A decrease in enzymatic activity of PAL is a plausible explanation for the reduction in total phenolic content of olive drupes during ripening [37,39]. In contrast to the above observations, Bouaziz *et al.* reported that the olive cultivar *Chemlali* presented the highest concentration of phenolics at the last stage of maturation [55].

Irrigation

In addition to the cultivars and maturity, irrigation also affects the composition of phenolic compounds in olive fruits [56]. Irrigation had negligible effects on free acidity, peroxide value, and fatty acid composition of VOO, but strongly influenced its phenolic concentration [57]. It is generally accepted that phenolic substances are more abundant in drought-stressed olive crops than in irrigated crops, which might be due to drought-related variation in the enzymatic activities involved in the formation of phenolic substances. It has been reported that the activity of the enzyme PAL in olive fruit decreased with increased irrigation [58,59].

Machado *et al.* observed that the phenolic content, PAL and antioxidant activity of olive fruits of the Portuguese cultivar *Cobrançosa* were significantly affected by the extent of water treatments applied, giving 61.3 to 15.7 g GA equivalents/kg under rain fed conditions (no irrigation), 24.5 to 11.2 g GA equivalents/kg under inadequate irrigation and 14.7 to 10.2 g GA equivalents/kg under good irrigation [56]. Providing irrigation from the second half of the fruit growing cycle (i.e. after pit hardening) can delay the ripening process leading to changes in the primary and secondary metabolisms, resulting in marked changes in the phenolic composition at harvest [60]. Thus, irrigation exerts a negative effect on the quality of VOO by way of diminished phenolic concentrations.

Transfer of phenolic compounds from olive fruit to VOO

The extraction process for VOO production comprises four main stages: crushing, malaxation, separation and clarification (Figure 3). During oil extraction, phenolic compounds in olive fruit are transferred from crushed fruit (paste) to olive oil and by-products (pomace and wastewater). The amount of phenolic compounds transferred from fruit to oil is influenced by the solubility of these compounds in the oil phase and the presence of surfactants and temperature [18]. In addition, reactions such as breakdown of glycosides and oligosaccharides by glucosidases, oxidation by phenol oxidases, and polymerization of free phenols result in modification of olive phenolics [61,62].

Crushing

Crushing is performed to break down cell walls and release oil droplets within the cells, which can also trigger several enzymatic

reactions resulting in the conversion of phenolic compounds into various products as well as generation of volatile flavor compounds. Thus, depending on the crushing conditions, the concentration of minor components intimately related to the taste, aroma and stability of VOO are modified. Generally, the stronger the crushing conditions, the higher the concentration of phenolic compounds in both olive paste and oil. This could be attributed to greater disintegration of tissue as well as activation of enzymes such as β -glucosidase [62]. Consequent enzyme activity results in the generation of aglycones such as 3,4-DHPEA-EDA and 4-HPEA-EDA. 3,4-DHPEA-EDA could be obtained from the hydrolysis of oleuropein, demethyloleuropein, and ligstroside [19]. A large fall in the oleuropein content of olive paste has been observed [26,63], suggesting extensive degradation/conversion during the crushing operation. In addition to enzymatic reactions, phenolic compounds may also be complex with several polysaccharides impairing their access to the oil phase during crushing and malaxation [64].

Malaxation

Malaxation induces coalescence of small oil droplets into larger droplets and subsequent formation of a continuous oil phase, and promotes release of phenolic compounds into oil phase. And the activation of endogenous β -glucosidases during crushing can catalyze the hydrolysis of oleuropein, demethyloleuropein and ligstroside leading to the generation of their aglyconic forms [65]. Both malaxation temperature and time could influence the phenolic composition of VOO [17,66,68]. Interactions between polysaccharides and phenolic compounds present in the olive pastes may also be involved in the loss of phenols during malaxation process [26].

In experiments performed under laboratory conditions, the concentration of hydrophilic phenols (simple phenols, secoiridoids and lignans) in oil increased with the malaxation temperature attaining a maximum at 27 °C [69]. The concentration of the simple phenols increased linearly up to 36°C which was the maximum malaxation temperature examined, and the concentration of lignans was virtually unaffected. Stefanoudaki *et al.* also found a large increase in the phenolic concentration of VOO when the malaxation temperature was raised between 15 and 42 °C for two popular Italian (cv. *Coratina*) and Cretan (cv. *Koroneiki*) varieties [68]. On the other hand, other studies have shown falls in the phenolic content when the malaxation temperature was increased beyond 27 °C [17,69]. This fall




Figure 3: Unit operations in olive oil extraction and their impact on phenolic compounds.

in the phenolic concentration at higher malaxation temperatures has been attributed to oxidative degradation of secoiridoids resulting from enhanced activity of native oxidoreductases, e.g. Polyphenoloxidases (PPO) and Peroxidases (POD) and Lipoxygenases (LOX) [70]. Hydrolysis of oleuropein and demethyloleuropein to their corresponding aglycones by β -glucosidase has also been suggested as a reason for the fall in phenolic concentration at higher malaxation temperatures [17].

The opinion that malaxation temperatures higher than 27 °C lowers the phenolic concentration and thereby the quality of VOO has been brought in to question by more recent studies [66,71]. The terms 'first cold pressing' and 'cold extraction' signifying better quality VOO can only be used for oils extracted below 27 °C. However, VOO produced in an industrial plant using a malaxation temperature of 35 °C has been shown to contain more phenolic compounds, greater storage stability and better sensory properties than oil produced at 25 °C [66]. Taticchi *et al.* reported that whilst POD was essentially stable up to 40 °C, PPO was thermally labile with low thermal stability even at 40 °C [71]. Other contributory factors currently not well understood are the influence of malaxation temperature on the release of phenolics from cell wall polysaccharides by endogenous hemicellulases [64], and polygalacturonases and the solubilisation of the released phenolics in VOO [71].

Malaxation time also influences the phenolic content of VOO. In general, the phenolic concentration decreases as the malaxation time is increased. This is probably due to oxidative degradation of the phenolics catalysed by PPO, POD and LOX which increases with the exposure time of olive paste to air. Ranalli *et al.* recommended that the malaxation time should not exceed 45 min in order to preserve the phenolics [72]. *Chemlali* and *Chetoui* VOOs presented the highest total phenolic concentrations equal to 230.58 and 828.72 mg GA equivalents/kg at malaxation time of 30 min while the lowest concentrations at 60 min. In addition, the composition of individual phenolic compounds was differentially influenced by malaxation time [73]. Artajo *et al.* reported that malaxation time had an important effect on the alcohols and secoiridoids: the concentration of 3,4-DHPEA and 4-HPEA decreased and their hydrophilic character improved through their presence in the wet pomace and wastewater; the concentrations of demethyloleuropein and oleuropein decreased significantly after 45 min of malaxation, which indicated the beginning of the degradation process of secoiridoid compounds or their transformation into secoiridoid derivatives, mainly 3,4-DHPEA-EDA [74]. Jiménez *et al.* also reported that the malaxation time of olive paste was negatively correlated with the concentrations of secoiridoid as well as 3,4-DHPEA and 4-HPEA, but had little effect on the concentrations of phenolic acids, flavonoids and lignan [71]. Nevertheless, the effects of malaxation time on the phenolic content of VOO are minor compared to the effects of malaxation temperature [17,63,68].

Phase separation

Commercial VOO production is performed using either a classical mill (batch process) or a centrifugal extractor (continuous process). The classical mill, used for centuries, has been almost replaced by centrifugal extractors because of the productivity gains it offers [68]. The centrifugal process can be either a two-phase or

three-phase separation system. The two-phase system, introduced relatively more recently, requires a minimal moisture value on the olive paste (50%) to facilitate the separation process [76]. This system yields higher levels of phenolics, especially 3,4-DHPEA and 4-HPEA, and more stable oil compared to three-phase systems [77,78]. Klen and Vodopivec reported that the phase separation method only impacts on the quantitative recovery of phenolics and had no obvious qualitative effect on the phenolic composition [79]. In contrast to the three-phase systems, the two-phase extractors do not produce waste water and produce VOO containing higher concentrations of phenolics as well as flavour compounds [80].

Storage

Compared to other edible oils, VOO is reputed to be stable to oil processing and storage conditions [32]. This has been attributed in part to the antioxidant effects of the phenolic compounds present in VOO [20]. It has been shown that the concentrations of total phenolics and the oleosidic forms of 3,4-DHPEA correlated ($r=0.97$) with the oxidative stability of VOO [81]. Both hydrolysis and nonenzymatic oxidation of phenolics can take place during storage, leading to a decrease in VOO secoiridoid content and an increase in simple phenolics (such as 3,4-DHPEA and 4-HPEA), elenolic acid, oxidised forms of elenolic acid and oxidised forms of secoiridoids, whereas lignans, 1-acetoxypinoresinol and pinoresinol were relatively stable [82-84]. Brenes *et al.* observed that hydrolysis of secoiridoid aglycons was the main change to the phenolic constituents in the VOOs of *Arbequina*, *Hojiblanca*, and *Picual* varieties during storage in the absence of light, resulting in an increase the contents of 3,4-DHPEA and 4-HPEA in these oils [83]. Okogeri and Tasioula-Margari studied the trends in the concentrations of phenolics during storage of VOO, and found that the concentrations of total phenolics decreased by 57–63% after 6 months under diffused light compared to 39–45% in the dark [85]. A decrease in the VOO phenolic concentration by as much as 81% after storage for 9 months has been reported [86].

A positive correlation has been found between the storage age of VOO and the ratio of 4-HPEA to total phenolics [87], and the ratio of 4-HPEA to total phenolics has been proposed as an indicator for distinguishing freshly made oils from oils stored for substantial periods (<4% for 6 month storage, >42% for longer than 18 month storage). 3,4-DHPEA-EDA degradation products are also suggested as molecular markers of the early auto-oxidation state of VOO, which could be an early evaluation index of VOO shelf life [88].

Conclusion

Consumption of olives and olive oil has been recognized as an important way to increase the dietary intake of phenolic compounds and derive the health benefits associated with these bioactive compounds. Olive cultivar, irrigation and the degree of ripening have major qualitative and quantitative effects on the phenolic constituents in olive fruit. Phenolic compounds are changed both qualitatively and quantitatively during the transfer from olive fruit to olive oil, resulting in quite different phenolic composition of VOO compared to that of olive fruit. Since only about 2% of phenolics in olive fruit are transferred to VOO and the rest ends up in the pomace or wastewater depending on the extraction system used, the phenolics present in pomace and wastewater could serve as potential sources of valuable phenolic compounds for edible use.

References

1. Lee OH, Lee BY. Antioxidant and antimicrobial activities of individual and combined phenolics in *Olea europaea* leaf extract. *Bioresour Technol.* 2010; 101: 3751-3754.
2. Lockyer S, Yaqoob P, Spencer JPE, Rowland I. Olive leaf phenolics and cardiovascular risk reduction: Physiological effects and mechanisms of action. *Nutrition and Aging.* 2012; 1: 125-140.
3. Casaburi I, Puoci F, Chimento A, Sirianni R, Ruggiero C, Avena P, et al. Potential of olive oil phenols as chemopreventive and therapeutic agents against cancer: a review of in vitro studies. *Mol Nutr Food Res.* 2013; 57: 71-83.
4. Huang CL, Sumpio BE. Olive oil, the mediterranean diet, and cardiovascular health. *J Am Coll Surg.* 2008; 207: 407-416.
5. Pontoniere P, Martiradonna D. Inflammation and olive phenolics A perspective review of supporting literature. *Agro Food Industry Hi-Tech.* 2012; 23:69-71.
6. Tuck KL, Hayball PJ. Major phenolic compounds in olive oil: metabolism and health effects. *J Nutr Biochem.* 2002; 13: 636-644.
7. Alonso A, Ruiz-Gutierrez V, Martinez-Gonzalez MA. Monounsaturated fatty acids, olive oil and blood pressure: epidemiological, clinical and experimental evidence. *Public Health Nutr.* 2006; 9: 251-257.
8. Covas MI. Bioactive effects of olive oil phenolic compounds in humans: reduction of heart disease factors and oxidative damage. *Inflammopharmacology.* 2008; 16: 216-218.
9. Konstantinidou V, Covas MI, Sola R, Fito M. Up-to date knowledge on the in vivo transcriptomic effect of the Mediterranean diet in humans. *Mol Nutr Food Res.* 2013; 57: 772-783.
10. Oi-Kano Y, Kawada T, Watanabe T, Koyama F, Watanabe K, Senbongi R, et al. Oleuropein supplementation increases urinary noradrenaline and testicular testosterone levels and decreases plasma corticosterone level in rats fed high-protein diet. *J Nutr Biochem.* 2013; 24: 887-893.
11. Bulotta S, Celano M, Lepore SM, Montalcini T, Pujia A, Russo D. Beneficial effects of the olive oil phenolic components oleuropein and hydroxytyrosol: focus on protection against cardiovascular and metabolic diseases. *J Transl Med.* 2014; 12: 219.
12. Impellizzeri D, Esposito E, Mazzon E, Paterniti I, Di Paola R, Bramanti P, et al. The effects of oleuropein aglycone, an olive oil compound, in a mouse model of carrageenan-induced pleurisy. *Clin Nutr.* 2011; 30: 533-540.
13. Kiritsakis AK. Flavor components of olive oil - A review. *J Amer Oil Chem Soc.* 1998; 75: 673-681.
14. Servili M, Selvaggini R, Esposito S, Taticchi A, Montedoro G, Morozzi G. Health and sensory properties of virgin olive oil hydrophilic phenols: agronomic and technological aspects of production that affect their occurrence in the oil. *J Chromatogr A.* 2004; 1054: 113-127.
15. Catalano P, Caponio F. Machines for olive paste preparation producing quality virgin olive oil. *Fett/Lipid.* 1996; 98: 408-412.
16. Morelló JR, Romero MP, Motilva MJ. Effect of the maturation process of the olive fruit on the phenolic fraction of drupes and oils from Arbequina, Farga, and Morrut cultivars. *J Agric Food Chem.* 2004; 52: 6002-6009.
17. Clodoveo ML. Malaxation: Influence on virgin olive oil quality. Past, present and future - An overview. *Trends Food Sci Technol.* 2012; 25: 13-23.
18. Rodis PS, Karathanos VT, Mantzavinou A. Partitioning of olive oil antioxidants between oil and water phases. *J Agric Food Chem.* 2002; 50: 596-601.
19. Fregapane G, Salvador MD. Production of superior quality extra virgin olive oil modulating the content and profile of its minor components. *Food Res Int.* 2013; 54: 1907-1914.
20. Boskou D. Olive oil: chemistry and technology. USA: AOCS Press. 1996.
21. Cardoso SM, Guyot S, Marnet N, Lopes-da-Silva JA, Silva AMS, Renard CMGC, et al. Identification of oleuropein oligomers in olive pulp and pomace. *J Sci Food Agr.* 2006; 86: 1495-1502.
22. Romani A, Mulinacci N, Pinelli P, Vincieri FF, Cimato A. Polyphenolic content in five tuscany cultivars of *Olea europaea* L. *J Agric Food Chem.* 1999; 47: 964-967.
23. Ryan D, Prenzler PD, Lavee S, Antolovich M, Robards K. Quantitative changes in phenolic content during physiological development of the olive (*Olea europaea*) cultivar Hardy's Mammoth. *J Agric Food Chem.* 2003; 51: 2532-2538.
24. Servili M, Baldioli M, Selvaggini R, Macchioni A, Montedoro G. Phenolic compounds of olive fruit: one- and two-dimensional nuclear magnetic resonance characterization of Nüzhenide and its distribution in the constitutive parts of fruit. *J Agric Food Chem.* 1999; 47: 12-18.
25. El Riachy M, Priego-Capote F, Rallo L, Luque-de, Castro MD, León L. Phenolic profile of virgin olive oil from advanced breeding selections. *Span J Agric Res.* 2012; 10: 443-453.
26. Servili M, Montedoro G. Contribution of phenolic compounds to virgin olive oil quality. *Eur J Lipid Sci Technol.* 2002; 104: 602-613.
27. Owen RW, Mier W, Giacosa A, Hull WE, Spiegelhalter B, Bartsch H. Phenolic compounds and squalene in olive oils: the concentration and antioxidant potential of total phenols, simple phenols, secoiridoids, lignans and squalene. *Food Chem Toxicol.* 2000; 38: 647-659.
28. Garcia A, Brenes M, Martínez F, Alba J, García P, Garrido A. High-performance liquid chromatography evaluation of phenols in virgin olive oil during extraction at laboratory and industrial scale. *J Amer Oil Chem Soc.* 2001; 78: 625-629.
29. Garcia B, Coelho J, Costa M, Pinto J, Paiva-Martins F. A simple method for the determination of bioactive antioxidants in virgin olive oils. *J Sci Food Agric.* 2013; 93: 1727-1732.
30. Tovar MJ, Motilva MJ, Romero MP. Changes in the phenolic composition of virgin olive oil from young trees (*Olea europaea* L. cv. Arbequina) grown under linear irrigation strategies. *J Agric Food Chem.* 2001; 49: 5502-5508.
31. de la Torre-Carbot K, Jauregui O, Gimeno E, Castellote AI, Lamuela-Raventós RM, López-Sabater MC. Characterization and quantification of phenolic compounds in olive oils by solid-phase extraction, HPLC-DAD, and HPLC-MS/MS. *J Agric Food Chem.* 2005; 53: 4331-4340.
32. Brenes M, Garcia A, Garcia P, Rios JJ, Garrido A. Phenolic compounds in Spanish olive oils. *J Agric Food Chem.* 1999; 47: 3535-3540.
33. Carrasco Pancorbo A, Cruces-Blanco C, Segura Carretero A, Fernández Gutierrez A. Sensitive determination of phenolic acids in extra-virgin olive oil by capillary zone electrophoresis. *J Agric Food Chem.* 2004; 52: 6687-6693.
34. Kesen S, Kelebek H, Selli S. Characterization of the volatile phenolic and antioxidant properties of monovarietal olive oil obtained from cv. Halhali. *J Amer Oil Chem Soc.* 2013; 90: 1685-1696.
35. Bianco A, Coccioli F, Guiso M, Marra C. The occurrence in olive oil of a new class of phenolic compounds: hydroxy-isochromans. *Food Chem.* 2002; 77: 405-411.
36. Mulinacci N, Romani A, Galardi C, Pinelli P, Giaccherini C, Vincieri FF. Polyphenolic content in olive oil waste waters and related olive samples. *J Agric Food Chem.* 2001; 49: 3509-3514.
37. Daqdelen A, Tumen G, Ozcan MM, Dundar E. Phenolics profiles of olive fruits (*Olea europaea* L.) and oils from Ayvalik, Domat and Gemlik varieties at different ripening stages. *Food Chem.* 2013; 136: 41-45.
38. Gomez-Rico A, Fregapane G, Salvador MD. Effect of cultivar and ripening on minor components in Spanish olive fruits and their corresponding virgin olive oils. *Food Res Int.* 2008; 41: 433-440.
39. Morello JR, Romero MP, Ramo T, Motilva MJ. Evaluation of L-phenylalanine ammonia-lyase activity and phenolic profile in olive drupe (*Olea europaea* L.) from fruit setting period to harvesting time. *Plant Sci.* 2005; 168: 65-72.
40. Therios IN. Olive: Crop production science in horticulture 18. UK: CABI Press, 2009.
41. Rigane G, Ben Salem R, Sayadi S, Bouaziz M. Phenolic composition, isolation, and structure of a new deoxyloganic acid derivative from Dhokar and Gemri-Dhokar olive cultivars. *J Food Sci.* 2011; 76: 965-973.

42. Jemai H, Bouaziz M, Sayadi S. Phenolic composition, sugar contents and antioxidant activity of Tunisian sweet olive cultivar with regard to fruit ripening. *J Agric Food Chem*. 2009; 57: 2961-2968.
43. Markakis EA, Tjamos SE, Antoniou PP, Roussos PA, Paplomatas EJ, Tjamos EC. Phenolic responses of resistant and susceptible olive cultivars induced by defoliating and nondefoliating verticillium dahliae pathotypes. *Plant Dis*. 2010; 94: 1156-1162.
44. Vinha AF, Ferreres F, Silva BM, Valentao P, Gonçalves A, Pereira JA, et al. Phenolic profiles of Portuguese olive fruits (*Olea europaea* L.): Influences of cultivar and geographical origin. *Food Chem*. 2005; 89: 561-568.
45. Esti M, Cinquanta L, La Notte E. Phenolic Compounds in Different Olive Varieties. *J Agric Food Chem*. 1998; 46: 32-35.
46. He ZY, Xia WS, Liu QH, Chen J. Identification of a new phenolic compound from Chinese olive (*Canarium album* L.) fruit. *Eur Food Res Technol*. 2009; 228: 339-343.
47. García JM, Sella S, Perez-Camino MC. Influence of fruit ripening on olive oil quality. *J Agr Food Chem*. 1996; 44: 3516-3520.
48. Bianco A, Scalzo RL, Scarpati ML. Isolation of cornoside from *Olea europaea* and its transformation into halleridone. *Phytochemistry*. 1993; 32: 455-457.
49. Ghanbari R, Anwar F, Alkharfy KM, Gilani AH, Saari N. Valuable Nutrients and Functional Bioactives in Different Parts of Olive (*Olea europaea* L.)-A Review. *Int J Mol Sci*. 2012; 13: 3291-3340.
50. Brenes M, Rejano L, Garcia P, Sanchez AH, Garrido A. Biochemical changes in phenolic compounds during Spanish-style green olive Processing. *J Agr Food Chem*. 1995; 43: 2702-2706.
51. Amiot MJ, Fleuriot A, Macheix JJ. Accumulation of oleuropein derivatives during olive maturation. *Phytochemistry*. 1989; 28: 67-69.
52. Damak N, Bouaziz M, Ayadi M, Sayadi S, Damak M. Effect of the maturation process on the phenolic fractions, fatty acids, and antioxidant activity of the Chétoui olive fruit cultivar. *J Agric Food Chem*. 2008; 56: 1560-1566.
53. Yorulmaz A, Erinc H, Tekin A. Changes in olive and olive oil characteristics during maturation. *J Amer Oil Chem Soc*. 2013; 90: 647-658.
54. Menz G, Vriesekoop F. Physical and chemical changes during the maturation of Gordal Sevillana olives (*Olea europaea* L. cv. Gordal Sevillana). *J Agric Food Chem*. 2010; 58: 4934-4938.
55. Bouaziz M, Chamkha M, Sayadi S. Comparative study on phenolic content and antioxidant activity during maturation of the olive cultivar Chemlali from Tunisia. *J Agric Food Chem*. 2004; 52: 5476-5481.
56. Machado M, Felizardo C, Fernandes-Silva AA, Nunes FM, Barros A. Polyphenolic compounds, antioxidant activity and L-phenylalanine ammonia-lyase activity during ripening of olive cv. "Cobrançosa" under different irrigation regimes. *Food Res Int*. 2013; 51: 412-421.
57. Caruso G, Gucci R, Urbani S, Esposito S, Taticchi A, Di Maiol, et al. Effect of different irrigation volumes during fruit development on quality of virgin olive oil of cv. Frantoio. *Agr Water Manage*. 2014; 134: 94-103.
58. Patumi M, d'Andria R, Fontanazza G, Morelli G, Giorio P, Sorrentino G. Yield and oil quality of intensively trained trees of three cultivars of olive (*Olea europaea* L.) under different irrigation regimes. *J Hortic Sci Biotechnol*. 1999; 74: 729-737.
59. Tovar MJ, Romero MP, Girona J, Motilva MJ. L-phenylalanine ammonia-lyase activity and concentration of phenolics in developing olive (*Olea europaea* L cv Arbequina) fruit grown under different irrigation regimes. *J Sci Food Agr*. 2002; 82: 892-898.
60. Martinelli F, Basile B, Morelli G, d'Andria R, Tonutti P. Effects of irrigation on fruit ripening behavior and metabolic changes in olive. *Sci Hort*. 2012; 144: 201-207.
61. Ryan D, Robards K. Phenolic compounds in olives. *Analyst*. 1998; 123: 31-44.
62. Ryan D, Antolovich M, Prenzler P, Robards K, Lavee S. Biotransformations of phenolic compounds in *Olea europaea* L. *Sci Hort*. 2002; 92: 147-176.
63. Gómez-Rico A, Inarejos-García AM, Salvador MD, Fregapane G. Effect of malaxation conditions on phenol and volatile profiles in olive paste and the corresponding virgin olive oils (*Olea europaea* L. Cv. Cornicabra). *J Agric Food Chem*. 2009; 57: 3587-3595.
64. Vierhuis E, Servili M, Baldioli M, Schols HA, Voragen AG, Montedoro GF. Effect of enzyme treatment during mechanical extraction of olive oil on phenolic compounds and polysaccharides. *J Agric Food Chem*. 2001; 49: 1218-1223.
65. Montedoro GF, Baldioli M, Selvaggini R, Begliuomini AL, Taticchi A, Servili M. Relationships between phenolic composition of olive fruit and olive oil: The importance of the endogenous enzymes. *Acta Hort*. 2002; 586: 551-556.
66. Boselli E, Di Lecce G, Strabbioli R, Perialisi G, Frega NG. Are virgin olive oils obtained below 27°C better than those produced at higher temperatures? *LWT-Food Sci Technol*. 2009; 42: 748-757.
67. Kalua CM, Bedgood DR, Bishop AG, Prenzler PD. Changes in volatile and phenolic compounds with malaxation time and temperature during virgin olive oil production. *J Agric Food Chem*. 2006; 54: 7641-7651.
68. Stefanoudaki E, Koutsaftakis A, Harwood JL. Influence of malaxation conditions on characteristic qualities of olive oil. *Food Chem*. 2011; 127: 1481-1486.
69. Parenti A, Spugnoli P, Masella P, Calamai L. The effect of malaxation temperature on the virgin olive oil phenolic profile under laboratory-scale conditions. *Eur J Lipid Sci Technol*. 2008; 110: 735-741.
70. Servili M, Baldioli M, Mariotti F, Montedoro GF. Phenolic composition of olive fruit and virgin olive oil: distribution in the constitutive parts of fruit and evolution during oil mechanical extraction process. *Acta Hort*. 1999; 474: 609-619.
71. Taticchi A, Esposito S, Veneziani G, Urbani S, Selvaggini R, Servili M. The influence of the malaxation temperature on the activity of polyphenoloxidase and peroxidase and on the phenolic composition of virgin olive oil. *Food Chem*. 2013; 136: 975-983.
72. Ranalli A, Pollastri L, Contento S, Iannucci E, Lucera L. Effect of olive paste kneading process time on the overall quality of virgin olive oil. *Eur J Lipid Sci Technol*. 2003; 105: 57-67.
73. Youssef O, Mokhtar G, Abdely C, Mohamed SN, Mokhtar Z, Guido F. Changes in volatile compounds and oil quality with malaxation time of Tunisian cultivars of *Olea europaea*. *Int J Food Sci Technol*. 2013; 48: 74-81.
74. Artajo LS, Romero MP, Motilva MJ. Transfer of phenolic compounds during olive oil extraction in relation to ripening stage of the fruit. *J Sci Food Agr*. 2006; 86: 518-527.
75. Jimenez B, Sanchez-Ortiz A, Rivas A. Influence of the malaxation time and olive ripening stage on oil quality and phenolic compounds of virgin olive oils. *Int J of Food Sci Technol*. 2014.
76. Frankel E, Bakhouché A, Lozano-Sanchez J, Segura-Carretero A, Fernandez-Gutierrez A. Literature review on production process to obtain extra virgin olive oil enriched in bioactive compounds. Potential use of byproducts as alternative sources of polyphenols. *J Agr Food Chem*. 2013; 61: 5179-5188.
77. Salvador MD, Aranda F, Fregapane G. Chemical composition of commercial Cornicabra virgin olive oil from 1995/96 and 1996/97 crops. *J Amer Oil Chem Soc*. 1998; 75: 1305-1311.
78. Salvador MD, Aranda F, Gomez-Alonso S, Fregapane G. Influence of extraction system, production year and area on Cornicabra virgin olive oil: a study of five crop seasons. *Food Chem*. 2003; 80: 359-366.
79. Klen TJ, Vodopivec BM. The fate of olive fruit phenols during commercial olive oil processing: Traditional press versus continuous two- and three-phase centrifuge. *Lwt-Food Sci Technol*. 2012; 49: 267-274.
80. Vossen, PM. Organic olive production manual. USA: UCANR Publications. 2007.
81. Baldioli M, Servili M, Perretti G, Montedoro GF. Antioxidant activity of tocopherols and phenolic compounds of virgin olive oil. *J Amer Oil Chem*. 1996; 73: 1589-1593.

82. Baiano A, Gambacorta G, Terracone C, Previtali MA, Lamacchia C, La Notte E. Changes in phenolic content and antioxidant activity of Italian extra-virgin olive oils during storage. *J Food Sci.* 2009; 74: 177-183.
83. Brenes M, García A, García P, Garrido A. Acid hydrolysis of secoiridoid aglycons during storage of virgin olive oil. *J Agric Food Chem.* 2001; 49: 5609-5614.
84. Morello JR, Motilva MJ, Tovar MJ, Romero MP. Changes in commercial virgin olive oil (cv Arbequina) during storage, with special emphasis on the phenolic fraction. *Food Chem.* 2004; 85: 357-364.
85. Okogeri O, Tasioula-Margari M. Changes occurring in phenolic compounds and alpha-tocopherol of virgin olive oil during storage. *J Agric Food Chem.* 2002; 50: 1077-1080.
86. Baiano A, Terracone C, Gambacorta G, La Notte E. Changes in quality indices, phenolic content and antioxidant activity of flavored olive oils during storage. *J Amer Oil Chem Soc.* 2009; 86: 1083-1092.
87. Cinquanta L, Esti M, La Notte E. Evolution of phenolic compounds in virgin olive oil during storage. *J Amer Oil Chem Soc.* 1997; 74: 1259-1264.
88. Di Maio I, Esposto S, Taticchi A, Selvaggini R, Veneziani G, Urbani S, et al. Characterization of 3,4-DHPEA-EDA oxidation products in virgin olive oil by high performance liquid chromatography coupled with mass spectrometry. *Food Chem.* 2013; 138: 1381-1391.