

Special Article - Antioxidants in Foods

Saffron, A Functional Spice

Anastasia Kyriakoudi¹, Stella A Ordoudi¹, Marta Roldán-Medina² and Maria Z Tsimidou^{1*}¹School of Chemistry, Aristotle University of Thessaloniki (AUTh), Greece²EU & International Projects Promoter, Universidad de Castilla-La Mancha (UCLM), Spain*Corresponding author: Tsimidou MZ, School of Chemistry, Aristotle University of Thessaloniki, 54124, Thessaloniki, Greece, Tel: +302310997796; Fax: +302310997847; E-mail: tsimidou@chem.auth.gr

Received: February 27, 2015; Accepted: April 20, 2015; Published: April 27, 2015

Abstract

Saffron, the dehydrated red stigmas of the flower of the plant *Crocus sativus* L., comprises the most expensive spice in the world. It is mainly used as a spice highly valued for its coloring power, bitter taste and unique aroma attributed primarily to crocins, picrocrocin and safranal, respectively. Apart from its known applications in the food industry, a number of pharmacological actions have been assigned to saffron and its constituents. Expansion of the use of natural products in the prevention of chronic diseases or cancer as well as the interest in functional foods is expected to attract the attention of consumers to include saffron in everyday diet. Only authentic high quality saffron exerts beneficial health effects.

Keywords: Saffron; *Crocus sativus* L.; Apocarotenoids; Authenticity; Antioxidant activity

Abbreviations

AP-1: Activator Protein 1; CNS: Central Nervous System; DNA: Deoxyribonucleic acid; FRAP: Ferric Reducing Antioxidant Power; GSH: Glutathione Peroxidase; i.p.: Intraperitoneally; LDL: Low Density Lipoprotein; MDA: Malondialdehyde; NFκB: Nuclear Factor Kappa B; NO: Nitric Oxide; NSAIDs: Non-Steroidal Antiinflammatory Drugs; p.o.: Per Os (by mouth); ROS: Reactive Oxygen Species; SOD: Superoxide Dismutase


Introduction

Saffron is comprised of the dehydrated red stigmas of the flower of the plant *Crocus sativus* L. (Figure 1A), a sterile triploid that belongs to the subfamily *Crocoideae* of the Iridaceae family [1]. Saffron is the most expensive spice in the world, highly valued in different cuisines for its distinctive color, exceptional taste and unique aroma. The compounds that elicit yellow color hues are certain highly water-soluble apocarotenoids, the sugar esters of crocetin (8,8'-diapocarotene-8,8'-dioic acid), selectively named as crocins. The *trans*-crocetin (di-β-d-gentiobiosyl) ester (*trans*-4-GG), commonly referred as crocin, is the most abundant member. The gustatory traits of the spice are given by the monoterpene glucoside, picrocrocin [4-(β-D-glucopyranosyloxy)-2,6,6-trimethyl-1-cyclohexane-1-carboxaldehyde] (bitterness) and certain volatiles, e.g. safranal (2,6,6-trimethyl-1,3-cyclohexadiene-1-carboxaldehyde) (aroma) [2]. Recently, light was shed on the biosynthetic pathway of these apocarotenoids through transcriptome sequencing [3]. The precursor of saffron apocarotenoids is considered to be zeaxanthin that through successive cleavages give rise to the formation of crocetin and 3-OH-β-cyclocitral, which are then glucosylated to yield crocins and picrocrocin, respectively. These apocarotenoids accumulate throughout the development of stigmas, reach a maximum concentration at a fully developed stage and decrease thereafter, a fact that justifies traditional collection practices [4]. Crocins and picrocrocin account for nearly 50 % (w/w) in dried stigmas. The moisture content of the latter is much lower and ranges between 9-12 %. Safranal is mainly formed during processing and storage and represents ca. 70% of the essential oil of the spice [5]. The chemical

structures of the mentioned compounds are given in Figure 1B.

Culinary herbs and spices have been traditionally used not only as flavor enhancers and food preservatives but also as folk medicines. Concerning saffron, its use as a spice or textile dye dates back to early antiquity era whereas in traditional Indian, Chinese and European medieval medicines its extracts and tinctures are reported as antispasmodic, eupeptic, gingival sedative, carminative, diaphoretic, expectorant, stimulant, stomachic, aphrodisiac and emmenagogue [5]. Today, there is a growing body of evidence from epidemiological studies that saffron apocarotenoids may present multiple biological actions including anticancer, anti-inflammatory, antimicrobial and antioxidant [6]. A search in the Scopus database using as keywords "saffron" and "biological" (11-02-2015) shows that the relevant scientific articles published the last five years ($n=53$) are almost two fold more than those published between 2005 and 2010. The results of these studies provide strong evidence for the effectiveness of either individual apocarotenoids (crocetin, crocins or safranal) or saffron extracts against various diseases. The most important health benefits concern activity against several types of cancer [7,8], gastric disorders [9], cardiovascular diseases [10], Alzheimer disease [11], and mild to moderate depression [12]. Administration of saffron has been also found to ameliorate the side effects of conventional treatments e.g. sexual dysfunction by antidepressant drugs [13].

Saffron is likely safe for most individuals when used at levels commonly found in diet [14]. In all the preclinical and clinical studies so far, saffron has been administered orally or intraperitoneally. The common effective doses in clinical trials (30–50 mg/day) were much lower than the LD50 of saffron. The latter depends on the form of intervention and can range from around 1.5 g/kg (after injection in mice) to >5 g/kg (after human consumption) indicating a high safety margin for the exploitation of saffron in pharmaceutical preparations [15]. Isolated constituents are also considered safe; a dose of 20 mg *trans*-4-GG/day over one month (in tablets) was found non-toxic for human consumption and safranal was non-toxic for both mice and rats after oral administration [16,17]. It should be stressed as a general remark that although intravenously injected extracts or compounds are considered fully bioavailable to exert any function *in*



vivo, this is not the case for orally administered substances. As far as saffron bioactive ingredients are concerned, the data regarding their absorption profile are limited [18,19]. Independent researchers have shown that crocins, although highly bioaccessible [20] are most likely hydrolyzed in the gastrointestinal tract to crocetin, which is then absorbed as such or as glucuronide conjugates [21]. Crocetin is reported to slowly penetrate the blood brain barrier to reach the central nervous system [22].

Physiological Benefits through Antioxidant Action

The physiological disorders that can be alleviated after the treatment with saffron, crocins, safranal or crocetin are usually the result of various, often complex and unclear mechanisms. In most circumstances elevated levels of oxidative stress and reduced antioxidant defenses are evidenced. In the following paragraphs, certain pharmacological effects that have been demonstrated so far *in vivo* and attributed to the antioxidant activity of saffron and its constituents are presented.

Effects on Central Nervous System (CNS)

The pathogenesis of acute Central Nervous System (CNS) disorders (e.g. dementia, ischemia) is associated with oxidative stress effects. After brain injury, Reactive Oxygen Species (ROS) are overproduced resulting in imbalance of prooxidant/antioxidant species and damage of tissues via several different cellular molecular pathways. In principle, ROS are considered to play a major role in the formation of Nitric Oxide (NO) and Malondialdehyde (MDA) and to suppress the activity of antioxidant enzymes such as Superoxide Dismutase (SOD) and Glutathione Peroxidase (GSH). The exceeding oxidant species may, thus, impair key cellular components (lipids,

proteins, and nucleic acids) leading to subsequent cell death (necrosis or apoptosis) [23].

Since 2005, when safranal was found to protect hippocampal tissue of ischemic rats from oxidative damage [23], a number of preclinical experiments in animal models were carried out using saffron extract [24], isolated crocins [25-27], picrocrocin [25] or crocetin [28] to study their effect on markers of cerebral ischemic-induced injuries. So far literature indicates that administration of saffron extracts at doses > 100 mg/kg b.w. (i.p. or p.o.), *trans*-4-GG at > 25 mg/kg b.w. (i.p.), safranal at > 700 mg/kg b.w. (i.p.) or crocetin at > 8 mg/kg b.w. (i.p.) is expected to be beneficial for the abovementioned pathological conditions.

Saffron, crocins and safranal have also shown cognitive-enhancing effects in animal studies [29]. Few clinical trials have been carried out so far and provide evidence that daily and chronic consumption of saffron extract capsules (standardized to 1.7 mg *trans*-4-GG/capsule) by patients suffering from Alzheimer disease is as much helpful as conventional treatments with less adverse effects [29]. The underlying mechanism is still not known; *in vitro* studies have shown that crocins and to a lesser extent, safranal, are likely to inhibit the oxidation-induced formation of toxic amyloid fibrils which signal the onset of various degenerative diseases such as the Alzheimer one e.g. [30,31].

The anti-depressant effect of saffron was evidenced by several clinical trials so far e.g. [32,33]. This type of action against inflammatory and immune responses, disruption of neurotransmitter balance and neurodegeneration may involve antioxidant, anti-inflammatory, serotonergic, hypothalamus-pituitary-adrenal (HPA) axis-modulating and neuroprotective mechanisms [12].

Effects on Cardiovascular System

Crocetin and its *trans*-4-GG ester have shown a potential in the treatment of atherosclerosis, hyperlipidemia and other cardiovascular disorders. In particular, oral administration of these constituents to quails (25, 50, 100 mg/kg/day) for a period of 9 weeks was found to attenuate atherosclerosis mainly by decreasing the levels of oxidized LDL [34,35]. Zheng et al. [10] reached the same conclusion after feeding hyperlipidemic rabbits with crocetin for 8 weeks. The latter research group reported also that the hypolipidemic effect of *trans*-4-GG crocetin can be attributed to the inhibition of pancreatic lipase activity, which mediates the hydrolysis of fat [36]. In addition, orally administered *trans*-4-GG at 20 mg/kg was found to improve cardiac functions in rats fed with a cardiotoxic agent (isoproterenol) by maintaining the redox status of the cell, restoring endogenous antioxidants, controlling lipid peroxide formation and preserving activities of CK-MB, LDH enzymes [37].

Effects on Diabetes

It is generally accepted that diabetes and its complication, that is, peripheral neuropathy and retinopathy are associated to oxidative stress. Although dietary supplementation of saffron may induce a short-term improvement in age-related macular degeneration, it is not clear yet whether the improvement of retinal function is due to antioxidant effects [38]. It has been suggested though that supplementation of saffron extract or *trans*-4-GG may be beneficial against glucose-induced neurotoxicity in PC12 cells by inhibiting the formation of ROS [39].

Effects against Non-Steroidal Antiinflammatory Drugs (Nsaids)-Induced Gastropathy

Recently, saffron, *trans*-4-GG and safranal were found to prevent the formation of gastric lesions and lipid peroxidation products as well as the depletion of glutathione induced by indomethacin in non-diabetic and diabetic rats. The efficiency was comparable to that of a conventional medicine, omeprazole and attributed to the antioxidant action of the plant constituents [9].

Mechanism of Antioxidant Action

Saffron and its constituents, *trans*-4-GG, crocetin and safranal, are potent antioxidants. This is proved in several *in vivo* studies with animals in which, after dietary supplementation of the medicinal plant, the levels of antioxidant enzymes such as superoxide dismutase and catalase as well as reduced glutathione increase e.g. [40,41]. On the other hand, the content in lipid peroxidation products such as malondialdehyde, 8-iso-prostaglandin F(2 beta), hippocampal nitrite levels, oxidized LDL, oxidized DNA decreases [41,42]. In several cases, the saffron, *trans*-4-GG, safranal or crocetin-treated animal tissues (e.g. brain, liver, hippocampus) showed a positive response toward the Ferric Reducing Antioxidant Power (FRAP) assay indicating elevated levels of reducing substances [10,23,40,43].

Despite the numerous studies that demonstrate the aforementioned effects, the exact mechanism of action is still unknown. In fact, saffron extracts or free crocetin are not expected to act like the phenolic-type antioxidants that is, by direct scavenging

of free radical species in cell-free models such as those employed in common antiradical activity assays (e.g. DPPH, TEAC) [20,44]. The strong activity of saffron constituents in comparison to that of known phenolics is revealed only intracellularly; crocins seem to have an intracellular antioxidant effect comparable to that of α -tocopherol [45] and crocetin is more powerful than curcumin and rosmarinic acid in the protection of insulin-stimulated human monocytes [44]. Thus, the most probable pathway is the interaction of crocins and crocetin (through the conjugated double bond system) with antioxidant enzymes or with signal transduction of free radicals in the cells. For example, they could inhibit generation of H_2O_2 , which acts as a signal transduction messenger to activate transcription factors NF κ B (Nuclear Factor Kappa B) and AP-1 (Activator Protein 1) for gene expression of various inflammatory cytokines and proteases that cause cell damage [9]. Safranal, as a lipophilic monoterpene aldehyde could also act as inhibitor of ROS through a pathway that remains to be elucidated.

The pharmacologically active compounds present in saffron seem to deliver their potency synergistically which highlights the benefits of dietary supplementation of the spice rather than isolated constituents [40]. Extracts rich in crocins or isolated crocetin are prone to autoxidation and photoxidation due to extensive conjugation. Their shelf life can be prolonged via encapsulation e.g. [46,47] as it is also reported for other bioactive ingredients.

Last but not least, it is necessary to emphasize that all the biological functions that are ascribed to saffron can be claimed only if the plant material used is authentic and of high quality. Peculiar adverse effects that have been reported e.g. mortality after ingestion of 4 g/day is rather associated with the use of fake saffron. Schmidt et al. [15] suggested that many such cases that are reported from Germany, refer to "meadow saffron" (*Colchicum autumnale*), which is a poisonous plant and relatively abundant there. Other mislabeled products such as "American saffron" that corresponds to *Carthamus tinctorius* L. (safflower) petals and "African saffron", which is actually the flowers of *Lyperia crocea* should be avoided. Since saffron is particularly susceptible to fraudulent practices [48] it may contain other parts of the flower of *Crocus sativus* L., or the flowers of other *crocus* species or other plants (e.g. *Calendula officinalis* L., marigold) with or without coloring properties. The ground rhizomes of *Curcuma longa* (turmeric) are often mislabeled as saffron powder [5]. Saffron of low quality or non-colored exogenous matter used as substitute can be also adulterated with synthetic dyes [49]. The development of modern high throughput analytical methods for detecting adulteration in saffron is a primary objective of the research network established through the COST Action FA 1101 with the acronym Saffron-OMICS (www.saffronomics.org, www.cost.eu/COST_Actions/fa/Actions/FA1101) [50]. The importance of this program lies on the involvement of scientists from countries in Europe and elsewhere (e.g. Egypt, New Zealand) with great interest in acquiring expertise in the cultivation and production of high quality saffron. The efforts of this network is to offer to the scientific community and stakeholders advanced tools for combating saffron fraud with lower quality plant materials (*Gardenia jasminoides* Ellis, *Buddleja officinalis* Maxim, etc) and other types of adulterants [51-53]. Introduction of the use of authentic saffron in everyday diet seems to exert beneficial health effects so that the spice can be considered as a functional food ingredient.

Acknowledgement

M.Z.T (Chair), M.R.M (Vice Chair, Grant Holder), S.A.O. (WG3 co-leader) and A.K. (WG2 member, ESR) thank the European Science Foundation (ESF) through the COST Action FA1101 (Saffron –OMICS) for giving them the opportunity to spread worldwide information on authentic saffron and its inherent culinary and health properties.

References

- Fernández JA, Santana O, Guardiola JL, Molina RV, Heslop-Harrison P, Borbely G, et al. The World Saffron and Crocus collection: strategies for establishment, management, characterization and utilization. *Genet. Resour. Crop. Evol.* 2011; 58: 125-137.
- Carmona M, Zalacain A, Alonso GL. The chemical composition of saffron: color, taste and aroma. 1st edn. Albacete, Spain: Editorial Bomarzo S.L. 2006.
- Frusciante S, Diretto G, Bruno M, Ferrante P, Pietrella M, Prado-Cabrero A, et al. Novel carotenoid cleavage dioxygenase catalyzes the first dedicated step in saffron crocin biosynthesis. *Proc Natl Acad Sci USA.* 2014; 111: 12246-12251.
- Rubio-Moraga A, Trapero A, Ahrazem O, Gómez-Gómez L. Crocins transport in *Crocus sativus*: the long road from a senescent stigma to a newborn corn. *Phytochemistry.* 2010; 71: 1506-1513.
- Ordoudi S, Tsimidou M. Saffron quality: Effect of agricultural practices, processing and storage. Dris R, Jain SM, editors. In: *Production Practices and Quality Assessment of Food Crops, Volume 1.* Kluwer Academic Publishers. 2004; 209-260.
- Kaefer CM, Milner JA. The role of herbs and spices in cancer prevention. *J Nutr Biochem.* 2008; 19: 347-361.
- Gutheil WG, Reed G, Ray A, Anant S, Dhar A. Crocetin: An agent derived from saffron for prevention and therapy for cancer. *Curr. Pharm. Biotechnol.* 2012; 13: 173-179.
- Zhang Z, Wang CZ, Wen XD, Shoyama Y, Yuan CS. Role of saffron and its constituents on cancer chemoprevention. *Pharm. Biol.* 2013; 51: 920-924.
- Kianbakht S, Mozaffari K. Effects of saffron and its active constituents, crocin and safranal, on prevention of indomethacin induced gastric ulcers in diabetic and nondiabetic rats. *J. Med. Plants.* 2009; 8: 30-38.
- Zheng S, Qian Z, Sheng L, Wen N. Crocetin attenuates atherosclerosis in hyperlipidemic rabbits through inhibition of LDL oxidation. *J Cardiovasc Pharmacol.* 2006; 47: 70-76.
- Moshiri M, Vahabzadeh M, Hosseinzadeh H2. Clinical Applications of Saffron (*Crocus sativus*) and its Constituents: A Review. *Drug Res (Stuttg).* 2014;.
- Lopresti AL, Drummond PD. Saffron (*Crocus sativus*) for depression: a systematic review of clinical studies and examination of underlying antidepressant mechanisms of action. *Hum. Psychopharmacol.* 2014; 29: 517-527.
- Kashani L, Raisi F, Saroukhani S, Sohrabi H, Modabbernia A, Nasehi AA, et al. Saffron for treatment of fluoxetine-induced sexual dysfunction in women: randomized double-blind placebo-controlled study. *Hum. Psychopharmacol.* 2013; 28: 54-60.
- Ulbricht C, Conquer J, Costa D, Hollands W, Kannuzzi C, Isaac R, et al. An evidence-based systematic review of saffron (*Crocus sativus*) by the Natural Standard Research Collaboration. *J. Diet. Suppl.* 2011; 8: 58-114.
- Schmidt M, Betti G, Hensel A. Saffron in phytotherapy: pharmacology and clinical uses. *Wien Med Wochenschr.* 2007; 157: 315-319.
- Hosseinzadeh H, Sadeghi Shakib S, Khadem Sameni A, Taghiabadi E. Acute and subacute toxicities of safranal, a constituent of saffron, in mice and rats. *Clin. Biochem.* 2011; 44: S122.
- Modagheh MH, Shahabian M, Esmaili HA, Rajbai O, Hosseinzadeh H. Safety evaluation of saffron (*Crocus sativus*) tablets in healthy volunteers. *Phytomedicine.* 2008; 15: 1032-1037.
- Xi L, Qian Z, Du P, Fu J. Pharmacokinetic properties of crocin (crocetin digentiobiose ester) following oral administration in rats. *Phytomedicine.* 2007; 14: 633-636.
- Chryssanthi DG, Lamari FN, Georgakopoulos CD, Cordopatis P. A new validated SPE-HPLC method for monitoring crocetin in human plasma--application after saffron tea consumption. *J Pharm Biomed Anal.* 2011; 55: 563-568.
- Kyriakoudi A, Tsimidou MZ, O'Callaghan YC, Galvin K, O'Brien NM. Changes in total and individual crocetin esters upon in vitro gastrointestinal digestion of saffron aqueous extracts. *J Agric Food Chem.* 2013; 61: 5318-5327.
- Asai A, Nakano T, Takahashi M, Nagao A. Orally administered crocetin and crocins are absorbed into blood plasma as crocetin and its glucuronide conjugates in mice. *J Agric Food Chem.* 2005; 53: 7302-7306.
- Lautenschläger M, Sendker J, Hüwel S, Galla HJ, Brandt S, Düfer M. et al. Intestinal formation of trans-crocetin from saffron extract (*Crocus sativus* L.) and in vitro permeation through intestinal and blood brain barrier. *Phytomedicine* 2015; 22: 36-44.
- Hosseinzadeh H, Sadeghnia HR. Safranal, a constituent of *Crocus sativus* (saffron), attenuated cerebral ischemia induced oxidative damage in rat hippocampus. *J Pharm Pharm Sci.* 2005; 8: 394-399.
- Saleem S, Ahmad M, Ahmad AS, Yousuf S, Ansari MA, Khan MB, et al. Effect of Saffron (*Crocus sativus*) on neurobehavioral and neurochemical changes in cerebral ischemia in rats. *J Med Food.* 2006; 9: 246-253.
- Ochiai T, Shimeno H, Mishima K, Iwasaki K, Fujiwara M, Tanaka H, et al. Protective effects of carotenoids from saffron on neuronal injury in vitro and in vivo. *Biochim Biophys Acta.* 2007; 1770: 578-584.
- Hosseinzadeh H, Sadeghnia HR, Ghaeni FA, Motamedshariaty VS, Mohajeri SA. Effects of saffron (*Crocus sativus* L.) and its active constituent, crocin, on recognition and spatial memory after chronic cerebral hypoperfusion in rats. *Phytother Res.* 2012; 26: 381-386.
- Vakili A, Einali MR, Bandegi AR. Protective effect of crocin against cerebral ischemia in a dose-dependent manner in a rat model of ischemic stroke. *Journal of Stroke and Cerebrovascular Diseases: The Official Journal of National Stroke Association.* 2014; 23: 106-113.
- Tashakori-Sabzevar F, Hosseinzadeh H, Motamedshariaty VS, Movassaghi AR, Mohajeri SA. Crocetin attenuates spatial learning dysfunction and hippocampal injury in a model of vascular dementia. *Curr Neurovasc Res.* 2013; 10: 325-334.
- Pitsikas N. The Effect of *Crocus sativus* L. and its constituents on memory?: Basic studies and clinical applications. *Evidence-Based Complementary and Alternative Medicine* 2015; 1-7.
- Ebrahim-Habibi MB, Amininasab M, Ebrahim-Habibi A, Sabbaghian M, Nemat-Gorgani M. Fibrillation of alpha-lactalbumin: effect of crocin and safranal, two natural small molecules from *Crocus sativus*. *Biopolymers.* 2010; 93: 854-865.
- Papandreou MA, Kanakis CD, Polissiou MG, Efthimiopoulos S, Cordopatis P, Margariti M, et al. Inhibitory activity on amyloid- β aggregation and antioxidant properties of *Crocus sativus* stigmas extract and its crocin constituents. *J. Agric. Food Chem.* 2006; 54: 8762-8768.
- Akhondzadeh S, Fallah-Pour H, Afkham K, Jamshidi AH, Khalighi-Cigaroudi F. Comparison of *Crocus sativus* L. and imipramine in the treatment of mild to moderate depression: a pilot double-blind randomized trial [ISRCTN45683816]. *BMC Complement Altern Med.* 2004; 4: 12.
- Shahmansouri N, Farokhnia M, Abbasi SH, Kassaian SE, Noorbala Tafti AA, Gougol A, et al. A randomized, double-blind, clinical trial comparing the efficacy and safety of *Crocus sativus* L. with fluoxetine for improving mild to moderate depression in post percutaneous coronary intervention patients. *J. Affect. Disord.* 2014; 155: 216-222.
- He SY, Qian ZY, Tang FT, Wen N, Xu GL, Sheng L. Effect of crocin on experimental atherosclerosis in quails and its mechanisms. *Life Sci.* 2005; 77: 907-921.
- He SY, Qian ZY, Wen N, Tang FT, Xu GL, Zhou CH. Influence of Crocetin on

- experimental atherosclerosis in hyperlipidemic-diet quails. *Eur J Pharmacol.* 2007; 554: 191-195.
36. Sheng L, Qian Z, Zheng S, Xi L. Mechanism of hypolipidemic effect of crocin in rats: crocin inhibits pancreatic lipase. *Eur J Pharmacol.* 2006; 543: 116-122.
37. Goyal SN, Arora S, Sharma AK, Joshi S, Ray R, Bhatia J, et al. Preventive effect of crocin of *Crocus sativus* on hemodynamic, biochemical, histopathological and ultrastructural alterations in isoproterenol-induced cardiotoxicity in rats. *Phytomedicine* 2010; 17: 227-232.
38. Falsini B, Piccardi M, Minnella A, Savastano C, Capoluongo E, Fadda A, et al. Influence of saffron supplementation on retinal flicker sensitivity in early age-related macular degeneration. *Invest Ophthalmol Vis Sci.* 2010; 51: 6118-6124.
39. Mousavi SH, Tayarani NZ, Parsaee H. Protective effect of saffron extract and crocin on reactive oxygen species-mediated high glucose-induced toxicity in PC12 cells. *Cell Mol Neurobiol.* 2010; 30: 185-191.
40. Asdaq SM, Inamdar MN. Potential of *Crocus sativus* (saffron) and its constituent, crocin, as hypolipidemic and antioxidant in rats. *Appl Biochem Biotechnol.* 2010; 162: 358-372.
41. Samarghandian S, Azimi-Nezhad M, Samini F. Ameliorative effect of saffron aqueous extract on hyperglycemia, hyperlipidemia, and oxidative stress on diabetic encephalopathy in streptozotocin induced experimental diabetes mellitus. *BioMed Res. Int.* 2014; 920857.
42. Bolhassani A, Khavari A, Bathaie SZ. Saffron and natural carotenoids: Biochemical activities and anti-tumor effects. *Biochim Biophys Acta.* 2014; 1845: 20-30.
43. Papandreou MA, Tsachaki M, Efthimiopoulos S, Cordopatis P, Lamari FN, Margariti M. Memory enhancing effects of saffron in aged mice are correlated with antioxidant protection. *Behav Brain Res.* 2011; 219: 197-204.
44. Ordoudi SA, Befani CD, Nenadis N, Koliakos GG, Tsimidou MZ. Further examination of antiradical properties of *Crocus sativus* stigmas extract rich in crocins. *J Agric Food Chem.* 2009; 57: 3080-3086.
45. Ochiai T, Ohno S, Soeda S, Tanaka H, Shoyama Y, Shimeno H. Crocin prevents the death of rat pheochromyctoma (PC-12) cells by its antioxidant effects stronger than those of alpha-tocopherol. *Neurosci Lett.* 2004; 362: 61-64.
46. Selim K, Tsimidou M, Biliaderis CG. Kinetic studies of degradation of saffron carotenoids encapsulated in amorphous polymer matrices. *Food Chem.* 2000; 71: 199-206.
47. Kyriakoudi A, Tsimidou MZ. A food-grade approach to isolate crocetin from saffron (*Crocus sativus* L.) extracts. *Food Anal. Methods* 2015.
48. Everstine K, Spink J, Kennedy S. Economically motivated adulteration (EMA) of food: common characteristics of EMA incidents. *J Food Prot.* 2013; 76: 723-735.
49. Zalacain A, Ordoudi SA, Blázquez I, Díaz-Plaza EM, Carmona M, Tsimidou MZ, et al. Screening method for the detection of artificial colours in saffron using derivative UV-Vis spectrometry after precipitation of crocetin. *Food Addit Contam.* 2005; 22: 607-615.
50. COST Action FA 1101: Omics Technologies for Crop Improvement, Traceability, Determination of Authenticity, Adulteration and Origin in Saffron. www.saffronomics.org, www.cost.eu/COST_Actions/fa/Actions/FA1101.
51. Ordoudi SA, de Los Mozos Pascual M, Tsimidou MZ. On the quality control of traded saffron by means of transmission Fourier-transform mid-infrared (FT-MIR) spectroscopy and chemometrics. *Food Chem.* 2014; 150: 414-421.
52. Ordoudi SA, Cagliani LR, Lalou S, Naziri E, Tsimidou MZ, Consonni R. ¹H NMR-based metabolomics of saffron reveals markers for its quality deterioration. *Food Res. Int.* 2015; 70: 1-6.
53. Petrakis EA, Cagliani LR, Polissiou MG, Consonni R. Evaluation of saffron (*Crocus sativus* L.) adulteration with plant adulterants by (¹) H NMR metabolite fingerprinting. *Food Chem.* 2015; 173: 890-896.