

Research Article

Preharvest Treatment of 1-methylcyclopropene Reduces the Occurrence of the Water-soaking and Enhances the Quality of Watermelons (*Citrullus Vulgaris*)

Chao Zhang, Jie Huo, Wu Li, Yue Ma, Guoyi Gong, Xiaoyan Zhao and Yong Xu*

Beijing Vegetable Research Center, Beijing Academy of Agriculture and Forestry Sciences, China

*Corresponding author: Yong Xu, Beijing Vegetable Research Center, Beijing Academy of Agriculture and Forestry Sciences, Banjing, Haidian District, Beijing 100097, China, Tel: +86-10-51503656; Fax: +86-10-88446286; Email: zhangchao@nercv.org

Received: February 27, 2015; Accepted: May 11, 2015;

Published: May 14, 2015

Abstract

Exposure to exogenous ethylene is closely related to occurrence of water-soaking in watermelons. 1-Methylcyclopropene (1-MCP) inhibits the physiological action of ethylene in fruits and vegetables. Consequently, a preharvest treatment of 1-MCP was a possible option to reduce the occurrence of the water-soaking in watermelons. Two cultivars were used to validate the effect of the preharvest 1-MCP treatment at 10 days prior to their harvest. The 1-MCP treatment of 1 μ L/L enhanced the soluble solid content and fruit weight of the 2 cultivars, and lowered the water-soaking percentage and respiration rate. The 1-MCP blocked the physiological action of exogenous ethylene and slowed down the maturation of the watermelons. In summary, the preharvest 1-MCP treatment of 1 μ L/L reduced the occurrence of the water-soaking and enhanced the quality of watermelons by slowing down the maturation.

Keywords: Watermelon; 1-MCP; Preharvest treatment; Water-soaking; Exogenous ethylene

Introduction

Water-soaking of watermelons (*Citrullus vulgaris*) is characterized by tissue translucency, softening and maceration of the endocarp and placental tissues [1]. A number of factors are suspected to contribute to this syndrome [1-5]. Among the factors, the exposure to exogenous ethylene has been demonstrated to induce the water-soaking in both the immature and fully ripe fruits [1,5-7]. Exposure to exogenous ethylene is difficult to avoid during the growth of watermelons, as ethylene may be produced from other plants or fruits [8]. Therefore, inhibiting the effect of exogenous ethylene could be an effective strategy to reduce the occurrence of the water-soaking in watermelons. 1-Methylcyclopropene (1-MCP) inhibits the physiological action of ethylene in fruits and vegetables [9,10]. Postharvest treatment of 1-MCP inhibits water-soaking and extends the shelf life of fresh-cut watermelons [8,11]. Moreover, the 1-MCP treatment to watermelons benefits the commercial storage conditions (up to 3 weeks at 13°C) even with exposure to exogenously ethylene [1]. However, the effect of preharvest treatment of 1-MCP on water-soaking of watermelon fruits has not been fully explored.

The aim of the current study was to evaluate the effect of preharvest treatments of 1-MCP on water-soaking and quality of watermelon fruits. Preharvest 1-MCP treatments either 1 μ L/L or 5 μ L/L were applied 10 days prior to their harvest. Water-soaking and quality parameters of the fruits were then evaluated. Moreover, 2 cultivars, one of whom is more susceptible to the water-soaking, were used to corroborate and reinforce our results.

Materials and Methods

Plant material and cultivation

Two cultivars of watermelon (*Citrullus vulgaris*, var Jingying

series), F and M that have not been used commercially, were used in this study. The Jingying series that are round with serrated stripes are harvested 26 d after the pollination. The average weight of the fruits is 2000 g. Both cultivars were planted in August and harvested in October (2010 and 2011) at the Sijiqing Farm of the Beijing Vegetable Research Center, Beijing, China, following standard commercial practices. Two separated greenhouses (4 m \times 6 m) were used as the controls to raise the fruit number. The other 2 separated greenhouses (4 m \times 6 m) were subjected to preharvest 1-MCP treatments either 1 μ L/L or 5 μ L/L. In each green houses, the cultivar F (about 60 seeds) and M (about 60 seeds) were sown on in rows that were 3 m apart. At the six-true-leaf stage, all vines except 2 per plant were pinched to promote branching. The female flowers on the 11th node were self-pollinated by using more than 2 staminate flowers per pistil at anthesis. Only one pollinated flower was left on each of the 2 branches, and all the other female flowers were removed. The fruits were harvested after the pollination for 26 d. For each cultivar, more than 40 fruits were harvested and weighed for each treatment.

Preharvest 1-MCP treatment and experiment design

Each fruit was sealed separately in a polyethylene bag for 24 h at 10 d prior to the harvest. An appropriate weight of the 1-MCP powder (Lanzhou Jiacheng Biotechnology Ltd., Lanzhou, China) and water were sealed in a bag to obtain an initial 1-MCP concentration of either 1 μ L/L or 5 μ L/L that was validated by the GC analysis. The control-F and control-M stood the control of the cultivars F and M, respectively. The 1-MCP treatments of 1 μ L/L were named as F1 and M1 for cultivars F and M, respectively. In a similar manner, the treatments of 5 μ L/L were named as F5 and M5 for cultivars F and M, respectively.

For each treatment, more than 40 fruits were harvested. Among

the fruits, 10 fruits were used to measure the respiration rate after harvest immediately, and the remaining fruits (more than 30 fruits) were used for the other analyses. The remaining fruits were cut in half through the placenta tissues and the cross surface was evaluated for the water-soaking. The half of the fruit was peeled and juiced in a HR1861 food mixer (Philips, Dongguan, China) to acquire the watermelon juice. The juice was used to measure the Soluble Solid Content (SSC), titratable acid, and vitamin C content. The juice for the vitamin C analysis was frozen at -80°C for the future analyses. The other half of the fruits was peeled and cut into a cube with a dimension of $5\text{ cm} \times 5\text{ cm} \times 3\text{ cm}$ for the texture analysis.

Determination of vitamin C content

The vitamin C content of the fruits was determined by an HPLC method that was recently reported [12]. An aliquot of 25 mL of the juice was mixed with 25 mL of a solution containing 45 g/L of metaphosphoric acid and 7.2 g/L of dithiothreitol. The mixture was centrifuged at $20,000 \times g$ for 15 min at 4°C and the supernatant was vacuum-filtered through the Whatman No. 1 paper. The sample was then passed through a millipore $0.45\ \mu\text{m}$ membranes into an opaque vial and kept at -80°C until being used. An aliquot of $20\ \mu\text{L}$ was injected into Agilent 1200 series HPLC (Agilent Technologies, Palo Alto, California) fitted with a reverse-phase C18 Spherisorb[®] ODS2 ($5\ \mu\text{m}$) stainless-steel column ($4.6\text{ mm} \times 250\text{ mm}$). The mobile phase was a 0.01% sulphuric acid solution adjusted to a pH of 2.6. The flow rate was fixed at 1 mL/min and monitored at 245 nm at 25°C . Vitamin C was quantified using a calibration curve based on ascorbic acid pure standards and results were expressed as relative vitamin C concentration.

Determination of SSC and titratable acid

An aliquot of 1 ml juice was dropped on a pocket digital refractometer (Pal- α , ATAGO Co., Ltd., Japan) to measure the SSC. An aliquot of 10 mL juice was titrated automatically with a standardized 10 mmol/L NaOH to the end point (pH = 8.1) by a pH meter (TitroLine- SI Analytics GmbH, Mainz, Germany). The volume of NaOH (by multiplying with 0.67) was converted to mg malic acid per ml juice.

Texture analysis

The fruit cube was placed on the test platform of a TA.XT2i plus texture analyzer (Stable Micro Systems Ltd., Godalming, and Surrey, UK). The trigger (10 mm stainless cylinder) was compressed to 50 % of the total height with a pre-test speed of 5 mm/s, test speed of 1 mm/s, and post-test speed of 10 mm/s. During the compression, the force of the trigger increased quickly, and then vibrated in a small range at a certain level. The time of the vibration was defined as the statistical interval. The average force of the statistical interval indicated the firmness of fruits. The ratio of the linear distance to the time difference in the statistical interval indicated the fragility of fruits. The higher the ratio, the more fragile is the fruit.

Determination of respiration rate

Respiration rate was determined at 22°C by an infrared CO_2 analyzer (LI-6400XT, LI-COR, Inc., Lincoln, NE, USA) with a 5 L sample chamber. The fruit was sealed in the chamber. The carrier gas (O_2 of 21%, and N_2 of 79%) was used to transport the CO_2 produced by the fruit to the CO_2 analyzer. The respiration rate was recorded




Figure 1: Preharvest 1-MCP treatments on the vitamin C (A), SSC (B), titratable acid (C), and average weight (D) of watermelons. Data are means \pm standard deviation ($n \geq 60$). Means with different letter represent a significant difference ($p < 0.05$). The control-M and control-F were the controls for the cultivars F and M, respectively. The M1 and M5 were the 1-MCP treatments of $1\ \mu\text{L/L}$ and $5\ \mu\text{L/L}$, respectively for cultivars M. The similar rule was applied for the cultivars F.

when the CO_2 reached equilibrium. The results were the average of about 10 replicates per treatment.

Water-soaking of fruits

For each treatment, about 30 fruits were cut in half through the placenta tissues for the water-soaking percentage evaluation. The water-soaking percentage, which was the average percentage of the water-soaking area in the total fruit area through the placenta tissues, was determined through the visual estimation by 3 experienced panelists.

Statistical analysis

The results were the average of 2 years (2010 and 2011) that were expressed as the average \pm standard deviation ($n \geq 3$). Analysis of variance was used to compare mean differences of the results. If the differences among means were detected, multiple comparisons were performed using Duncan's Multiple Range Test. All analyses were conducted by SPSS (Window Version 19).

Results and Discussion

Preharvest treatments of 1-MCP enhanced the quality of watermelons

The vitamin C, SSC, titratable acid, and average weight of the fruits are shown in Figure 1. The data described herein were the summary of the fruits from two harvest seasons. The 1-MCP treatment showed no influence on the vitamin C content of the cultivar M, while the treatment of the 5 μ L/L enhanced the vitamin C content of the cultivar F significantly. This difference maybe resulted from the nature of the cultivars.

SSC increases with the ripening of watermelons and reaches a plateau at maturation [13], while titratable acid decreases during the maturation of watermelons [13,14]. Remarkably, the 1-MCP treatments influence the SSC in different manners that depend on the fruit type or cultivar [15]. For the 2 cultivars, the SSC of the 1 μ L/L treatment was significantly higher than that of the control, while the SSC of the 5 μ L/L treatment was similar to that of the control. Meanwhile, the 1-MCP treatment of 5 μ L/L reduced the titratable acid content of the fruits when compared to the control. The average weight of the 1 μ L/L treatment was higher than that of the control, while the average weight of the 5 μ L/L treatment was similar to that of the control. Hence, the preharvest 1-MCP treatments of 1 μ L/L raised the SSC and average weight of the fruits.

The preharvest 1-MCP treatment showed no influence on the firmness of the fruits (Figure 2). The 1-MCP treatment of 1 μ L/L showed on influence on both the fragility and hardness of the fruits. The 1-MCP treatment of 5 μ L/L reduced the fragility of the fruits compared with the control, while showed no influence on their hardness. Being similar to our results, the postharvest treatment of 1-MCP (1 μ L/L) shows no significant influence on the firmness of the fresh-cut watermelons [8].

The 1-MCP treatment of 1 μ L/L showed the similar influence on the quality parameters of the 2 cultivars, except the vitamin C content. These results validated that the preharvest 1-MCP treatment of 1 μ L/L enhanced the quality of the watermelons.




Figure 2: Preharvest 1-MCP treatments on the texture of watermelons. Data are means \pm standard deviation ($n \geq 180$). Means with different letter represent a significant difference ($p < 0.05$). The control-M and control-F were the controls for the cultivars F and M, respectively. The M1 and M5 were the 1-MCP treatments of 1 μ L/L and 5 μ L/L, respectively for cultivars M. The similar rule was applied for the cultivars F.




Figure 3: Preharvest 1-MCP treatments on the water-soaking occurrence ($n \geq 60$) (A) and the respiration rate ($n \geq 20$) (B) of watermelons. Data are means \pm standard deviation. Means with different letter represent a significant difference ($p < 0.05$). The control-M and control-F were the controls for the cultivars F and M, respectively. The M1 and M5 were the 1-MCP treatments of 1 μ L/L and 5 μ L/L, respectively for cultivars M. The similar rule was applied for the cultivars F.

Preharvest treatments of 1-MCP reduced the water-soaking and respiration rate of watermelons

The effect of the preharvest treatments of 1-MCP on the water-soaking and respiration rate of watermelons is shown in Figure 3. The preharvest 1-MCP treatment lowered the water-soaking percentage of the fruits, which was speculated as the slowing down the maturation of watermelons by 1-MCP regulation. The similar phenomenon also presents on tomato [16]. The results of the respiration rate of the fruits further confirmed our speculation. The respiration rate of the treated fruits was significantly lower than that of the control. Consequently, the preharvest 1-MCP treatment was supposed to reduce the water-soaking by slowing down the maturation of fruits, which would also be good for their postharvest storage. Furthermore, the water-soaking percentage of the 1 μ L/L treatment was lower than that of the 5 μ L/L treatment. The reason for this phenomenon is not clear. Based on previous studies, the effective concentrations of 1-MCP vary widely for different commercial purposes that differ with respect to time, temperature and the method of application [17,18].

The water-soaking percentage of the cultivar M was much higher than that of the cultivar F, which proved that the cultivar M was more susceptible to the water-soaking than the cultivar F under the same conditions.

Acknowledgement

The authors are grateful to the financial support of the earmarked fund for Modern Agro-industry Technology Research System (CARS-26-22) and Beijing Academy of Agricultural and Forestry Sciences Special Funding (KJXC20140204).

References

- Mao L, Karakurt Y, Huber DJ. Incidence of water-soaking and phospholipid catabolism in ripe watermelon (*Citrullus lanatus*) fruit: induction by ethylene

- and prophylactic effects of 1-methylcyclopropene. *Postharvest Biology and Technology*. 2004; 33: 1-9.
2. Lee SG, Shin YA, Kim KY, Chung JH, Lee YB. Effect of rootstocks on the growth, fruit quality and ethylene evolution from harvested fruits in watermelon. *Journal of Korean Society Horticultural Science*. 1998; 39: 238-241.
 3. Cho M, Hurr B, Jeong J, Lim C, Huber D. Postharvest senescence and deterioration of 'Thoroughbred' and 'Carlo' green beans (*Phaseolus vulgaris* L.) in response to 1-methylcyclopropene. *HortScience*. 2008; 43: 427-430.
 4. Lee SG, Ko KD. Ethephon application induces symptoms of fruit tissue degeneration in watermelon. *Journal of Plant Biology*. 2008; 51: 337-340.
 5. Hurr MB, Huber JD, Vallejos EC, Lee E, Sargent AS. Ethylene-induced overproduction of reactive oxygen species is responsible for the development of water-soaking in immature cucumber fruit. *J Plant Physiol*. 2013; 170: 56-62.
 6. Karakurt Y, Huber DJ. Ethylene-induced gene expression, enzyme activities, and water soaking in immature and ripe watermelon (*Citrullus lanatus*) fruit. *J Plant Physiol*. 2004; 161: 381-388.
 7. Hurr MB, Huber JD, Vallejos EC. Features of programmed cell death precede water-soaking development in ethylene-treated immature cucumber fruit. *Postharvest Biol Technol*. 2010; 58: 13-20.
 8. Zhou B, McEvoy JL, Luo Y, Saftner RA, Feng H, Beltran T. 1-Methylcyclopropene counteracts ethylene-induced microbial growth on fresh-cut watermelon. *J Food Sci*. 2006; 71: M180-M184.
 9. Sisler E, Serek M. Inhibitors of ethylene responses in plants at the receptor level: recent developments. *Physiologia Plantarum*. 1997; 100: 577-582.
 10. Watkins C, Miller W. 1-methylcyclopropene (1-MCP) based technologies for storage and shelf life extension. *Acta Hort*. 2005; 687: 217-224.
 11. Mao L, Jeong J, Que F, Huber DJ. Physiological properties of fresh-cut watermelon (*Citrullus lanatus*) in response to 1-methylcyclopropene and post-processing calcium application. *J Sci Food Agri*. 2006; 86: 46-53.
 12. Oms-Oliu G, Odriozola-Serrano I, Soliva-Fortuny R, Martin-Belloso O. Effects of high-intensity pulsed electric field processing conditions on lycopene, vitamin C and antioxidant capacity of watermelon juice. *Food Chemistry*. 2009; 115: 1312-1319.
 13. Corey AK, Schlimme VD. Relationship of rind gloss and groundspot color to flesh quality of watermelon fruits during maturation. *Scientia Horticulturae*. 1988; 334: 211-218.
 14. Bangalore DV, McGlynn WG, Scott DD. Effects of fruit maturity on watermelon ultrastructure and intracellular lycopene distribution. *J Food Sci*. 2008; 73: S222-S228.
 15. Watkins CB, Nock JF, Whitaker BD. Responses of early, mid and late season apple cultivars to postharvest application of 1-methylcyclopropene (1-MCP) under air and controlled atmosphere storage conditions. *Postharvest Biology and Technology*. 2000; 19: 17-32.
 16. Wills RBH, Ku VVV. Use of 1-MCP to extend the time to ripen of green tomatoes and postharvest life of ripe tomatoes. *Postharvest Biology and Technology*. 2002; 26: 85-90.
 17. Jiang Y, Joyce DC. Effects of 1-methylcyclopropene alone and in combination with polyethylene bags on the postharvest life of mango fruit. *Annals of Applied Biology*. 2002; 173: 321-327.
 18. Blankenship MS, Dole MJ. 1-Methylcyclopropene: a review. *Postharvest Biology and Technology*. 2003; 28: 1-25.