

Special Article – Food Safety

First Report of Human Gastroenteritis Caused by *Escherichia Coli* O157:H⁻

Santos RFC¹, Nascimento JDS², Geimba MP³, Hessel CT⁴ and Tondo EC^{4*}

¹Superintendency of Sanitary Surveillance, Inspection and Nutrition in Food of the City Hall of the City of Rio de Janeiro, (SMS/SUBVISA/SVFSAN), Brazil

²Laboratory of Microbiology, Federal Institute of Education, Science and Technology-Rio de Janeiro Campus, Brazil

³Institute of Basic Health Sciences, Federal University of Rio Grande do Sul (UFRGS), Brazil

⁴Institute of Food Science and Technology, Federal University of Rio Grande do Sul (ICTA/UFRGS), Brazil

*Corresponding author: Eduardo Cesar Tondo, Instituto de Ciência e Tecnologia de Alimentos, Universidade Federal do Rio Grande do Sul (ICTA/UFRGS), Porto Alegre, RS, Brazil

Received: November 08, 2016; Accepted: January 16, 2017; Published: January 20, 2017

Introduction

Escherichia coli are a broad group of bacteria including harmless commensals up to highly pathogenic strains [1]. Among them, Enterohaemorrhagic *Escherichia coli* (EHEC) constitute a subset of serotypes of Shiga-Toxin (Stx)-Producing *E. coli* (STEC) firmly associated with severe human illnesses like bloody diarrhea and Hemolytic Uremic Syndrome (HUS). STEC/EHEC strains have been isolated from several foods, including an increasing number of unusual food vehicles associated with human infections [2-5]. Currently, the most common STEC associated with human diseases is *E. coli* O157:H7 [6] and the identification of the H7 flagellar antigen is critical for the confirmation of this strain, however, some clinical isolates of *E. coli* O157 are Nonmotile (NM) and do not produce detectable H antigen [7], being designated *E. coli* O157:NM or *E. coli* O157:H⁻.

E. coli O157: H⁻ was first recognized as food pathogen in 1988, during an HUS outbreak, in Bavaria, Germany [8]. The strain involved with this outbreak had *stx2* gene, responsible for the expression of Shiga-toxin, was non-motile and was able to ferment sorbitol. The finding of this uncommon O157 strain causing HUS in humans motivated in-deep studies. For example, in Germany, during the period between 1988 and 1991 [9], calculated the percentage of cases of *E. coli* O157:H⁻ responsible for diarrhea and HUS and have shown 7.4 to 25% and 13.3 to 40.5% of the cases of these two syndromes were caused by *E. coli* O157:H⁻, respectively. Also in Bavaria, during the winter of 1995/1996, one big outbreak was caused by *E. coli* O157:H⁻, resulting in 28 children-cases of HUS and 3 fatal victims [10]. The mechanisms of pathogenicity of *E. coli* O157:H⁻ are not completely understood and some hypotheses raised. For example, in Austria [8], suggested that Shiga-toxin production was not essential for pathogenicity of *E. coli* O157 in humans. Similarly, in Germany [11],

Abstract

In September 2005, Sanitary Surveillance Service of the Municipality of Rio de Janeiro, Brazil, investigated a gastroenteritis case involving a 13-year-old teenager hospitalized due to bloody diarrhea and severe abdominal pain. Special attention was given due to the severity of symptoms and an epidemiological investigation was conducted in two states of Brazil. *Escherichia coli* O157:H⁻ was isolated from stools and from a tomato salad added with cheese prepared in the canteen of high school where the teenager studied. This is the report of the first case of gastroenteritis related to *E. coli* O157:H⁻ in Brazil.

Keywords: Foodborne disease; Enteropathogenic *Escherichia coli*; Shiga-toxin

have reported that serotypes H7 and H⁻ may be involved in foodborne outbreaks even without a uniform production of Shiga-toxin. Other study reported that some strains of *E. coli* O157:H⁻ may lost *stx* genes during infections, isolation and cultures and others strains of *E. coli* O157:H⁻ that are Shiga-toxin positive evolved from non-Shiga-toxin producing *E. coli* O157:H⁻ strains, acquiring *stx* genes from bacteriophages [9]. Shiga-toxin production generally occurs due to the induction of *stx* genes after stress conditions, mainly oxidative stress [12]. Karch and Bielaszewska [9] have suggested that *E. coli* O157:H⁻ is frequent involved in foodborne outbreaks during cold periods of the year, mainly involving children <3 years-old. Bielaszewska [13] also demonstrated that *E. coli* O157:H⁻ was isolated from dairy cattle, indicating that these kind of animals can be natural reservoirs, just as they are for *E. coli* O157:H7. Nielsen, *et al.* [14] reported that sorbitol fermenting and Shiga-toxin producing *E. coli* O157:H⁻ are rare but emergent in Europe and syndromes produced by these strains frequently result in HUS. Brandal *et al.* [15] have reported that *E. coli* O157:H⁻ cases were frequent in Europe, but its reservoir and exposure routes remained unknown. Sallam *et al.* [16] demonstrated that sorbitol fermenter *E. coli* O157:H⁻ emerged as an important foodborne pathogen in Africa and this microorganism has been responsible for HUS cases, even with higher rates than non-fermenting sorbitol *E. coli* O157:H7.

In Brazil, *E. coli* O157:H7 has been rarely isolated, however few studies demonstrated its presence in bovine feces [17,18] and recently, in ground meat [19] and on bovine carcasses [20]. Additionally, there were only two registered gastroenteritis cases attributed to *E. coli* O157:H7 after the consumption of rare red meat, in 2001, in Campinas/SP [21], but there is no register of human gastroenteritis involving *E. coli* O157:H⁻. The aim of this study was to describe the first case of human foodborne disease caused by *E. coli* O157:H⁻ in Brazil.

Materials and Methods

Outbreak investigation

The foodborne outbreak was investigated by the Sanitary Surveillance Service of the municipality of Rio de Janeiro, Brazil, following recommendations of Pan-American Health Organization/World Health Organization [22]. Sanitary Surveillance Service of the State of Minas Gerais helped Sanitary Surveillance Service of Rio de Janeiro in epidemiological investigation, because the victim and involved people had travelled to Minas Gerais some days before the burden of symptoms.

Notification and general investigation procedures

The outbreak notification occurred by telephone (September 13, 2005) informing a case of severe diarrhea to the Technical Center of Sanitary Surveillance Service of Rio de Janeiro (RJ), starting routine foodborne investigation. Officers were mobilized to carry out in-depth investigation with the victim and involved persons and to collect stool and food samples, if possible. Case patient and controls were interviewed personally or by telephone using routine questionnaire for foodborne outbreaks investigation. Hypothesis-generating questions were done in order to identify commonly consumed foods and places of consumption. Suspect establishments were visited in Rio de Janeiro and Minas Gerais in order to verify Good Hygiene Practices (GHP), Standard Operating Procedures (SOP) and adherence to the Brazilian food safety regulation. Food handlers were interviewed regarding food preparation and recent illnesses. Stools and suspect food samples were collected and sent to the Reference Laboratory of Public Health of Rio de Janeiro, Noel Nutels in order to be analyzed following methods described by FDA/CFSAN [23]. The identification of bacterial isolates was carried out in Fundacao Oswaldo Cruz, Rio de Janeiro as follow:

Microbiological testing

Stool samples (10 g) were homogenized in 90 ml peptone water diluent. After that, serial tenfold dilution was prepared using the same diluents, according to the expected level of contamination. 0.1 ml of each dilution in duplicate was spread onto dried surface of sorbitol-MacConkey (SMAC) agar (Oxoid, England). The plates were incubated at 35°C and read after 18 h. Food samples (25 g) were homogenized in 225 ml of modified Trypticase Soy Broth (mTSB, Oxoid, England) containing 0.45 mg of novobiocin (Laborclin, Brazil) and incubated at 37°C for 24 h. After incubation, 0.1 ml of enrichment broth were spread onto Tellurite-Cefixime-Sorbitol MacConkey Agar (TC SMAC, Potassium tellurite 2.50 mg/liter and Cefixime 0.05 mg/liter) and incubated at 37°C for 24 h. Sorbitol-negative colonies (pale colonies) were picked from TC SMAC and characterized by biochemical tests (indole and beta-glucuronidase assay). Motility was carried out successively growing suspect isolates in semisolid agar (concentrations of 0.2% to 0.8%) according to Ewing [24]. Non-motile, indole-positive and beta-glucuronidase-negative isolates were tested for agglutination with *E. coli* O157 antiserum in the Reference Laboratory of *Enterobacteriaceae* in Fundacao Oswaldo Cruz-FIOCRUZ, following methods of FDA/CFSAN [21] and Orskov and Orskov [25]. O157-positive isolates were further analyzed by tissue culture assay for verotoxins (Shiga-like toxins) [23] as described below. Non-O157 STEC isolates were not investigated.

Tissue culture assay for verotoxins

Bacterial culture preparation: *E. coli* O157 isolates were inoculated in 20 ml Trypticase Soy Broth (TSB) and incubated at 37°C for 20-24 h. Bacterial cultures were centrifuged at 7000xg for 30 min and the supernatant was filtered through 0.45 µm membrane in order to remove residual cells. The filtrate was stored at 4°C and before use, was 1:5 diluted in Dulbecco's phosphate-buffered saline (DPBS), pH 7.0.

Preparation of vero monolayer's: Vero culture was maintained in Eagle's minimal essential medium (MEME-L15) containing 2% milk serum added of gentamicin sulfate (50 µg/ml). Before use, the purity of culture was checked in 5% CO₂ incubator held at 36°C for 72 h. Normal cells were treated with trypsin to remove monolayer. Cells were suspended to the density of 10⁵ per ml in growth medium and portions of 0.5 ml were transferred to 16 mm wells in sterile plastic dishes. Cells were incubated for 3-4 days at 36°C in CO₂ incubator. Growth medium was removed and replaced with 0.5 ml fresh medium.

Toxicity test

Portions of 0.05 ml of diluted culture filtrates were transferred to wells and incubated for 4 days at 36°C in CO₂ incubator. Cytopathic effect was examined daily for rounded and shriveled cells and detachment). Dilute TSB 1:5 in DPBS was used as control [23].

Results

Foodborne outbreak description

At night of September 13, 2005, a 13-years-old boy was hospitalized in the Pediatric Medical Center of Barra da Tijuca, Rio de Janeiro (RJ), resulting in the notification of the case. Medical reports demonstrated intense abdominal pain and severe diarrhea. Computerized tomography and ultra-sound revealed mesenteric adenitis. Clinical symptoms evolved to bloody diarrhea, cramps and one emetic episode. Intravenous antibiotic treatment was used each 12 hours (Rocefin, Roche). At September 16, 2005, the patient returned to home. Coproculture revealed a sorbitol-negative, indole positive, non-motile *E. coli*. The strain agglutinated O157 antiserum, being identified as *E. coli* O157:H⁻.

The victim declared that the week before the burden of symptoms, had lunch and consumed different meals at the canteen of school where he studied, at the city of Jacarepagua, RJ. The canteen was served by an outsourced catering service. At home, the meals were composed mainly by red meat, fruits and vegetables, purchased in three commercial food establishments. The day before the beginning of symptoms, the teenager and his family had lunch at the restaurant of the residential condominium where they lived, at Barra da Tijuca, RJ. The dairy routine of victim was going to school, play with friends at residential condominium, play football and swim in the swimming pool. The family did not get sick even eating the same foods and meals of teenager, at the same restaurant and at home, except at school canteen.

Five days before get sick, the victim traveled to Tiradentes city, State of Minas Gerais (from 08 to 10/09/2005), in a travel organized by school. He had several meals at different restaurants of diverse cities, eating barbecue (including rare bovine meat), pizza, snacks,

lettuce, pepper, homemade ice cream and water from a font in a touristic city. Teachers and other students that also were in the tour were interviewed in order to identify possible consumption of foods commonly linked to transmission of *E. coli* O157 (rare red meat, non-potable water, fruits and vegetables). Among the 62 people that were in the tour, 58 (52 teenagers and 6 six adults) were interviewed. Five teenagers declared gastrointestinal symptoms after travel. One of them reported intense watery diarrhea, low fever, prostration, body pain and abdominal pain during four days. There were no nausea or vomit episodes. The same teenager had sore throat with oropharyngeal hyperemia. During the travel, students have eaten hamburgers, snacks, rare red meat, vegetables, fruits, non-potable water, homemade ice cream and Minas's type cheese.

At September 16, 2005, Sanitary Surveillance Service of the State of Minas Gerais sent a technical note and a sanitary alert informing about the foodborne case notified at Rio de Janeiro and the possible involvement of food establishments where the group had their meals during the travel to Minas Gerais. Official inspections were carried out in the establishments of Minas Gerais, as well. Inspections were done aiming at the verification of GHP and SOP, traceability of raw material and manipulation of suspect foods, as hamburgers and unpasteurized cheeses. Sanitary officers did not find important irregularities inside food establishments, turning the attention to food contamination inside the canteen school at Rio de Janeiro city where teenagers had meals before travel.

Based on this, several foods served at school canteen were sampled and sent to the Reference Laboratory of Public Health of Rio de Janeiro Noel Nutels. A Tomato salad added with Minas type cheese demonstrated inadequate sensorial characteristics and counts of thermo tolerant coli forms above the limit established by Brazilian Microbiological Standard regulation [26]. The sampling was carried out some days after the beginning of victim symptoms and the analysis demonstrated *E. coli* O157:H⁻, *stx* gene negative. Based on the results, this food was considered the most probable vehicle of the foodborne disease. Additional sampling of Minas type cheese was carried out at supermarket where the cheese was purchased, but *E. coli* O157:H⁻ was not found. Other sources of contamination also could be tomato used for salad or food handlers, but these were not investigated.

Discussion

This study describes a foodborne outbreak occurred in Rio de Janeiro, Brazil, with characteristics indicating a foodborne illness caused by *E. coli* O157:H⁻. The incubation time, victim's symptoms and the isolation of *E. coli* O157:H⁻ from the patient stool and from a tomato salad added of a regional cheese prepared in a food establishment where the victim had made some meals strongly indicated that the foodborne outbreak was caused by *E. coli* O157:H⁻. An interesting fact was that the *E. coli* O157:H⁻ isolated from patient's coproculture was verotoxin-positive, while the strain isolated from the tomato salad did not produce verotoxin in culture tissue assay. Some possible explanations for these results are the loss of *stx* genes or the incapacity of expression of these genes outside of intestine. For example, Themphachana, et al. [27] have reported a *stx*-negative *E. coli* O157 causing 228 cases of diarrhea in Thailand and suggested that the causative agent was originally *stx*-positive, but lost the gene after establishing infection. Similar results were demonstrated

by Whatahiki, et al. [28] who characterized strains of *E. coli* O111 and O157 isolated from raw beef dishes responsible for 181 infected patients and five deaths, in Japan. *E. coli* O157 isolated from stools demonstrated diverse *stx* gene profiles and molecular analyses indicated that isolates originated from a single clone. The same authors suggested that *E. coli* O111 *stx*2-positive converted in *stx*-negative during infection. Recently, Licznarska, et al. [12] reported that the induction of *stx* genes of some *E. coli* O157 only happens during human infections, mainly inside intestine, explaining why some strains have the *stx* genes, but do not produce Shiga-toxin during *in vitro* tests.

Based on epidemiological investigation and laboratorial results, Sanitary Surveillance Service of Rio de Janeiro assumed that the foodborne outbreak was caused by *E. coli* O157:H⁻. In order to avoid new illness cases, officers carried out new inspections in the food establishments where the victim used to buy foods, i.e. restaurant of the residential condominium where teenager lives, his school and supermarkets next to his home. Further, in the school, a discussion session was done with students, focusing on clinical symptoms and most probable ways of contamination of *E. coli* O157. Guideline information about food consumption in trips was sent to school, as well.

In Brazil, *E. coli* O157 is not routinely investigated by reference laboratories, because this food pathogen is not frequently found on foods or causing reported illnesses. However, there is the possibility of occurrence of non-notified cases caused by this microorganism. The lack of reported cases in Brazil can be reflect of the low prevalence of *E. coli* O157 on foods [29,30] or because this microorganism has not been investigated routinely by food microbiology laboratories, making difficult to know the true incidence of *E. coli* O157 in Brazil.

Even though information about *E. coli* O157 is not abundant in Brazil, the importance of this food pathogen is well recognized and based on this fact Brazil has started to implement monitoring programs and control measures in order to prevent *E. coli* O157 illnesses. For example, the Ministry of Agriculture, Livestock and Food Supply published in 17, July 2015, the regulation DIPOA/SDA 01, approving sampling and analytical procedures for investigation of verotoxigenic *E. coli* (serovars O26, 045, O103, O111, O121, O145 and O157) in raw beef products. This regulation will increase database about real prevalence of enteropathogenic *E. coli* serovars and will contribute to food safety of Brazilian food products.

Conclusion

The present study reported the first foodborne disease human case caused by *E. coli* O157:H⁻ case in Brazil. The isolation of the same microorganism in a tomato salad added with Minas type cheese prepared in the same local where the victim had lunch some days before the burden of symptoms, strongly indicated that the source of contamination was the canteen of school where the victim studied.

References

1. Bodouei MA, Morabito S, Najaffar A, Mazendarani E. Molecular characterization of enterohemorrhagic Escherichia coli hemolysin gene (EHEC-hlyA)-harboring isolates from cattle reveals a diverse origin and hybrid diarrheagenic strains. *Infection, genetics and evolution*. 2016; 39: 342-348.
2. Majowicz S, Scallan E, Jones-Bitton A, Saggert J, Stapleton J, Angulo F.

- Global Incidence of Human Shiga Toxin-Producing *Escherichia coli* Infections and Deaths: A Systematic Review and Knowledge Synthesis. *Foodborne Pathog Dis.* 2014; 11: 477-485.
3. Khalil, RK, Frank, JF. Behavior of *Escherichia coli* O157:H7 on Damaged Leaves of Spinach, Lettuce, Cilantro and Parsley Stored at Abusive Temperatures. *J Food Prot.* 2010; 73: 212-220.
 4. Caprioli A, Morabito S, Brugere H, Oswald E. 2005. Enterohaemorrhagic *Escherichia coli*: emerging issues on virulence and modes of transmission. *Vet Res.* 2005; 36: 289-311.
 5. Rangel J, Sparling P, Crowe C, Griffin P, Swerdlow D. Epidemiology of *Escherichia coli* O157:H7 Outbreaks, United States, 1982-2002. *Emerging and Infection Diseases.* 2005; 11.
 6. Gould LH, Mody RK, Ong KL, Clogher P, Cronquist AB, Garman KN, et al. Increased recognition of non-O157 Shiga toxin-producing *Escherichia coli* infections in the United States during 2000-2010: Epidemiologic features and comparison with *E. coli* O157 infections. *Foodborne Pathogens and Disease.* 2013; 10: 453-460.
 7. Fields PI, Blom K, Hughes HJ, Hesel LO, Feng P, Swaminathan B. Molecular Characterization of the Gene Encoding H Antigen in *Escherichia coli* and Development of a PCR-Restriction Fragment Length Polymorphism Test for Identification of *E. coli* O157:H7 and O157:NM. *Journal of Clinical Microbiology.* 1997; 1066-1070.
 8. Allerberger F, Dierich MP, Gruber-Moesenbacher U, Liesegang A, Prager R, Hartmann G, et al. Nontoxigenic sorbitol-fermenting *Escherichia coli* O157:H- associated with a family outbreak of diarrhea. *Wien Klin Wochenschr.* 2000; 112: 846-850.
 9. Karch H, Bielaszewska M. Sorbitol-Fermenting Shiga Toxin Producing *Escherichia coli* O157:H- Strains: Epidemiology, Phenotypic and Molecular Characteristics and Microbiological Diagnosis. *J Clin Microbiol.* 2001; 39: 2043-2049.
 10. Ammon A, Petersen LP, Karch H. A large outbreak of hemolytic uremic syndrome caused by an unusual sorbitol-fermenting strain of *E. coli* O157:H-. *J Infect Dis.* 1999; 179: 1274-1277.
 11. Schmidt H, Scheef J, Huppertz HI, Frosch M, Karch H. *Escherichia coli* O157:H7 and O157:H- strains that do not produce Shiga toxin: phenotypic and genetic characterization of isolates associated with diarrhea and hemolytic-uremic syndrome. *J Clin Microbiol.* 1999; 37: 3491-3496.
 12. Licznarska K, Falenczyk BN, Bloch S, Dydecka A, Topka G, Gdsior T, et al. Oxidative Stress in Shiga Toxin Production by Enterohemorrhagic *Escherichia coli*. *Oxid Med Cell Longev.* 2016; 1-8.
 13. Bielaszewska M, Schmidt H, Liesegang A, Prager R, Rabsch W, Tschape H, et al. Cattle can be a reservoir of sorbitol-fermenting shiga-toxin-producing *Escherichia coli* O157:H-strains and a source of human diseases. *J Clin Microbiol.* 2000; 38: 3470-3473.
 14. Nielsen S, Frank C, Fruth A, Spode A, Prager R, Graff A, et al. Desperately seeking diarrhoea: outbreak of haemolytic uraemic syndrome caused by emerging sorbitol-fermenting shiga toxin-producing *Escherichia coli* O157:H-, Germany. 2009. *Zoonoses Public Health.* 2011; 58: 567-572.
 15. Brandal LT, Lobersli I, Stavnes TI, Wester AI, Lindstedt BA. First Report of the Shiga Toxin 1 Gene in Sorbitol-Fermenting *Escherichia coli* O157:H-. *J Clin Microbiol.* 2012; 50: 1825-1826.
 16. Sallam Ki, Mohammed MA, Ahdy AM, Tamura T. Prevalence, genetic characterization and virulence genes of sorbitol-fermenting *Escherichia coli* O157:H-and *Escherichia coli* O157:H7 isolated from retail beef. *Int J Food Microbiol.* 2013; 165: 295-301.
 17. Cerqueira AM, Guth BE, Joaquim RM, Andrade Jr. High occurrence of Shiga toxin-producing *Escherichia coli* (STEC) in healthy cattle in Rio de Janeiro State, Brazil. *Vet Microbiol.* 1999; 70: 111-121.
 18. Irino K, Kato Ma, Vaz Tm, Ramos II, Souza Ma, Cruz AS. Serotypes and virulence markers of Shiga toxin producing *Escherichia coli* (STEC) isolated from dairy cattle in Sao Paulo State, Brazil. *Vet microbiol.* 2005; 105: 29-36.
 19. Paula CMD, Casarin LS, Tondo EC. *Escherichia coli* O157:H7-patogeno alimentar emergente. *Vig Sanit Debate.* 2014; 2: 23-33.
 20. Loiko MR, Paula CMD, Langone ACJ, Rodrigues RQ, Cibulski S, Rodrigues RO, et al. Genotypic and antimicrobial characterization of pathogenic bacteria at different stages of cattle slaughtering in southern Brazil. *Meat Science.* 2016; 116: 193-200.
 21. Centro De Vigilancia Epidemiologica. Casos confirmados e coeficientes de incidencia de casos autoctones de doencas de notificacao compulsoria no estado de Sao Paulo, no periodo de 1998 a 2008. 2013.
 22. OPAS/WHO. Procedimientos para la Investigacion de Enfermedades transmitidas por alimentos. Manual de la International Association of Milk, Food and Environmental Sanitarias Inc. Publicacion Cientifica N° 367. 2nd edn. 1978.
 23. FDA/CFSAN. Isolation Methods for Enterohemorrhagic *E. coli* O157:H7. In: *Bacteriological Analytical Manual. Food and Drug Administration, Centers for Food Safety and Applied Nutrition.* 2001.
 24. Ewing WH. Edwards and Ewing's Identification of Enterobacteriaceae, 4th edn. Elsevier Science Publishing Co., Inc., New York. 1986; 536.
 25. Orskov F, Orskov I. Serotyping of *Escherichia Coli*. *Methods on Microbiology.* 1984; 14: 43-112.
 26. Brasil. Agencia Nacional de Vigilancia Sanitaria. Resolucao RDC n° 12 de 02 de janeiro de 2001. Aprova o Regulamento Tecnico sobre Padroes Microbiologicos para Alimentos. *Diario Oficial da Republica Federativa do Brasil, Brasilia, Janeiro de.* 2001.
 27. Themphachana M, Nakaguchi Y, Nishibuchi M, Seto K, Rattanachuy P, Singkhamanan K, et al. First report in Thailand of a stx-negative *Escherichia coli* O157 strain from a patient with diarrhea. *Southeast Asian Journal of Tropical Medicine and Public Health.* 2014; 45: 881-889.
 28. Watahiki M, Isobe J, Kimata K, Shima T, Kanat J, Shimizu M, et al. Characterization of enterohemorrhagic *Escherichia coli* O111 and O157 strains isolated from outbreak patients in Japan. *J Clin Microbiol.* 2014; 52: 2757-2763.
 29. Rodrigues R, Loiko M, Paula CMD, Hessel CT, Jacxsens L, Uyttendaele M, et al. Microbiological contamination linked to implementation of good agricultural practices in the production of organic lettuce in Southern Brazil. *Food Control.* 2014.
 30. Ceuppens S, Hessel CT, Rodrigues RQ, Bartz S, Tondo Ec, Uyttendaele M. Microbiological quality and safety assessment of lettuce production in Brazil. *Int J Food Microbiol.* 2104; 181: 67-76.