

Special Article - Role of Antioxidants

Antioxidant Activity Assay of Phenolic Compounds Isolated from *Origanum Onites* L. Aromatic Water by High Performance Liquid Chromatography

Aybastier Ö and Demir C*

University of Uludag, Department of Chemistry, Turkey

*Corresponding author: Cevdet Demir, University of Uludag, Department of Chemistry, Faculty of Science and Arts, 16059 Bursa, Turkey

Received: June 22, 2017; Accepted: July 20, 2017;

Published: July 27, 2017

Abstract

A High Performance Liquid Chromatographic (HPLC) method was developed to investigate the antioxidant performance of *Origanum onites* L. aromatic water and its phenolic compounds in terms of the free radical scavenging activity in Fenton reaction. The remaining concentrations of carvacrol, thymol and thymoquinone were measured after the oxidation reaction by HPLC to evaluate the scavenging potential of these compounds for reactive hydroxyl radicals. The main oxidation product of thymoquinone showed the highest scavenging activity in Fenton reaction. The aromatic water of *Origanum onites* L. could be used as a rich source of natural antioxidants with potential applications in pharmaceutical industry.

Keywords: Fenton reaction; Hydroxyl radical; Carvacrol; Thymol; Thymoquinone; HPLC

Introduction

Phenolic compounds such as carvacrol, thymol and thymoquinone have been reported to exhibit a wide range of biological effects, including analgesic, anti-inflammatory, protection of organs against oxidative damage induced by a variety of free radical generating agents [1,2]. Carvacrol and its isomer thymol are one of the main components of the essential oils of *Labiatae* (*Laminaceae*) members like oregano, thyme and savory [3]. Thymoquinone is the main constituent of *Nigella sativa* essential oil from most sources [4]. Thymoquinone is also the main oxidation product of carvacrol and thymol. These compounds inhibit lipid peroxidation, thus exerting their effects as antioxidant and free radical scavengers [5]. The scavenging properties of phenolic compounds are generally accepted to be due to aromatic hydroxyl substituents.

Antioxidants can react by depleting molecular oxygen or decreasing its local concentration, scavenging chain initiating radicals like hydroxyl HO•, alkoxyl RO• or peroxy ROO• and breaking the chain of radical sequence [6]. The chain-breaking antioxidants are able to scavenge free radicals due to their hydrogen donating ability with subsequent stabilization of the resulting phenoxyl radical [7]. It has been reported that carvacrol, thymol and thymoquinone have free radical scavenging activity and good reducing power [8,9]. The reducing properties of phenolic compounds are generally associated with the presence of reductones, which have been shown to exert protective action of antioxidant due to inhibition of free radical induced chain reaction through the donation of hydrogen atom. The antioxidant ability of carvacrol and thymol and their mixture has been evaluated taking into account their capacity to protect Caco-2 cells against a further exposure to H₂O₂. It was found that carvacrol and thymol possess a strong free radical scavenging ability during oxidative stress [10]. Hydroxyl free radicals are highly reactive and have short half-lives. Furthermore, direct detection

of their concentrations in natural systems is inadequate with ESR spectroscopy. Spin-trapping is a chemical reaction that provides an approach to help overcome this problem [11].

Materials and Methods**Chemicals and reagents**

Carvacrol, thymol, thymoquinone, catalase and H₂O₂ were purchased from Sigma (Sigma-Aldrich Company, USA). Acetonitrile and methanol were obtained from Merck. FeSO₄·5H₂O was purchased from Riedel-de Haën (Seelze, Germany).

Quantification of oxidation products

The scavenging capacity of carvacrol, thymol and thymoquinone to hydroxyl radical was followed and remaining concentrations of compounds after oxidation were quantified by HPLC. The reaction mixture contained the following reagents for carvacrol and thymol separately: 400µM carvacrol, 400µM thymol, 200µM H₂O₂ and 100µM FeSO₄·5H₂O in water. The reaction mixture contained 40µM thymoquinone, 20µM H₂O₂ and 10µM FeSO₄·5H₂O to monitor the thymoquinone Fenton reaction. Because of the high absorption, the concentration of reaction mixture for thymoquinone is ten times lower than carvacrol and thymol. The reaction solutions were incubated at 37°C for each phenolic compound. The Fenton reaction was stopped by adding 50µL 0.6mg/mL catalase. The quantification of compounds was performed by HPLC at wavelength 273nm for standard carvacrol and thymol and 254nm for thymoquinone. Peaks were identified based on comparison of retention times and UV spectra with standards of carvacrol, thymol and thymoquinone.

HPLC-DAD conditions

An Agilent Technologies 1200 HPLC system (Waldbronn, Germany), consisting of a vacuum degasser, binary pump, autosampler and a diode-array detector, was used for determination of phenolic compounds. Chromatographic separations were carried out




Figure 1: The chemical structures of carvacrol, thymol and thymoquinone.




Figure 2: (a) HPLC chromatogram of carvacrol at the beginning of the oxidation reaction. (b) HPLC chromatogram of carvacrol after 10 min of oxidation reaction. Reaction conditions: 400 μ M carvacrol; 200 μ M H₂O₂; 100 μ M Fe²⁺; temperature 37°C; wavelength 254 and 273nm.

using an XBridge C18 (4.6 \times 250 mm, 3.5 μ m) column from Waters. Mobile phase consists of water and acetonitrile (58:42, v/v). Total run time is 18 min. Flow rate was 0.5mL/min and injection volume was 20 μ L. Data acquisition and preprocessing was done with Chemstation for LC (Agilent Technologies).

Results and Discussion

Hydroxyl radical is a highly aggressive radical species resulted from Fenton-type reactions, responsible for the oxidative damage of most biomolecules. Antioxidants prevent hydroxyl radical occurrence and act as radical scavenger in antioxidative defense systems. Antioxidant compounds present in those systems maintain an important balance between the formation of radicals and their removal. The phenolic compounds can scavenge free radicals by rapid donation of the hydrogen atom to radical form. In this study, the hydroxyl radical scavenging reaction was based on the oxidation of phenolic compounds with Fenton reaction at 37°C. The scavenging of hydroxyl radical was followed by HPLC measurements of carvacrol, thymol and thymoquinone during the oxidation reactions. The chemical structures of these compounds are shown in Figure 1.

The Fenton reactions for carvacrol (Figure 2a) and thymol (Figure 3a) were followed at 400 μ M initial concentrations. HPLC measurements revealed that carvacrol and thymol transformed to thymoquinone by catalytic transformation after 10 min oxidation (Figures 2b & 3b). It has been reported that oxidation of carvacrol




Figure 3: (a) HPLC chromatogram of thymol at the beginning of the oxidation reaction. (b) HPLC chromatogram of thymol after 10 min of oxidation reaction. Reaction conditions: 400 μ M thymol; 200 μ M H₂O₂; 100 μ M Fe²⁺; temperature 37°C; wavelength 254 and 273nm.




Figure 4: (a) HPLC chromatogram of thymoquinone at the beginning of the oxidation reaction. (b) HPLC chromatogram of thymoquinone after 10 min of oxidation reaction. Reaction conditions: 40 μ M thymoquinone; 20 μ M H₂O₂; 10 μ M Fe²⁺; temperature 37°C; wavelength 254nm.

and thymol in the presence H₂O₂ with different catalysts yielded thymoquinone as a main product [12].

The oxidation reaction of thymoquinone was followed at 40 μ M initial concentration (Figure 4a). Thymoquinone reaction was performed at a concentration 10-fold lower than other compounds because of the higher molar absorptivity. The molar absorptivity

coefficient of thymoquinone ($20266\text{L mol}^{-1}\text{cm}^{-1}$) is 26 times higher than carvacrol ($781\text{L mol}^{-1}\text{cm}^{-1}$) and 33 times higher than thymol ($621\text{L mol}^{-1}\text{cm}^{-1}$) at 254nm and approximately 3 times higher at 273nm ($2031\text{L mol}^{-1}\text{cm}^{-1}$). Thymoquinone was also transformed into small-unidentified compounds during the oxidation reaction (Figure 4b).

The increase in the absorbance for carvacrol and thymol revealed that thymoquinone and other compounds developed during the oxidation process could have remarkable antioxidant abilities and probably play a crucial antioxidative role. These results are in agreement with the literature [13]. According to Jukic and Milos, when exposed to light for five days, thymoquinone is gradually converted to dithymoquinone, thymohydroquinone and some other derivatives [14]. Kruk et al. have shown by two different methods of measurement of the chemiluminescence intensity and bleaching of p-nitrosodimethylvaniline that even small quantities of dithymoquinone and thymohydroquinone have significant antioxidant properties [15]. Antioxidant activities of these three phenolic compounds were determined by the ABTS radical scavenging method as described in our previous work [16]. The antioxidant power decreased in the order: Thymoquinone > thymol > carvacrol.

The scavenging activities of carvacrol, thymol and thymoquinone were examined by measuring the remaining concentrations of phenolic compounds after 10 min oxidation by HPLC. The concentrations of phenolic compounds were determined in the linear range 50-500 μM for carvacrol and thymol, 4-100 μM for thymoquinone at five concentration levels. Calibration plots with correlation coefficient $R^2 \geq 0.998$ were obtained by reporting peak areas as a function of phenolic compounds concentrations. All the compounds were measured above the quantification level.

The remaining concentrations of carvacrol and thymol were 356.3 ± 0.9 and $341.3 \pm 1.8 \mu\text{M}$ from 400 μM initial concentrations. Thymoquinone remained $22.4 \pm 0.1 \mu\text{M}$ from 40 μM initial concentration. Initial concentration of thymoquinone and Fenton reagent were 10 times lower than other phenolic compounds due to high molar absorptivity coefficient of thymoquinone. Thymoquinone is also an oxidation product of carvacrol and thymol. After 10min oxidation, 6.5 and 10 μM of thymoquinone was produced from carvacrol and thymol, respectively. This indicates that the hydroxyl radicals are previously scavenged by thymoquinone produced during the oxidation reactions of carvacrol and thymol. Then, the rest of the hydroxyl radicals were scavenged by carvacrol and thymol at 400 μM initial concentrations. The decrease in the concentration of phenolic compounds during the Fenton reaction confirms that these compounds are able to scavenge $\text{HO}\cdot$ directly. The scavenging property seems to be more pronounced for thymoquinone, whose ability to scavenge $\text{HO}\cdot$ is higher when compared to that of the carvacrol and thymol.

Stable DPPH \cdot radicals have been used to evaluate the free radical scavenging ability of oregano extracts containing high content of carvacrol, thymol and thymoquinone (Stamenic et al.; Kulisic et al.). The single electron transfer reactions from phenolic compounds to DPPH \cdot has been monitored by recording the decay of the DPPH \cdot ESR signal. The IC₅₀ value is also a parameter widely

used to measure the free radical scavenging activity [17]. A smaller IC₅₀ value corresponds to a higher antioxidant activity. In the case of the investigated phenolic compounds in our study, the free radical scavenging ability was evaluated by measuring the remaining concentrations of phenolic compounds after oxidation reaction by chromatographic technique. The antioxidant activity values of 400 μM carvacrol, thymol and thymoquinone were 0.22, 0.28 and 0.30 mg Trolox/mL, respectively. The antioxidant activity of phenolic compounds is mainly due to their simultaneous hydrogen atom donation to free radicals, electron transfer and metal chelating [18]. The antioxidant activity of the investigated compounds on $\text{HO}\cdot$ is suggested to take place via hydrogen atom donation. The decrease in the concentrations of phenolic compounds corresponds to the scavenging of $\text{HO}\cdot$. Complete elimination of hydroxyl radicals was achieved with the concentration $356.3 \pm 0.9 \mu\text{M}$ of carvacrol, $341.3 \pm 1.8 \mu\text{M}$ of thymol and $22.4 \pm 0.1 \mu\text{M}$ of thymoquinone.

The developed Fenton reaction was applied to *Origanum onites* L. plant material due to its main constituents of carvacrol compound. Some other *Origanum* species such as *Origanum vulgare* L. also contains thymol together with carvacrol [19]. Carvacrol was determined in aromatic water of *Origanum onites* L. plant. Essential oil of *Origanum onites* L. also contains almost mainly carvacrol, but aromatic water was used for Fenton reaction in this study because of its consumption for food purposes. Aromatic water (40mL) was obtained from 40g of dried *Origanum onites* L. plant material using Clevenger apparatus. The amount of carvacrol was found as $8.73 \pm 0.12 \text{mM}$ in aromatic water. The Fenton reaction was applied to aromatic water by diluting 400 μM of carvacrol and the reaction was followed as carvacrol compound. The remaining concentration of carvacrol is $330.0 \pm 1.2 \mu\text{M}$ after 15min reaction period. Carvacrol containing aromatic water oxidized higher than the standard carvacrol as described above. The formation of thymoquinone was also followed in Fenton reaction together with carvacrol oxidation. The concentration of thymoquinone was found as $15.1 \pm 0.1 \mu\text{M}$ after 15min reaction period. Application of Fenton reaction to real sample revealed that carvacrol was more oxidized and thymoquinone formed higher concentration when compared with standard carvacrol.

Conclusion

Antioxidant ability of carvacrol, thymol and thymoquinone was investigated in Fenton reaction at 37°C. The chromatographic methods can provide more detailed information about the reactivity of these phenolic compounds with free radicals. It was demonstrated that the decrease in concentrations of carvacrol, thymol and thymoquinone detected by HPLC shows the scavenging of free radicals generated during the oxidation reaction. Chromatographic results revealed that thymoquinone is an oxidation product of carvacrol and thymol. It has also the highest antioxidant potential when compared with carvacrol and thymol. The present results show that carvacrol, thymol and thymoquinone can serve as effective scavengers of reactive free radicals. Similar results were observed with the application of the method to real sample of aromatic water obtained from *Origanum onites* L.

Acknowledgement

This work was supported by the Uludag University Research Foundation of Turkey (Project No. 2013/13).

References

1. Santos EH, Kamimura JA, Hill LE, Gomes CL. Characterization of carvacrol beta-cyclodextrin inclusion complexes as delivery systems for antibacterial and antioxidant applications. *LWT-Food Sci Technol.* 2015; 60: 583-592.
2. Sobczak M, Kalembe D, Ferenc B, Zylinska L. Limited protective properties of thymol and thyme oil on differentiated PC12 cells with downregulated Mgst1. *J Appl Biomed.* 2014; 12: 234-243.
3. Stamenic M, Vulic J, Djilas S, Masic D, Tadic V, Petrovic S, et al. Free-radical scavenging activity and antibacterial impact of Greek oregano isolates obtained by SFE. *Food Chem.* 2014; 165: 307-315.
4. Lupidi G, Scire A, Camaioni E, Khalife KH, Desanctis G, Tanfani F, et al. Thymoquinone, a potential therapeutic agent of *Nigella sativa*, binds to site o human serum albumin. *Phytomedicine.* 2010; 17: 714-720.
5. Nagi MN, Almakki HA, Sayed-Ahmed MM, Al-Bekairi AM. Thymoquinone supplementation reverses acetaminophen-induced oxidative stress, nitric oxide production and energy decline in mice liver. *Food Chem Toxicol.* 2010; 48: 2361-2365.
6. Martysiak-Zurowska D, Wentka W. A comparison of ABTS and DPPH methods for assessing the total antioxidant capacity of human milk. *ActaSci Pol.* 2012; 11: 83-89.
7. Siquet C, Pavia-Martins F, Lima JL, Reis S, Borges, F. Antioxidant profile of dihydroxy- and trihydroxyphenolic acids-A structure-activity relationship study. *Free Radical Res.* 2006; 40: 433-442.
8. Ince S, Kucukkurt I, Demirel HH, Turkmen R, Zemheri F, Akbe E. The role of thymoquinone as antioxidant protection on oxidative stress induced by imidacloprid in male and female Swiss albino mice. *Toxicol Environ Chem.* 2013; 95: 318-329.
9. Darakhshana S, Poura AB, Colagarc AH, Sisakhtnezhad S. Thymoquinone and its therapeutic potentials. *Pharmacol Res.* 2015; 138-158.
10. Llana-Ruiz-Cabello M, Gutierrez-Praena D, Puerto M, Pichardo S, Jos A, Camean AM. *In vitro* pro-oxidant/antioxidant role of carvacrol, thymol and their mixture in the intestinal Caco-2 cell line. *Toxicol in Vitro.* 2015; 29: 647-656.
11. Polyakov NE, Leshina TV, Konovalova TA, Kispert LD. Carotenoids as scavengers of free radicals in a fenton reaction: antioxidants or pro-oxidants?. *Free Radical Bio Med.* 2001; 31: 398-404.
12. Martins RRL, Neves MGPMS, Silvestre AJD, Silva AMS, Cavaleiro JAS. Oxidation of aromatic monoterpenes with hydrogen peroxide catalysed by Mn III/ porphyrin complexes. *J MolCatal A-Chem.* 1999; 137: 41-47.
13. Santos ICMS, Simoes MMQ, Pereira, MMMS, Martins RRL, Neves MGPMS, et al. Oxidation of monoterpenes with hydrogen peroxide catalysed by Keggin-type tungstoborates. *J MolCatal A-Chem.* 2003; 195: 253-262.
14. Jukic M, Milos M. Catalytic oxidation and antioxidant properties of thyme essential oils (*Thymus vulgaris* L.). *Croat ChemActa.* 2005; 78: 105-110.
15. Kruk I, Michalska T, Lichszteid K, Kladna A, Aboul-Enein HY. The effect of thymol and its derivatives on reactions generating reactive oxygen species. *Chemosphere.* 2000; 41: 1059-1064.
16. Aybastier Ö, Sahin S, Demir C. Response surface optimized ultrasonic-assisted extraction of quercetin and isolation of phenolic compounds from *Hypericum perforatum* L. By column chromatography. *SeparSci Technol.* 2013; 48: 1665-1674.
17. Kulisic T, Radonic, A, Katalinic V, Milos M. Use of different methods for testing antioxidative activity of oregano essential oil. *Food Chem.* 2004; 85: 633-640.
18. Calliste CA, Trouillas P, Allais DP, Simon A, Duroux JL. Free radical scavenging activities measured by electron spin resonance study and B16 cell antiproliferative behaviors of seven plants. *J Agr Food Chem.* 2001; 49: 3321-3327.
19. Cirino ICS, Menezes-Silva SMP, Silva HTD, De Souza EL, Siqueira-Junior JP. The Essential Oil from *Origanum vulgare* L. and Its Individual Constituents Carvacrol and Thymol Enhance the Effect of Tetracycline against *Staphylococcus aureus*. *Chemotherapy.* 2014; 60: 290-293.