

Research Article

Considering Costs of Complications in Bariatric Surgery for the Evaluation of New Technologies based on the Example of Surgical Robotics

Hagen ME^{1*}, Douissard J¹, Cohen G², Jung MK¹, Buchs NC¹ and Torso C¹

¹Division of Digestive and Transplant Surgery, Department of Surgery, University Hospital Geneva, Switzerland

²Medical Controlling, University Hospital Geneva, Switzerland

*Corresponding author: Monika E Hagen, Division of Digestive and Transplant Surgery, Department of Surgery, University Hospital Geneva, 4 Rue Gabrielle-Perret-Gentil, 1211 Geneva, Switzerland

Received: June 08, 2020; Accepted: July 03, 2020;

Published: July 10, 2020

Abstract

Purpose: The purpose of this analysis is to analyze the cost of complications of bariatric surgery to support the economic evaluation of new surgical technologies based on the example of robotic surgery.

Methods: Patients who underwent robotic bariatric surgery from 2014 to 2015 with complete economic data were included. The itemized and treatment costs were derived using the REKOLE method and stratified using the Clavien-Dindo classification.

Results: A total of 195 patients were included. 88.7% of patients underwent primary Roux-en-Y gastric bypass surgery, 6.2% underwent gastric sleeve resection, and 5.1% underwent revisional surgery. 136 patients had no complication, 42 were classified with a Clavien I complication, 6 with a Clavien II complication, and 11 with a Clavien III complication. The mean treatment costs were USD 19,857 for patients without complication, USD 20,575 for patients with a Clavien I complication, USD 29,069 for a Clavien II complication, and USD 52,473 for a Clavien III complication.

Conclusion: Complications are important cost drivers with an incremental correlation to the Clavien-Dindo classification. While minor complications have a clear impact, major complications have an exponential effect on overall costs. As such, a decrease in complications might justify higher procedural costs due to the use of new technologies.

Keywords: Cost; Complications; Robotic Surgery; Gastric Bypass; RYGB; Bariatric Surgery

Introduction

New surgical techniques like minimally invasive surgeries, including robotics provide patient benefits such as shorter hospital stay and reduced surgical trauma when compared to the traditional open approach [1]. However, several studies have shown higher costs for robotic surgery when compared with laparoscopy or open surgery [2-4]. Although new technologies are integral to the advancement of surgical care, they often have – sometimes obstructive - incremental costs that are typically associated with significant upfront investments for the initial purchase as well as procedure-related costs, which is also the case for surgical robotics [5,6].

Since the first published robotic procedures more than two decades ago, the adoption of this technology has increased across surgical specialties with a clear uptake in recent years [7]. However, adoption of robotic surgery still limited to the minority of overall world-wide performed surgical procedures [7]. An important factor that contributes to the limited adoption are the high costs of these systems while the clinical value is still under evaluation as independent high impact factor. As an example, the da Vinci Surgical System – the currently most widely used robotic system - accrues capital costs ranging from USD 0.5M to 2.5M, the annual service costs range from USD 80K to 190K, and the per procedure instruments and accessories'

costs range from USD 700 to 3,500 [7]. These costs are – depending on the structure of the respective healthcare environment – at the burden of either the healthcare provider, the insurer, the patient, or another third party [5]. If these upfront costs cannot be balanced with other factors, they might become prohibitive, particularly in healthcare systems that utilize a flat-fee reimbursement. In addition, high-quality and industry-independent research supporting clinical superiority of robotics over conventional approaches is missing at present. As such, technologies should not only be evaluated for its safety and efficacy, but also its clinical impact and cost efficiency.

Potential options for reducing the cost of surgeries using expensive equipment include savings on other material and the improvement of surgical quality, resulting in fewer complications. This concept has been demonstrated for robotic gastric bypass surgeries [8]. However, detailed research is complex, and it seems intriguing to understand the cost structures of surgical complications to estimate the impact of clinical improvements on the cost of surgery.

To date, there is no available reliable model that estimates the specific costs of complications, which could facilitate modeling the cost-effectiveness of new technologies. This study aims to analyze the costs of complications during bariatric surgery and discuss these costs in light of incremental costs for new technologies using the example

of robotics.

Materials and Methods

Patients

This is a retrospective single center analysis of patients who underwent bariatric surgery between January 2014 and December 2015 at the University Hospital Geneva (Switzerland). Patients with missing economic data were excluded.

Patients were operated with either a fully robotic or hybrid approach using the da Vinci Si or Xi Surgical System (Intuitive Surgical Inc., Sunnyvale, CA, USA). The procedures have been previously described [8-11].

Patients routinely entered the hospital the morning of the procedure and stayed in the intermediate care unit until released from anesthesia to the standard ward. Routine blood testing was performed on day 1 and patients were discharged after successful re-alimentation, pain control with oral medication, and a willingness to return home. If medically necessary, deviations from this path were planned and executed.

Data collection

Clinical data was derived from our prospective database and individual cases' charts were reviewed if information was missing. Patients were stratified using the Clavien-Dindo Classification as per the original publication [12]. In case patients underwent additional imaging or other testing outside the previously described clinical pathway, they were classified as a Clavien-Dindo I case, even if imaging or other testing was negative and not strictly indicated (for example ordered by a resident without approval).

The medical cost data included in this study was obtained using the REKOLE method [13]. The REKOLE method is the Swiss national cost accounting system used in hospitals in Switzerland. The REKOLE system was introduced because the Swiss government opted for comparable and transparent hospital costs. Under the REKOLE method, costs are defined as total direct costs of inpatient care and all hospitals adhere to a minimum standard. The REKOLE system was established at the same time as the change of the reimbursement system from the per-diem payment system to the Diagnose Related Groups (DRG) payment system after January 2012, and it has been used since then for academic cost analyses [5].

Statistical analysis

Statistical analysis was carried with Stata 15.0 software (Stata Corp., College Station, USA). P-value lower than 0.05 was considered statistically significant. Descriptive statistics included means and standard deviation for continuous data, or frequency and percentage for discrete data. Considering more than 2 groups of complications, costs comparisons were carried through one-way analysis of variances (ANOVA) assuming the normal distribution of costs within groups, independence of observations and normal distribution of variances. Tukey's honest significant difference (Tukey HSD) test was chosen as post-hoc test to explore differences between groups and determine their relative statistical significance.

Results

195 patients were included in this analysis. 173 (88.7%) underwent

Table 1: Demographic parameters.

Parameter	Bariatric Surgery Patients 2014–2015 (n=195)
Procedure, n (%)	
Primary gastric bypass	173 (88.7)
Primary gastric sleeve	12 (6.2)
Revisional surgery	10 (5.1)
Age, years	
Mean (Stdev.)	42.9 (11.0)
Gender, n (%)	
Female	145 (74.4)
Male	50 (25.6)
BMI, kg/m ²	
Mean (Stdev.)	43.6 (5.6)
ASA, n (%)	
I and II	126 (64.6)
III and IV	69 (35.4)

primary gastric bypass, 12 (6.2%) underwent gastric sleeve resection, and 10 (5.1%) underwent a revisional procedure. The mean age in this cohort was 42.9 (+/- 11) years and 145 (74.4%) were female. The mean BMI was 43.6 (+/- 5.6) kg/m². 126 (64.6%) were classified as American Society of Anesthesiologists (ASA) 1 or 2 and 69 (35.4%) as ASA 3 or 4. The details regarding demographic parameters can be found in Table 1.

136 patients had no complication, 42 a Clavien I, 6 a Clavien 2 and 11 a Clavien 3. The mean 18 days treatment costs for patients without complications were USD 19,857 (+/- 6,740), USD 20,575 (+/- 5,712) for patients with a Clavien I, USD 29,069 (+/- 9,231) for patients with a Clavien II, and USD 52,473 (+/- 32,185.97) for patients with a Clavien III complication. There was a statistically significant difference between groups as determined by one-way ANOVA ($F(3,191) = 39.132, p < 0.001$). The details regarding costs per complication and incremental costs can be found in Table 2.

The costs were itemized into operating room, anesthesia, intensive care unit, nursing, imaging, medical personnel, hotel, laboratory, material, and drugs. No statistically significant differences were observed for all patient groups for materials. All other categories showed significantly higher costs for patients with a Clavien III complication *versus* patients without complications and patients with a Clavien I complication. Differences for all categories were non-significant between a Clavien I complication and no complications. A significant difference between a Clavien II complication and no complications were only observed for laboratory. The itemized costs are listed in Table 3 and Table 4.

Discussion

To our knowledge, this is the first attempt to systematically evaluate the specific cost of complications in bariatric surgery. While it appears logical that more significant complications generate higher perioperative treatment costs, more detailed insights are essential to allow more precise economic modeling when evaluating the financial impact of postoperative complications. A detailed cost data of complications may also offer insights into the source of expenses

Table 2: Treatment costs stratified by Clavien-Dindo classification*.

USD	Clavien I, n=42 20 575 ± 5 712	Clavien II, n=6 29 069 ± 9 231	Clavien III, n=11 52 473 ± 32 186
No complication, n = 136 19 857 ± 6 740			
Difference (%)	718.0 (3.6)	9212.0 (46.0)	32616.0 (164.0)
p-value	0.976	0.112	<0.001
Clavien I, n = 42 20 575 ± 5 712			
Difference (%)		8494.0 (42.8)	31898.0 (160.6)
p-value		0.195	<0.001
Clavien II, n = 6 29 069 ± 9 231			
Difference (%)			23404.0 (117.9)
p-value			<0.001

*Tukey HSD post-hoc test after one-way ANOVA

Table 3: Itemized costs stratified by Clavien-Dindo classification: Operating Room, Anesthesia, Intensive Care Unit, Nursing, Imaging.

Cost in USD	Operating Room	Anesthesia	Intensive care unit	Nursing	Imaging
No complication	8456	3546	0	3311	40
Mean	2218	689	0	1684	173
Clavien I					
Mean	8361	3474	0	3804	206
Stdev.	2280	766	0	1889	350
Clavien II					
Mean	9 234	3661	0	4 695	892
Stdev.	3 720	419	0	600	209
Clavien III					
Mean	12 251	5 690	3 839	11 960	1 560
Stdev.	2 958	2040	7 638	7 767	1 821
One-way ANOVA	F(3,191)=9.4, p<0.001	F(3,191)=23.7, p<0.001	F(3,191)=16.7, p<0.001	F(3,191)=43.0, p<0.001	F(3,191)=40.0, p<0.001
Tukey HSD post-hoc test	Group 1 vs Group 2: Diff=-95.0000, 95% CI=-1159.3547 to 969.3547, p=0.9956	Group 1 vs Group 2: Diff=-72.0000, 95% CI=-450.3289 to 306.3289, p=0.9605	Group 1 vs Group 2: Diff=0.0000, 95% CI=-799.4922 to 799.4922, p=1.0000	Group 1 vs Group 2: Diff=493.0000, 95% CI=-621.7571 to 1607.7571, p=0.6615	Group 1 vs Group 2: Diff=166.0000, 95% CI=-49.6409 to 381.6409, p=0.1935
	Group 1 vs Group 3: Diff=778.0000, 95% CI=-1737.1829 to 3293.1829, p=0.8536	Group 1 vs Group 3: Diff=115.0000, 95% CI=-779.0313 to 1009.0313, p=0.9872	Group 1 vs Group 3: Diff=0.0000, 95% CI=-1889.2847 to 1889.2847, p=1.0000	Group 1 vs Group 3: Diff=1384.0000, 95% CI=-1250.2892 to 4018.2892, p=0.5251	Group 1 vs Group 3: Diff=852.0000, 95% CI=-342.4178 to 1361.5822, p=0.0001
	Group 1 vs Group 4: Diff=3795.0000, 95% CI=1904.9933 to 5685.0067, p=0.0000	Group 1 vs Group 4: Diff=2144.0000, 95% CI=1472.1899 to 2815.8101, p=0.0000	Group 1 vs Group 4: Diff=3839.0000, 95% CI=2419.3177 to 5258.6823, p=0.0000	Group 1 vs Group 4: Diff=8649.0000, 95% CI=6669.4922 to 10628.5078, p=0.0000	Group 1 vs Group 4: Diff=1520.0000, 95% CI=1137.0800 to 1902.9200, p=0.0000
	Group 2 vs Group 3: Diff=873.0000, 95% CI=-1758.4240 to 3504.4240, p=0.8255	Group 2 vs Group 3: Diff=187.0000, 95% CI=-748.3497 to 1122.3497, p=0.9547	Group 2 vs Group 3: Diff=0.0000, 95% CI=-1976.5994 to 1976.5994, p=1.0000	Group 2 vs Group 3: Diff=891.0000, 95% CI=-1865.0349 to 3647.0349, p=0.8364	Group 2 vs Group 3: Diff=686.0000, 95% CI=152.8670 to 1219.1330, p=0.0056
	Group 2 vs Group 4: Diff=3890.0000, 95% CI=1847.8510 to 5932.1490, p=0.0000	Group 2 vs Group 4: Diff=2216.0000, 95% CI=1490.1104 to 2941.8896, p=0.0000	Group 2 vs Group 4: Diff=3839.0000, 95% CI=2305.0357 to 5372.9643, p=0.0000	Group 2 vs Group 4: Diff=8156.0000, 95% CI=6017.1452 to 10294.8548, p=0.0000	Group 2 vs Group 4: Diff=1354.0000, 95% CI=940.2556 to 1767.7444, p=0.0000
	Group 3 vs Group 4: Diff=3017.0000, 95% CI=-43.0109 to 6077.0109, p=0.0549	Group 3 vs Group 4: Diff=2029.0000, 95% CI=941.3075 to 3116.6925, p=0.0000	Group 3 vs Group 4: Diff=3839.0000, 95% CI=1540.4667 to 6137.5333, p=0.0001	Group 3 vs Group 4: Diff=7265.0000, 95% CI=4060.0825 to 10469.9175, p=0.0000	Group 3 vs Group 4: Diff=668.0000, 95% CI=48.0343 to 1287.9657, p=0.0292

when perioperative complications occur. In addition, if the structure of the cost data is well understood in the clinical context, it may be used as a tool to minimize the financial impact of complications.

Our data show the baseline cost outcomes of patients without complications, which are in line with previous publications from our group [5,8]. Interesting observations can be found when closely evaluating the itemized costs as outlined in Table 3: operating room

and anesthesia costs are similar for patients with and without a Clavien I complication, but this post is slightly higher for patients with a Clavien II and significantly higher for Clavien III complication. It appears logical that a Clavien III complication incurs higher costs for the operating room on average because, as per its definition, this class of complication requires a re-intervention. However, this should not be the case for patients suffering from a Clavien II complication.

Table 4: Itemized costs stratified by Clavien-Dindo classification: Medical Personnel, Hotel, Laboratory, Material, Drugs.

Cost in USD	Medical personnel	Hotel	Laboratory	Material	Drugs
No complication					
Mean	1862	541	606	180	154
Stdev.	679	283	519	193	289
Clavien I					
Mean	1894	597	819	201	204
Stdev.	456	222	614	242	325
Clavien II					
Mean	2114	923	1308	155	106
Stdev.	422	207	774	122	69
Clavien III					
Mean	4292	2002	2599	337	3768
Stdev.	2511	1492	1553	742	6898
One-way ANOVA	F(3,191)=28.8, p<0.001	F(3,191)=40.2, p<0.001	F(3,191)=34.1, p<0.001	F(3,191)=1.3, p=0.281	F(3,191)=17.5, p<0.001
Tukey HSD post-hoc test	Group 1 vs Group 2: Diff=32.0000, 95% CI=-352.1762 to 416.1762, p=0.9964	Group 1 vs Group 2: Diff=56.0000, 95% CI=-140.6837 to 252.6837, p=0.8817	Group 1 vs Group 2: Diff=213.0000, 95% CI=-81.0808 to 507.0808, p=0.2414	Group 1 vs Group 2: Diff=21.0000, 95% CI=-98.3905 to 140.3905, p=0.9684	Group 1 vs Group 2: Diff=50.0000, 95% CI=-683.7970 to 783.7970, p=0.9979
	Group 1 vs Group 3: Diff=252.0000, 95% CI=-655.8492 to 1159.8492, p=0.8893	Group 1 vs Group 3: Diff=382.0000, 95% CI=-82.7844 to 846.7844, p=0.1474	Group 1 vs Group 3: Diff=702.0000, 95% CI=7.0558 to 1396.9442, p=0.0467	Group 1 vs Group 3: Diff=-25.0000, 95% CI=-307.1324 to 257.1324, p=0.9957	Group 1 vs Group 3: Diff=-48.0000, 95% CI=-1782.0399 to 1686.0399, p=1.0000
	Group 1 vs Group 4: Diff=2430.0000, 95% CI=1747.8067 to 3112.1933, p=0.0000	Group 1 vs Group 4: Diff=1461.0000, 95% CI=1111.7429 to 1810.2571, p=0.0000	Group 1 vs Group 4: Diff=1993.0000, 95% CI=1470.7918 to 2515.2082, p=0.0000	Group 1 vs Group 4: Diff=157.0000, 95% CI=-55.0053 to 369.0053, p=0.2235	Group 1 vs Group 4: Diff=3614.0000, 95% CI=2310.9746 to 4917.0254, p=0.0000
	Group 2 vs Group 3: Diff=220.0000, 95% CI=-729.8061 to 1169.8061, p=0.9319	Group 2 vs Group 3: Diff=326.0000, 95% CI=-160.2647 to 812.2647, p=0.3073	Group 2 vs Group 3: Diff=489.0000, 95% CI=-238.0615 to 1216.0615, p=0.3045	Group 2 vs Group 3: Diff=-46.0000, 95% CI=-341.1714 to 249.1714, p=0.9777	Group 2 vs Group 3: Diff=-98.0000, 95% CI=-1912.1799 to 1716.1799, p=0.9990
	Group 2 vs Group 4: Diff=2398.0000, 95% CI=1660.8913 to 3135.1087, p=0.0000	Group 2 vs Group 4: Diff=1405.0000, 95% CI=1027.6283 to 1782.3717, p=0.0000	Group 2 vs Group 4: Diff=1780.0000, 95% CI=1215.7549 to 2344.2451, p=0.0000	Group 2 vs Group 4: Diff=136.0000, 95% CI=-93.0714 to 365.0714, p=0.4166	Group 2 vs Group 4: Diff=3564.0000, 95% CI=2156.0833 to 4971.9167, p=0.0000
Group 3 vs Group 4: Diff=2178.0000, 95% CI=1073.4965 to 3282.5035, p=0.0000	Group 3 vs Group 4: Diff=1079.0000, 95% CI=513.5361 to 1644.4639, p=0.0000	Group 3 vs Group 4: Diff=1291.0000, 95% CI=445.5200 to 2136.4800, p=0.0006	Group 3 vs Group 4: Diff=182.0000, 95% CI=-161.2468 to 525.2468, p=0.5172	Group 3 vs Group 4: Diff=3662.0000, 95% CI=1552.3399 to 5771.6601, p=0.0001	

While not statistically significant – also due to the relatively small cohort of patients with a Clavien II complication - we found higher OR costs for patients with a Clavien II complication when compared to patients with or without a Clavien I complication. In this cohort, the cost increases most likely stem from the initial surgery as no re-intervention was conducted. This can be caused either by a prolonged procedure or by higher usage of instruments during the initial surgery. It seems reasonable to assume that a deviation from the standard surgical path (standard length case and/or standard set of instruments) might put the patients at a higher risk of perioperative complications. Still, this dataset does not allow the exact identification of cost drivers in each itemized segment and this comparison is not adequately powered. As such, there might also be a bias since longer procedures can be more complex in patients with high BMIs and other conditions that may lead to longer OR times or higher risk of postoperative complications. It would be interesting to explore further if cost-driving events in the OR can be used as predictors for perioperative complications and as such, if warning systems could be developed depending on the course of the surgical procedures. However, this dataset is not sufficiently deep for such analyses.

A closer look at the itemized costs of the patients with a Clavien I complication show comparable costs to the cohort without complications. However, we found an increase - although not statistically significant - in imaging and laboratory costs. This observation seems reasonable in the clinical context as a Clavien I complication is a deviation from the standard clinical path not requiring specific treatment. As a matter of fact, most of these patients did not have a real medical problem, but received additional imaging or laboratory work prescribed by an inexperienced resident. As a teaching institution, residents are primarily responsible for the post-operative patients at night and over weekends and have the freedom to order these tests. Under these circumstances, we often observe a rather generous use sometimes without clear medical indication. However, as per definition these patients need to be classified as a Clavien I and it easily explains increased costs for imaging and laboratory. Another logical observation is the absence of intensive care unit costs for patients with or without a Clavien I and Clavien II complication, but the presence of intensive care unit costs when a re-intervention is needed. All the above mentioned clinically relevant consistencies suggest that this set of cost data is robust and can be

used as a solid baseline for economic modeling.

Overall, Clavien I complications within the 18 days of robotic bariatric surgery seem to incur a moderate cost increase of about 3.6%, but the further increase of about 36% and 164% for Clavien II and III complications are rather exponential. It appears evident that any complication requiring treatment is not only devastating for the patient, but also an economic challenge to the healthcare system. As such, there is room for investments to avoid complications and from an economic perspective. As this is a first dataset quantifying the economic impact of complications in bariatric surgery, it can be used as a guide to show the extent of upfront investments that can be afforded to improve outcomes. While these investments might include a wide range of measures including surgeon training, patient preparation, and others, we want to focus on the investment in new surgical technologies. Albeit the absence of high quality and high level of evidence supporting robotics for a widespread application in bariatric surgery, this technology finds an increasing adoption, also driven by the desire to reduce perioperative complications of bariatric surgery. As an example, a few groups have published the positive impact of surgical robotics for gastric bypass surgery regarding procedure times per BMI and a reduction in severe complications such as intestinal leaks [8,14-16]. One study showed a cost reduction with a robotic approach *versus* laparoscopy for gastric bypass surgery due to a reduction in complications and the use of fewer surgical staplers [8]. This dataset shows that a Clavien III complication costs about USD 33,000. Therefore, a 1% reduction in a postoperative re-intervention rate would save USD 33,000 in 100 patients, and this money can be spent on new surgical technologies such as robotics.

Currently, the da Vinci Surgical System is the most widely used robot and its published costs have been listed above [7]. While these costs may fluctuate depending on the exact surgical approach and the instruments required, this post has been calculated to be USD 1,582.91 in a previous academic publication for gastric bypass surgery in 2011 [8]. Applying this cost structure, it becomes obvious, that a 1% reduction in re-interventions (saving about USD 33,000) would not be sufficient to counterbalance this upfront investment, but a significantly greater reduction of Clavien III complications would be required to financially justify the use of the da Vinci Surgical System. While this is a very simplified model, it outlines the practical applicability of this dataset and the extent of financial considerations that have to be made before widespread clinical adoption.

Still, several shortcomings of this analysis have to be mentioned: first, this dataset is mostly based on economic data and many conclusions require more data breadth and depth. In addition, this analysis includes a limited number of subjects in the healthcare environment of Switzerland. As such, the data might not be applicable in other geographic regions or might not be the same when looking at larger cohorts. Also, this is a retrospective analysis with all commonly known sources of bias. Lastly, the cost of robotic surgery is currently under development: Intuitive Surgical as well as new competitors address the cost issue with alternative models including flat-fee payments and lease options [17,18]. Still, this is a unique approach for analyzing the bariatric cohort and might be a decent source of reflection for both clinicians and economists.

Conclusion

Complications are important cost drivers and correlate to the Clavien-Dindo classification with an exponential increase for more significant complications. While this is a good rationale to justify higher procedural costs if complications can be avoided, major investments including current surgical robotics might not be offset by a clinically realistic reduction of complications. More systematic research, including the analyses of broader and deeper datasets, is needed to generate more detailed insights into the correlation of perioperative complications and costs.

Compliance with Ethical Standards

Conflict of interest

Dr. Hagen reports grants from Intuitive Surgical Inc., personal fees and non-financial support from Quantgene Inc., personal fees and non-financial support from Verb Surgical, personal fees and non-financial support from Johnson & Johnson, outside the submitted work.

Dr. Jung reports personal fees and non-financial support from Intuitive Surgical Inc., personal fees and non-financial support from Ethicon Endosurgery, outside the submitted work.

Dr. Douissard reports grants from Intuitive Surgical Inc., personal fees and non-financial support from Verb Surgical, outside the submitted work.

Dr. Cohen has nothing to disclose.

Dr. Buchs has nothing to disclose.

Dr. Toso reports grants from Intuitive Surgical Inc., personal fees and non-financial support from Ethicon Endosurgery Inc., outside the submitted work.

Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Acknowledgement

We thank Françoise Bernardi (University Hospital Geneva) for data collection.

References

1. Fuchs KH. Minimally invasive surgery. *Endoscopy*. 2002; 34: 154-159.
2. Abdelmoaty WF, Dunst CM, Neighorn C, Swanstrom LL, Hammill CW. Robotic-assisted *versus* laparoscopic unilateral inguinal hernia repair: a comprehensive cost analysis. *Surg Endosc*. 2018.
3. Hagen ME, Balaphas A, Podetta M, Rohner P, Jung MK, Buchs NC, et al. Robotic single-site *versus* multiport laparoscopic cholecystectomy: a case-matched analysis of short- and long-term costs. *Surg Endosc*. 2018; 32: 1550-1555.
4. Trehan A, Dunn. TJ. The robotic surgery monopoly is a poor deal. *BMJ*. 2013; 347: f7470.
5. Hagen ME, Rohner P, Jung MK, Amirghasemi N, Buchs NC, Fakhro J, et al. Robotic Gastric Bypass Surgery in the Swiss Health Care System: Analysis of Hospital Costs and Reimbursement. *Obes Surg*. 2017; 27: 2099-2105.

6. Niklas C, Saar M, Berg B, Steiner K, Janssen M, Seiemer S, et al. da Vinci and Open Radical Prostatectomy: Comparison of Clinical Outcomes and Analysis of Insurance Costs. *Urol Int*. 2016; 96: 287-294.
7. Inc., IS. Investor Presentation Q4 2019. Accessed. 11-19-2019.
8. Hagen ME, Pugin F, Chassot G, Huber O. Reducing cost of surgery by avoiding complications: the model of robotic Roux-en-Y gastric bypass. *Obes Surg*. 2012; 22: 52-61.
9. Buchs NC, Hagen ME, Pugin F, Volonte F, Bucher P, Schiffer E, et al. Intra-operative fluorescent cholangiography using indocyanin green during robotic single site cholecystectomy. *Int J Med Robot*. 2012; 8: 436-440.
10. Buchs NC, Pugin F, Chassot G, Volonte F, Koutny-Fong P, Hagen ME, et al. Robot-assisted Roux-en-Y gastric bypass for super obese patients: a comparative study. *Obes Surg*. 2013; 23: 353-357.
11. Hagen ME, Jung MK, Fakhro J, Buchs NC, Buehler L, Mendoza JM, et al. Robotic versus laparoscopic stapling during robotic Roux-en-Y gastric bypass surgery: a case-matched analysis of costs and clinical outcomes. *Surg Endosc*. 2018; 32: 472-477.
12. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004; 240: 205-213.
13. Besson P. REKOLE® - Betriebliches Rechnungswesen im Spital, (Comptabilité analytique à l'hôpital) 2013: Die Spitäler der Schweiz H+, Bern.
14. Sanchez BR, Mohr CJ, Morton JM, Safadi BY, Alami RS, Curet MJ. Comparison of totally robotic laparoscopic Roux-en-Y gastric bypass and traditional laparoscopic Roux-en-Y gastric bypass. *Surg Obes Relat Dis*. 2005; 1: 549-554.
15. Mohr CJ, Nadzam GS, Alami RS, Sanchez BR, Curet MJ. Totally robotic laparoscopic Roux-en-Y Gastric bypass: results from 75 patients. *Obes Surg*. 2006; 16: 690-696.
16. Tieu K, Allison N, Snyder B, Wilson T, et al. Robotic-assisted Roux-en-Y gastric bypass: update from 2 high-volume centers. *Surg Obes Relat Dis*. 2013; 9: 284-288.
17. Guthard G, Morgan JP. Health care Conference. 2020; 2020.
18. CMR. Accessed 02-16-2020.