

## Clinical Image

# Neurogenic Paraosteoarthropathy Following Severe Head Injury

**Akhouayri M\***; Benabbouha ADepartment of Orthopedic Surgery and Traumatology,  
University Cadi Ayyad, Morocco**\*Corresponding author: Akhouayri M**Department of Orthopaedic Surgery and Traumatology,  
Avicenne Military Hospital, Faculty of Medicine and  
Pharmacy, University Cadi Ayyad, 40000 Marrakech,  
Morocco.

Tel: + 212 659-660-561

Email: makhouayri@gmail.com

**Received:** July 10, 2024**Accepted:** August 26, 2024**Published:** September 03, 2024

## Clinical Image

Standard radiograph and CT scan (Figure 1) of the pelvis of a 40-year-old man with disabling bilateral hip stiffness making ambulation impossible. The patient had suffered a severe cerebral trauma one year earlier, resulting in a 45-day stay in the intensive care unit. These images revealed heterotopic peri-articular ossifications with bone bridges between the greater trochanter and the acetabulum. Neurogenic para-osteoarthropathy remains a significant and disabling complication occurring in 5-20% of severe head injuries. The pathophysiology involves complex interactions between neural and inflammatory pathways, leading to ectopic bone formation [1]. This case underscores the importance of early diagnosis and intervention to manage symptoms and improve patient outcomes. This article aims to contribute to the relatively limited literature on this condition.

**Keywords:** Neurogenic paraosteoarthropathy

## Author Statements

## Disclosure of Interest

None of the authors has any conflicts of interest to declare.

**Figure 1:** Imaging of neurogenic paraosteoarthropathy of the hip.

## References

1. Sullivan MP, Torres SJ, Mehta S, Ahn J. Heterotopic ossification after central nervous system trauma: A current review. *Bone & Joint Research*. 2013; 2: 51-7.