

Review Article

Definition and Self-Reported Pain Intensity in Adolescents with Dysmenorrhea: A Debate Report

De Sanctis V^{1*}, Ashraf S², Sergio B³, Luigi B⁴, Gianni B⁵, Mauro B⁶, Fabio B⁷, De Sanctis C⁸, Giorgio T⁹, Franco R¹⁰ and Perissinotto E¹¹

¹Private Accredited Hospital Quisisana, Pediatric and Adolescent Outpatients Clinic, Italy

²Department of Pediatrics, Division of Endocrinology, Alexandria University Children's Hospital, Egypt

³University "G. D'Annunzio" Chieti-Pescara, Italy

⁴Rehabilitation Centre, La Nostra Famiglia, Italy

⁵Division of Pediatrics, Department of Mother and Child Health, Azienda Ospedaliero-Universitaria Maggiore della Carità, Italy

⁶Internal Medicine and Therapeutics, Section of Childhood and Adolescence, University of Pavia, Foundation IRCCS San Matteo, Italy

⁷Department of Pediatrics, "Carlo Poma" Hospital, Italy

⁸Department of Pediatric Endocrinology, Ospedale Infantile Regina Margherita, Italy

⁹Centre of Pediatric Diabetology, Burlo Garofolo Hospital, Italy

¹⁰Department of Pediatrics, University of Padua, Italy

¹¹Department of Cardiac, Thoracic and Vascular Sciences, Unit of Biostatistics, Epidemiology and Public Health, University of Padua, Italy

*Corresponding author: De Sanctis V, Pediatric and Adolescent Outpatient Clinic, Quisisana Hospital, Italy

Received: April 20, 2016; Accepted: June 25, 2016;

Published: June 30, 2016

Abstract

Background: Dysmenorrhea is one of the most common complaints among adolescents and women. Patho-physiologically, it is categorized into primary and secondary dysmenorrhea. Primary Dysmenorrhea (PD) refers to pain with no obvious pathological pelvic disease; Secondary Dysmenorrhea (SD) is caused by an underlying pelvic condition or pathology. The prevalence of primary dysmenorrhea varies between 67% and 90% in adolescents, with severe pain perceived in 7%-15% of the women studied.

The lowest prevalence of 16% was reported in a random sample of Japanese women aged 17-51 years through daily diary recording for 1 month. The highest prevalence of 91% was reported in a random sample of Iranian women aged 16-56 years, with the majority younger than 30 years of age without children. The prevalence of primary dysmenorrhoea is difficult to determine because many affected women do not seek medical treatment. The majority accept pain as a part of their normal menstrual cycle.

Aim of the study: The authors reviewed the epidemiology data reported in the literature on 4 aspects of Dysmenorrhea: definition, pain severity, prevalence and development of chronic pain.

Results: The authors highlight the necessity for finding a uniform definition of dysmenorrhea. The wide variation in the prevalence rates may be due to using different definitions, employing selected groups of subjects and/or the absence of a universally established method for measuring pain. Variation in the methods that assess the severity of dysmenorrhea (ranging from occasional menstrual cramps to severe pain that interferes with daily activities and/or to require medication) adds to the variation in the prevalence among different studies.

Definition: Dysmenorrhea: Pain severity, prevalence and development of chronic pain.

Conclusions: Studying the epidemiology of menstrual pain is an important issue that deserves further attention because of its high prevalence and negative effect on women's health.

Keywords: Dysmenorrhea; Adolescents; Epidemiology; Health Problems

Background

Dysmenorrhea is one of the most common complaints among adolescents and women of reproductive age. It is categorized into two types: primary and secondary. Primary Dysmenorrhea (PD) refers to pain with no obvious pathological pelvic disease. Secondary Dysmenorrhea (SD) is caused by an underlying pelvic pathology and occurs more in women older than 20 years. SD can be caused by endometriosis, pelvic inflammatory disease, intrauterine devices, ovarian cysts, adenomyosis, uterine myomas or polyps, intrauterine adhesions or cervical stenosis [1-5].

PD is characterized by a crampy suprapubic pain that begins between several hours before and a few hours after the onset of the menstrual bleeding. Symptoms peak with maximum blood flow and usually last less than one day, but the pain may persist up to 2-3 days. Symptoms are more or less reproducible, from one menstrual period

to the other [3,4]. The pain is characteristically colicky and located in the midline of the lower abdomen, but may be dull and may extend to both lower quadrants. Pain can radiate to the back of the legs or the lower back. Associated nausea, vomiting, diarrhea, fatigue, mild fever and headache or lightheadedness are fairly common.

Dysmenorrhoea may impair the quality of (personal and social) life. In many women it is associated with mood disorders, sleep disturbance and limitations in performance of daily activities (school and work).

Dysmenorrhea shortly after menarche or in a patient who is clearly anovulatory should alert the physician to the possibility of an obstructing malformation of the genital tract. However, adolescents may experience menstrual pain with their first periods (anovulatory) without any demonstrable underlying cause, when the bleeding is heavy and accompanied in background, by clots [2,4]. Menstrual pain

Table 1: Some Criteria used in Literature to define Dysmenorrhea.

Authors and References	Criteria used to define Dysmenorrhea
Dawood MY. [33]	Painful menstrual cramps which appear to have no macroscopically identifiable pelvic pathology
Tangchai K. [56]	Any type of pain or discomfort associated with menstrual period
Lefebvre G, et al. [57]	A difficult menstrual flow
Proctor M, et al. [58]	A painful menstrual cramp of uterine origin
Rostami M. [59]	A painful syndrome which occurs at the time of menstrual flow in ovulatory cycles
Begum J, et al. [60]	Any type of pain or discomfort associated with menstrual period
Suresh K, et al. [61]	A pelvic pain associated with onset of menses and lasting for 8-72 hours
Grandi G. [62]	A cramp-like, dull, throbbing pain that usually emanates from the lower abdomen, and that occurs just before and/or during menstruation
Shah M. [63]	Menstrual pain without organic pathology

appearing after several years of painless periods is suggestive of SD [2-5].

Pain intensity in PD can be mild (pain that does not disturb daily activities or require painkillers), moderate (pain that slightly interferes with daily routines, but can be managed with painkillers), and severe (pain that entirely prevents daily life activities) [6,7].

Pathophysiology of PD

At the end of the luteal phase in non-pregnant women, the corpus luteum regresses, with a consequent decline in the progesterone level. This decline favors the production of prostaglandin precursor, arachidonic acid. Arachidonic acid, then enters the cyclooxygenase pathway that leads to the production of prostaglandins. Consequently, there is increased production and release of prostaglandins causing Prostaglandins stimulates contraction of vascular and uterine smooth muscle, causing an excessive uterine contractions and constriction of endometrial blood vessels. The vasoconstriction causes ischemia of the endometrium and expelling the menstrual effluent from the uterine cavity. Both the ischemia and the myometrial contraction explain the cramping of ovulatory menstrual cycles [8-14]. Increased leukotrienes and vasopressin and diminution of prostacyclin levels are contributing factors. Leukotrienes amplify myometrial contraction and vasoconstriction. Women who fail to respond to prostaglandin inhibitors may have elevated levels of leukotrienes [15]. Leukotrienes can increase the sensitivity of pain fibres [15,19]. Increased vasopressin levels, without an accompanying increase in oxytocin levels, can produce dysrhythmic uterine contractions that are more likely to produce uterine hypoxia and ischemia [17-19]. Stimulation of pain fibres in the uterus causes activation of the afferent pain pathways transmitted up to the central nervous system. It has also been suggested that women with PD have increased expression of pro-inflammatory cytokines and decreased expression of growth factors in the secretory and regenerative phases of MP. These factors may be involved in the regulation of endometrium breakdown and repair and indirectly exacerbate pain [16].

In summary, little is known about the various patho-physiological, vascular, molecular and neural mechanisms that produce and control the pain of dysmenorrhea.

Prevalence and Risk Factors for PD

Dysmenorrhea is a common symptom for a large proportion of young women. Severe pain, limiting daily activities is less common.

The reported prevalence of dysmenorrhea varies substantially. A greater prevalence is generally observed in young women (17-24 years), with estimates ranging from 67% to 90% [20,21]. In Australia, the prevalence of PD in senior high school girls is high (93%), while severe pain, sufficient to limit daily activities, is less common (7%-15%) [22,20].

Studies suggest that the importance the interplay of four main categories of factors in the prevalence of dysmenorrhea. These are: menstrual history, lifestyle characteristics, works related factors and personal variation of pain perception [23-25]. The prevalence of primary Dysmenorrhoea is difficult to determine because many affected women do not seek medical treatment. The majority accept pain as a part of their normal menstrual cycle.

Aim of the Study

The authors reviewed the epidemiology data reported in the literature on 4 aspects of dysmenorrhea: definition, pain severity, prevalence and development of chronic pain.

Results

Definition

The definition of dysmenorrhea varied considerably between different studies (Table 1). In 3 large studies in the United States, variable selection of women with dysmenorrhea, based on different definitions, produced big variation in the prevalence. One of the studies considered only those with moderate to extreme uterine cramping pain [27], the second recruited participants with one or more episode of severe pain [28] and the third selected participants with some degree of dysmenorrheal [29]. Other studies defined menstrual pain as dysmenorrheal [30-32]. Some considered dysmenorrhea to be any menstrual pain associated with "the need for medication or inability to function normally" [33]. A more recent study denied any association between intensity of menstrual pain and need for medication or work intolerance [34]. These definitions are clearly quite different and result in great variability in prevalence of dysmenorrhea.

Pain severity

Two instruments are commonly used to measure the subjects' severity of dysmenorrheal are Verbal Multidimensional Scoring System (VMS) and the Visual Analogue Scale (VAS) [35]. The VMS measures pain severity and takes into account the impacts of pain on daily. Activities, systemic symptoms and analgesic requirements.

Table 2: Summary of Studies on Dysmenorrhea in different Countries from 2011 to 2015.

References	Study Characteristics and Number of Subjects	Prevalence and Severity (%) of Dysmenorrhea
Khamdan HY, et al.[64]	Cross sectional study. 226 students. Mean Age: 21 years.	90.7%
Haque SE, et al. [65]	Intervention study. 416 Students Aged: 11–16 yrs.	Severe: 59.8%
Pitangui AC, et al.[66]	Public school 218 Adolescents Aged:12-17 yrs.	73%
Nooh AM. [67]	283 questionnaires. Mean Age: 17.8±0.7 yrs.	65.4%; 27.9%: Mild, 23.3%: Moderate and 14.1%: Severe.
Zegegne TK, et al.[68]	A mixed-method research combining quantitative and qualitative methods 562 students. Age range: 9-12 years, mean age of 14.96 yrs.	Severe: 20.9%
Gagua T, et al. [69]	Cross-sectional study including 2,561 women. Mean age 16±1.3 yrs.	52%
Gumanga SK, et al. [70]	Cross-sectional descriptive study. 456 girls Age Range: From 14-19 yrs; Mean age 16±0.93 yrs.74.4% Mild:18.1%; Moderate: 37.5%.	Severe: 18.8%.
Chia CF, et al. [71]	Cross-sectional questionnaire. 240 students. (128 medical and 112 non medical). Mean Age: 20.1±1.4 yrs.	80%
Rani A, et al. [72]	Cross-sectional study. 300 girls Age Range: 11-18 yrs	61.3%
Al Asadi JN, et al. [73]	Cross sectional study 384 students, mean age 17.1 yrs.	89.4% Moderate: 54.3%; Severe: 43.9%
Baghianimoghadam MH. [74]	Cross-sectional study. 300 students of university. Age Range of Participants: 18-35 years (Mean: 21±4.3 yrs).	38.1%
De Sanctis V, et al. [75]	Cross-sectional multicenter study. 4,992 girls (Mean Age: 17.1 yrs).	56% In 56% subject it was severe
Nohara M, et al. [76]	Anonymous self-administered questionnaire 2,166 women. 10.4% <25 yr.	77.6% 2.8% of female workers responded that they have "Very Serious Pain", 25.8% "Serious Pain", and 49.7% responded that "The Pain Could be Tolerated".
Bata MS. [77]	Self-administered questionnaire. 596 secondary-school students. Mean age:15.7±1.5 yrs.	Mid-cycle pain was reported by 30.9% and dysmenorrhea by 37.6% of the students
Al-Jefout M, et al. [78]	Cross-sectional study. 272 female medical students (Aged 19-25 years).	In 55.8% from moderate to severe
Lee JC, et al. [79]	Anonymous questionnaire 538 teenage girls, aged 14 to 18 yrs (Mean Age: 16.1 yrs).	82% Severe in 0.9% of subjects
Wong LP. [80]	Cross-sectional study. 1,295 adolescent girls (Aged 13-19 yrs) of a rural school	76.0%
Al-Kindi R, Al-Bulushi A. [81]	Cross-sectional survey. 404 girls from two public high schools. Age range:15-23 yrs.	94% Mild : 27% ; Moderate: 41% ; Severe: 32%
Gulzar, et al. [82]	Cross-sectional descriptive study. 337 school and college going female students aged between 12-19 years.	78%.
Rodrigues AC, et al. [83]	Observational transversal study. 274 adolescents and young adults (Age ≤26)	62.8%
Bano R, et al. [84]	Cross-sectional descriptive study using self-administered questionnaire. 100 participants. Age Range:18-24 yrs.	Mild: 20%; Moderate: 43% and severe in 37%.
Wijesiri HS, et al. [85]	A descriptive study. 200 adolescent students.	84%
Narring F, et al. [86]	Cross sectional survey. 3,340 females, aged 16 to 20 yrs who attended post-mandatory education	86.6% In 12.4%: Severe and in 74.2% moderate dysmenorrhea.
Pembe AB, Ndolele NT. [87]	Cross sectional study. 880 girls of eight secondary schools.	74.1%
Sahin S, et al. [88]	The study group included 520 students. Mean age of the students was 20.2±1.5 years (Range 17-25yrs).	69% Frequency of dysmenorrhea was higher in smokers, those with menstrual irregularity, those who used drugs for menstrual regulation and having a family history.
Jang IA, et al. [89]	Questionnaires. 3,017 Vietnamese women aged 17 to 42yrs (Mean: 25.5 yrs).	58.8% Mild to moderate: 7.8% Moderate: 65.9% Moderate to severe: 23.3% Severe: 2%

Based on this system, pain levels are defined based according to the criteria listed below:

Mild Dysmenorrhea: Painful menses that do not limit or hinder normal daily activities, and results in little or no systemic symptoms and/or analgesic need.

Moderate Dysmenorrhea: Painful menses that slightly limit or hinders normal daily activities, and result in moderate systemic symptoms and/or analgesic need.

Severe Dysmenorrhea: Painful menses that severely limit or hinder normal daily activities result in noticeable symptoms (such as fainting, vomiting) and respond poorly to analgesics.

The VAS method entails using a 100mm horizontal line scale that begins with no pain on the far left to the worst possible pain on the far right. Participants will be asked to indicate a perception of pain intensity scored from one to 100 by drawing a mark on this scale representing their subjective pain.

Pain is an extremely subjective symptom very difficult to quantify [36]. Unfortunately, several prevalence studies did not use any standardized scale for evaluation of menstrual pain severity. They used unreliable (not validated) tools for assessing severity of dysmenorrheal [36]. In addition, many symptoms related to menstruation are not well-defined, and thus may increase the misclassification for symptoms as 'menstrual' when they are not [37]. Moreover, the great personal difference in pain perception and reaction to this pain by different women can markedly affect the judgement of pain. Adolescents may not recall the time and severity of pain because they are still learning about their menstrual cycle.

Sixteen studies examined the associations between the perceptions of experimentally induced pain across menstrual cycle phases of healthy females. In healthy Hong Kong Chinese women the pain level was higher compared to Westerns women. Sex hormones may contribute to inter- and intra-individual differences in perception and reaction to pain. Pain inhibition is more effective in the ovulatory phase of the menstrual cycle than in the early follicular phase [38-40]. Therefore, research in this area has flaws in methodology.

In summary, menstrual pain should be regarded as a multidimensional phenomenon and should be measured by a multidimensional scoring system. Future research needs to specify a comprehensive list of dysmenorrhea-specific symptoms based on the pathogenesis of PD, explore relevant symptom dimensions (severity, frequency, duration, and timing) and test validity of the adapted methodology [41].

Prevalence of PD

The prevalence of PD is difficult to determine, because few women seek medical treatment. Many women accept menstrual pain as "normal" part of the menstrual cycle [42]. Thus, many cases remain undocumented [43-46].

Previous studies were retrospective, using different definitions of dysmenorrhea and different periods of recall of symptoms (weeks, months, and year) [43,47]. In addition, there was a lack of standard (validated) method for assessing its severity. Therefore, prevalence estimates vary widely (67% to 90%) [20,21,46]. The lowest prevalence

(16%) was reported by daily diary recording in Japanese women (17-51years). The highest prevalence (91%) was reported in Iranian women (16-56years) with the majority younger women than 30years, with no children [25]. Other studies focused on selected groups of subjects instead of taking a representative sample from the population of women in the community [48,49].

The reported downward trend in the age of menarche in the developed countries over the past century may change the experience of menstruation and related symptoms and prevalence reported in older studies. A summary of studies on dysmenorrhea in different countries from 2011 to 2015 is reported in (Table 2).

Development of chronic pain

There is a scarcity of longitudinal studies on the natural history of dysmenorrhea and the effects of a range of possible modifiable risk factors over time. In addition, the duration of follow-up in these studies was relatively short. Several aspects, therefore, are still open.

A longitudinal study of young women between the ages of 19 and 24 conducted by Sundell et al. showed that the severity of dysmenorrhea only decreased in women who gave birth to children during those 5 years, but was unchanged in those who remained nulliparous, had a miscarriage or an abortion [50]. Another study of Weissman et al. found a significant effect of age on the severity of dysmenorrhea. After adjusting for parity, older women were more likely to experience a decrease in the severity of dysmenorrhea, independent of childbearing [51]. Vincent et al. made an important observation, that the longer is the duration of reported dysmenorrheic symptoms (from 2-28 years), the greater is the suppression of the woman's hypothalamic-pituitary-adrenal axis, as manifested by a reduction in cortisol levels [52].

Conclusion

Dysmenorrhea represents a substantial public health burden. It has a high prevalence worldwide and is one of the leading causes of absenteeism from school and work, loss of earnings and diminished quality of life. Therefore, studying the epidemiology of menstrual pain deserves further attention [1-5].

Various prevalence rates are attributed to the use of selected groups of subjects, absence of a universally accepted method for defining dysmenorrhea or its severity, and reluctance of many women to report pain and/or seek medical treatment. In addition, the majority of studies did not distinguish between the types of dysmenorrhea (primary versus secondary). Teenagers, unlike adults, may not be experienced enough to report menstrual pain because they are still learning about the character of their cycles. Therefore, studies that included a good number of girls who just started their menarche have limited value as these symptoms often begin after regular ovulation (2 to 3 years post menarche) [37]. Adolescent receive information regarding menstruation mainly from their mothers [53,54]. Only 2% receive information about menstruation from health care providers [34]. Therefore, it is recommended that health care providers should be knowledgeable and well trained to educate and provide proper care for adolescents with dysmenorrheal [53-55].

In conclusion, this report highlights 4 main issues that deserve further consideration and research. High quality large population-

based longitudinal studies are still needed to delineate the natural history of dysmenorrhea and its potential effect on the reproductive life. A more uniform definition of dysmenorrhea and accurate grading system for severity, agreed upon by experts, are required. Age-specific incidence and prevalence studies of dysmenorrhea in different populations are still needed. There is also a high demand for studying the effect/s of possible risk factors on menstrual pain using appropriate statistical methods.

References

- Rosenwaks Z, Seegar-Jones G. Menstrual pain: its origin and pathogenesis. *J Reprod Med.* 1980; 25: 207-212.
- Coco AS. Primary dysmenorrhea. *Am Fam Physician.* 1999; 60: 489-496.
- Iglesias EA, Coupey SM. Menstrual cycle abnormalities: Diagnosis and management. *Adolesc Med.* 1999; 10: 255-273.
- Schroeder B, Sanfilippo JS. Dysmenorrhea and pelvic pain in adolescents. *Pediatr Clin North Am.* 1999; 46: 555-571.
- Durain D. Primary dysmenorrhea: Assessment and management update. *J Midwifery Womens Health.* 2004; 49: 520-528.
- Turk DC and Melzack R. *Handbook of Pain Assessment.* 2011.
- Dawood MY. Primary dysmenorrhea: advances in pathogenesis and management. *Obstet Gynecol.* 2006; 108: 428-441.
- Rosenwaks Z, Seegar-Jones G. Menstrual pain: Its origin and pathogenesis. *J Reprod Med.* 1980; 25: 207-212.
- Bieglmayer C, Hofer G, Kainz C, Reinthaller A, Kopp B, Janisch H. Concentrations of various arachidonic acid metabolites in menstrual fluid are associated with menstrual pain and are influenced by hormonal contraceptives. *Gynecol Endocrinol.* 1995; 9: 307-312.
- Akerlund M. Contractility in the nonpregnant uterus. *Ann N Y Acad Sci.* 1997; 828: 213-222.
- Owen PR. Prostaglandin synthetase inhibitors in the treatment of primary dysmenorrhea. Outcome trials reviewed. *Am J Obstet Gynecol.* 1984; 148: 96-103.
- Naves LA, McCleskey EW. An acid-sensing ion channel that detects ischemic pain. *Braz J Med Biol Res.* 2005; 38: 1561-1569.
- Maia H Jr, Haddad C, Coelho G, Casoy J. Role of inflammation and aromatase expression in the eutopic endometrium and its relationship with the development of endometriosis. *Womens Health (Lond Engl).* 2012; 8: 647-658.
- Bieglmayer C, Hofer G, Kainz C, Reinthaller A, Kopp B, Janisch H. Concentrations of various arachidonic acid metabolites in menstrual fluid are associated with menstrual pain and are influenced by hormonal contraceptives. *Gynecol Endocrinol.* 1995; 9: 307-312.
- Demers LM, Hahn DW and McGuire JL. Newer concepts in dysmenorrhea research: leukotrienes and calcium channel blockers. In: Dawood MY, McGuire JL, Demers LM, editors. *Premenstrual Syndrome and Dysmenorrhea.* Baltimore: Urban & Schwarzenberg. 1985: 205.
- Ma H, Hong M, Duan J, Liu P, Fan X, Shang E, Su S. Altered cytokine gene expression in peripheral blood monocytes across the menstrual cycle in primary dysmenorrhea: A case-control study. *PLoS One.* 2013; 8: 55200.
- Akerlund M, Strömberg P, Forsling ML. Primary dysmenorrhoea and vasopressin. *Br J Obstet Gynaecol.* 1979; 86: 484-487.
- Melin P, Trojnar J, Johansson B, Vilhardt H, Akerlund M. Synthetic antagonists of the myometrial response to vasopressin and oxytocin. *J Endocrinol.* 1986; 111: 125-131.
- Guo SW, Mao X, Ma Q, Liu X. Dysmenorrhea and its severity are associated with increased uterine contractility and over expression of Oxytocin Receptor (OTR) in women with symptomatic adenomyosis. *Fertil Steril.* 2013; 99: 231-240.
- Harlow SD, Ephross SA. Epidemiology of menstruation and its relevance to women's health. *Epidemiol Rev.* 1995; 17: 265-286.
- Kennedy S. Primary dysmenorrhoea. *Lancet.* 1997; 349: 1116.
- Parker MA, Sneddon AE, Arbon P. The menstrual disorder of teenagers (MDOT) study: determining typical menstrual patterns and menstrual disturbance in a large population-based study of Australian teenagers. 2010; 117: 185-192.
- Sundell G, Milsom I, Andersch B. Factors influencing the prevalence and severity of dysmenorrhoea in young women. *Br J Obstet Gynaecol.* 1990; 97: 588-594.
- Patel V, Tanksale V, Sahasrabhojane M, Gupte S, Nevrekar P. The burden and determinants of dysmenorrhoea: a population-based survey of 2262 women in Goa, India. *BJOG.* 2006; 113: 453-463.
- Ju H, Jones M, Mishra G. The prevalence and risk factors of dysmenorrhea. *Epidemiol Rev.* 2014; 36: 104-113.
- Tavallae M, Joffres MR, Corber SJ, Bayanzadeh M, Rad MM. The prevalence of menstrual pain and associated risk factors among Iranian women. *J Obstet Gynaecol Res.* 2011; 37: 442-451.
- Jarrett M, Heitkemper MM, Shaver JF. Symptoms and self-care strategies in women with and without dysmenorrhea. *Health Care Women Int.* 1995; 16: 167-178.
- Harlow SD, Park M. A longitudinal study of risk factors for the occurrence, duration and severity of menstrual cramps in a cohort of college women. *Br J Obstet Gynaecol.* 1996; 103: 1134-1142.
- Jamieson DJ, Steege JF. The prevalence of dysmenorrhea, dyspareunia, pelvic pain, and irritable bowel syndrome in primary care practices. *Obstet Gynecol.* 1996; 87: 55-58.
- Ortiz MI, Rangel-Flores E, Carrillo-Alarcón LC, Veras-Godoy HA. Prevalence and impact of primary dysmenorrhea among Mexican high school students. *Int J Gynaecol Obstet.* 2009; 107: 240-243.
- Seven M, Güvenç G, Akyüz A, Eski F. Evaluating dysmenorrhea in a sample of Turkish nursing students. *Pain Manag Nurs.* 2014; 15: 664-671.
- Polat A, Celik H, Gurates B, Kaya D, Nalbant M, Kavak E, Hanay F. Prevalence of primary dysmenorrhea in young adult female university students. *Arch Gynecol Obstet.* 2009; 279: 527-532.
- Dawood MY. Dysmenorrhoea and prostaglandins: pharmacological and therapeutic considerations. *Drugs.* 1981; 22: 42-56.
- Grandi G, Ferrari S, Xholli A, Cannoletta M, Palma F, Romani C, Volpe A. Prevalence of menstrual pain in young women: what is dysmenorrhea? *J Pain Res.* 2012; 5: 169-174.
- Melzack R and Katz J. Pain measurement in persons in pain. In: Wall PD, Melzack R (eds). *Textbook of Pain.* 1994, 337-351.
- Riley JL 3rd, Robinson ME, Wise EA, Price DD. A meta-analytic review of pain perception across the menstrual cycle. *Pain.* 1999; 81: 225-235.
- Negriff S, Dorn LD, Hillman JB, Huang B. The measurement of menstrual symptoms: factor structure of the menstrual symptom questionnaire in adolescent girls. *J Health Psychol.* 2009; 14: 899-908.
- Chang AM, Holroyd E, Chau JP. Premenstrual syndrome in employed Chinese women in Hong Kong. *Health Care Women Int.* 1995; 16: 551-561.
- Granot M, Yarnitsky D, Itskovitz-Eldor J, Granovsky Y, Peer E, Zimmer EZ. Pain perception in women with dysmenorrhea. *Obstet Gynecol.* 2001; 98: 407-411.
- Rezaii T, Hirschberg AL, Carlström K, Ernberg M. The influence of menstrual phases on pain modulation in healthy women. *J Pain.* 2012; 13: 646-655.
- Chen CX, Kwekkeboom KL, Ward SE. Self-report pain and symptom measures for primary dysmenorrhoea: A critical review. *Eur J Pain.* 2015; 19: 377-391.
- Wong LP, Khoo EM. Dysmenorrhea in a multiethnic population of adolescent Asian girls. *Int J Gynaecol Obstet.* 2010; 108: 139-142.

43. Daley AJ. Exercise and primary dysmenorrhoea: A comprehensive and critical review of the literature. *Sports Med.* 2008; 38: 659-670.
44. Gould D. Facing the pain of dysmenorrhoea. *Nurs Stand.* 1994; 8: 25-28.
45. Chen CH, Lin YH, Heitkemper MM, Wu KM. The self-care strategies of girls with primary dysmenorrhea: a focus group study in Taiwan. *Health Care Women Int.* 2006; 27: 418-427.
46. Zhou HG, Yang ZW and Students Group. Prevalence of dysmenorrhea in female students in a Chinese university: A prospective study. 2010; 2: 311-314.
47. Banikarim C, Chacko MR, Kelder SH. Prevalence and impact of dysmenorrhea on Hispanic female adolescents. *Arch Pediatr Adolesc Med.* 2000; 154: 1226-1229.
48. Andersch B, Milsom I. An epidemiologic study of young women with dysmenorrhea. *Am J Obstet Gynecol.* 1982; 144: 655-660.
49. Ng TP, Tan NC, Wansaicheong GK. A prevalence study of dysmenorrhoea in female residents aged 15-54 years in Clementi Town, Singapore. *Ann Acad Med Singapore.* 1992; 21: 323-327.
50. Sundell G, Milsom I, Andersch B. Factors influencing the prevalence and severity of dysmenorrhoea in young women. *Br J Obstet Gynaecol.* 1990; 97: 588-594.
51. Weissman AM, Hartz AJ, Hansen MD, Johnson SR. The natural history of primary dysmenorrhoea: a longitudinal study. *BJOG.* 2004; 111: 345-352.
52. Vincent K, Warnaby C, Stagg CJ, Moore J, Kennedy S, Tracey I. Dysmenorrhoea is associated with central changes in otherwise healthy women. *Pain.* 2011; 152: 1966-1975.
53. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: A rural school-based survey. *Aust J Rural Health.* 2011; 19: 218-223.
54. Aziato L, Dedey F, Clegg-Lampsey JN. Dysmenorrhea Management and Coping among Students in Ghana: A Qualitative Exploration. *J Pediatr Adolesc Gynecol.* 2015; 28: 163-169.
55. Houston AM, Abraham A, Huang Z, D'Angelo LJ. Knowledge, attitudes, and consequences of menstrual health in urban adolescent females. *J Pediatr Adolesc Gynecol.* 2006; 19: 271-275.
56. Dawood MY. Dysmenorrhoea and prostaglandins: Pharmacological and therapeutic considerations. *Drugs.* 1981; 22: 42-56.
57. Tangchai K. Dysmenorrhea in Thai adolescents: Prevalence, impact and knowledge of treatment. *J Med Assoc Thai.* 2004; 87: 69-73.
58. Lefebvre G, et al. Primary Dysmenorrhea Consensus Guideline SOGC Clinical Practice Guideline. 2005; 169: 1117-1130.
59. Proctor M, et al. Diagnosis and management of dysmenorrhea. *BMJ.* 2006; 332: 1134-1138.
60. Rostami M. The Study of Dysmenorrhea in High School Girls. *Pak J Med Sci.* 2007; 23: 928-931.
61. Begum J, et al. Menstrual Pattern and Common Menstrual Disorders among Students in Dinajpur Medical College. *Dinajpur Med Col J.* 2009; 2: 37-43.
62. Suresh K, et al. Prevalence of Dysmenorrhea among Adolescent Girls (14-19 Yrs) of Kadapa District and its Impact on Quality of Life: A Cross Sectional Study. *Nat J Comm Med.* 2011; 2: 265-268.
63. Grandi G. Prevalence of menstrual pain in young women: what is dysmenorrhea? *J Pain Res.* 2012; 5: 169-174.
64. Shah M. A study of prevalence of primary dysmenorrhea in young students - A cross-sectional study. *Health line.* 2013.
65. Khamdan HY, Aldallal KM, Almoosa EM, AlOmani NJ, Haider ASM, et al. The Impact of Menstrual Period on Physical Condition, Academic Performance and Habits of Medical Students. *J Women's Health Care.* 2014; 3: 185.
66. Haque SE, Rahman M, Itsuko K, Mutahara M, Sakisaka K. The effect of a school-based educational intervention on menstrual health: an intervention study among adolescent girls in Bangladesh. *BMJ Open.* 2014; 4: e004607.
67. Pitangui, Ana CR, et al. Menstruation Disturbances: Prevalence, Characteristics, and Effects on the Activities of Daily Living among Adolescent Girls from Brazil. *Journal of Pediatric and Adolescent Gynecology.* 2003; 26:148-152.
68. Ahmed MN. Menstrual disorders among Zagazig University Students, Zagazig, Egypt. 2015; 20: 198-203.
69. Zegegne TK, et al. Prevalence of female genital mutilation and its effect on women's health in Bale zone, Ethiopia: a cross-sectional study. *Ethiopia. BMC Public Health.* 2014; 14: 1118-1122.
70. Gagua T, et al. Primary dysmenorrhea: prevalence in adolescent population of Tbilisi, Georgia and risk factors Georgia. *J Turk Ger Gynecol Assoc.* 2012; 13: 162-168.
71. Gumanga SK et al. Menstrual characteristics in some adolescent girls in Accra, Ghana. *Ghana Med J.* 2012; 46: 3-7.
72. Chia CF, et al. Dysmenorrhoea among Hong Kong university students: prevalence, impact, and management. *Hong Kong Med J.* 2013; 19: 222-228.
73. Rani A et al. *Int J Adolesc Med Health.* India. 2015.
74. Al-Asadi JN et al. Iraq. *J Fac Med Baghdad.* 2013; 55: 339-344.
75. Baghianimoghadam MH. A Survey about the Prevalence of Dysmenorrhea in Female Students of Shahid Sadoughi University of Medical Sciences and Their Knowledge, and Practice toward it. *Iran.* 2012; 1: 93-98.
76. De Sanctis V et al. nset of menstrual cycle and menses features among secondary school girls in Italy: A questionnaire study on 3,783 students. *Italy. Indian J Endocr Metab.* 2014; 18: 84-92.
77. Nohara M et al. Menstrual Cycle and Menstrual Pain Problems and Related Risk Factors among Japanese Female Workers. *Japan. Ind Health.* 2011; 49: 228-234.
78. Bata MS. Age at menarche, menstrual patterns, and menstrual characteristics in Jordanian adolescent girls. *International Journal of Gynecology and Obstetrics.* 2012; 119: 281-283.
79. Al-Jefout M, et al. Dysmenorrhea: prevalence and impact on quality of life among young adult jordanian females. (*Jordan*) *J Pediatr Adolesc Gynecol.* 2015; 28: 173-185.
80. Lee JC et al. *Korean J Pediatr.* 2011; 54: 201-206
81. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: a rural school-based survey. *Malaysia Aust J Rural Health.* 2011; 19: 218-223
82. Al-Kindi R and Al-Bulushi A. Prevalence and Impact of Dysmenorrhoea among Omani High School Students. *Oman SQU Med J.* 2011; 11: 485-491.
83. Rodrigues AC, et al. Dysmenorrhea in adolescents and young adults: prevalence, related factors and limitations in daily living. *Portugal Acta Med Port.* 2011; 24 Suppl 2: 383-388.
84. Bano R, et al. Study of the Prevalence and Severity of Dysmenorrhea among the University Students of Hail City. *Saudi Arabia JHRSR.* 2013; 3: 15-22.
85. Wijesiri HS, et al. Knowledge and attitudes towards dysmenorrhea among adolescent girls in an urban school in Sri Lanka. *Nurs Health Sci.* 2013; 15: 58-64.
86. Narring F, et al. Dysmenorrhea: a problem for the pediatrician? (*Switzerland*) *Arch Pediatr.* 2012; 19: 125-130.
87. Pembe AB, Ndolele NT. Dysmenorrhoea and coping strategies among secondary school adolescents in Ilala District, Tanzania. *East Afr J Public Health.* 2011; 8: 232-236.
88. Sahin S, et al. Review of Frequency of Dysmenorrhea and Some Associated Factors and Evaluation of the Relationship between Dysmenorrhea and Sleep Quality in University Students. *Gynecol Obstet Invest. Turkey.* 2014; 78: 179-185.
89. Jang IA, et al. Factors related to dysmenorrhea among Vietnamese and Vietnamese marriage immigrant women in South Korea. *Vietnamese women living in South Korea. Obstet Gynecol Sci.* 2013; 56: 242-248.