

Research Article

Juglans Regia Contain Polyunsaturated Fats, and Essential Fatty Acid Omega-3

Murad S¹, Arain AQ², Saif S³, Shaikh DM⁴, Ghaffar A⁵, Murad JS⁶, Shekhzaid⁷ and Tufail S^{8*}

¹Professor of Pharmacology, at IMDC and DANTh, Islamabad, Pakistan

²Associate Professor of Pharmacology at HBSMDC, Islamabad, Pakistan

³Gynecology Department, NMC Karachi, Pakistan

⁴Professor of Microbiology at JSMU Karachi, Pakistan

⁵CWO at DANTh and IMDC, Pakistan

⁶Psychologist at Bahria University Karachi, Pakistan

⁷Medical College, Rahim Yar Khan, Pakistan

⁸Assistant Professor of Pharmacology, Pakistan

*Corresponding author: Dr. Sana Tufail, Assistant Professor of Pharmacology, Pakistan and Shekhzaid, Medical College, Rahim Yar Khan, Pakistan

Received: July 26, 2019; Accepted: August 29, 2019;

Published: September 05, 2019

Abstract

Background: Hyperlipidemia is one of the important risk factor to develop Coronary Artery Disease (CAD) leading to main cause of morbidity and mortality due to heart attack and cardiac arrhythmias. Decreased level of plasma HDL-cholesterol independently can cause CAD. Recent research studies have proved that walnuts can increase plasma HDL-cholesterol if taken in specific amount per day regularly.

Place and Duration of Study: Research study was conducted at Lahore General Hospital, Lahore from January 2014 to April 2014.

Patient's Inclusion/exclusion Criteria and Grouping: Forty hyperlipidemic patients were selected from Lahore General Hospital for the study. Age of patients ranged from 20 to 65 years. Both gender male and female patients were enrolled. They were advised to discontinue any hypolipidemic medications and foods. Patients suffering from any metabolic disease, renal impairment, liver disease, and already having any cardiac problems were excluded from the research study. Forty patients were divided in two groups, twenty individuals in each group. Group-1 was advised to take 30 grams of walnuts (without shell) per day for the period of eight weeks. Group-II was considered as control group and was advised not to take any dry fruit including walnuts.

Methodology: Their baseline values of HDL-cholesterol were measured at day-0 and day-60, by separating other lipoprotein fractions using chemical precipitation with Mg²⁺, then coupling the products of a cholesterol oxidase reaction.

Results: In twenty hyperlipidemic patients, two months therapy with 30 grams of walnuts per day increase in HDL-cholesterol was 6.3mg/dl which was significant change when analyzed bio statistically, showing p-value <0.01.

Statistical Analysis: Mean values with SD and SEM were analyzed statistically by using SPSS version 16.1. P-value of <0.001 was considered as significant and p-value of >0.05 was considered as non-significant change.

Conclusion: It was concluded from the study that regular intake of specific amount of walnuts without shell can increase good cholesterol (HDL-cholesterol) in male and female patients of age range from 18 to 65 years.

Introduction

High-Density Lipoprotein Cholesterol (HDL-cholesterol) are combinations of fats (lipids) and proteins, by which lipids are transported in the blood. HDLs transport cholesterol from the tissues of the body to the liver, so the cholesterol can be eliminated in the bile. HDL cholesterol is therefore considered the 'good' cholesterol: The higher the HDL cholesterol level, the lower the risk of coronary artery disease. The average man has an HDL cholesterol level of 40 to 50 mg/dl. In the average woman, HDL levels range from 50 to 60 mg/dl. An HDL cholesterol of 60 mg/dL or higher gives some protection against heart disease. Regular aerobic exercise, loss of excess weight (fat), and cessation of cigarette smoking increase HDL cholesterol levels. When lifestyle modifications are insufficient, medications can be used [1-5]. Walnuts (*Juglans regia*) are a tree nut belonging to the walnut family. They originated in the Mediterranean region and Central Asia, and have been part of the human diet for thousands of years⁶. Walnuts

are rich in omega-3 fats and contain higher amounts of antioxidants than most other foods. Eating walnuts may improve brain health while also helping to prevent heart disease and cancer [7]. Besides being a source of high monounsaturated fat, walnuts are a rich source of antioxidants and omega-3 fatty acids [8-10]. Walnuts contain l-arginine, an amino acid that is linked to reduced hypertension, because it causes blood cells to relax and stay smooth. Since all nuts are high in fat, eating too much can cause to gain weight [11]. The fat found in walnuts is mainly omega-3 fats, a type of monounsaturated fatty acid that has heart-protective qualities. Thus, walnuts, Lower total blood cholesterol, Increase HDL [good cholesterol], Decrease LDL [bad cholesterol], Reduce chances of blood clot formation, Reduce inflammation of blood vessels, Relax blood vessels that help to control high blood pressure. Research shows that walnuts [rich in omega-3 fats] boost brain cell activity. Adding some to daily diet helps boost memory in kids and even older people. Recent studies

have also indicated the use of walnuts in the treatment of Alzheimer's disease [12-13].

Subjects and Methods

Research study was conducted at Lahore General Hospital, Lahore from January 2014 to April 2014. Forty hyperlipidemic patients were selected from the hospital for the study. Age of patients ranged from 20 to 65 years. Both gender male and female patients were enrolled. They were advised to discontinue any hypolipidemic medications and foods. Patients suffering from any metabolic disease, renal impairment, liver disease, and already having any cardiac problems were excluded from the research study. Forty patients were divided in two groups, twenty individuals in each group. Group-I was advised to take 30 grams of walnuts (without shell) per day for the period of eight weeks. Group-II was considered as control group and was advised not to take any dry fruit including walnuts. Their baseline values of HDL-cholesterol were measured by separating other lipoprotein fractions using chemical precipitation with Mg^{2+} , then coupling the products of a cholesterol oxidase reaction. Mean values with SD and SEM were analyzed statistically by using SPSS version 16.1. Statistical significant was considered as p-value of changed parameter from day-0 to day-60. P-value of <0.001 was labeled as significant and p-value of >0.05 was considered as non-significant change.

Results

After eight weeks of taking 30 grams of walnuts by 20 patients, their HDL-cholesterol raised from 31.80 ± 1.65 mg/dl to 38.11 ± 1.54 mg/dl. When analyzed statistically it was significant change in the parameter (p-value <0.01). This change in percentage was 6.3 %. In controlled group (n=20) rise in HDL-cholesterol was insignificant statistically (p-value >0.05) (Table 1 & 2).

Discussion

Walnuts are densely packed with vitamin E, copper, manganese and polyphenols. These act as antioxidants that help human body get rid of free radicals created in the body because of exposure to pollution, UV radiations, smoke, stress and junk foods. In research studies conducted at high profile public based hospital settings in India, Bangladesh, Pakistan, Iran, and Srilanka it was proved that oxidative stress is reduced by taking walnuts because these contain omega-3 fatty acids. In our research study rise in HDL-cholesterol taken by 20 hyperlipidemic patients for two months, was 6.3mg/dl. When statistically analyzed it was significant change. These results match with results of study conducted by Rose E et al [14] who observed 8.97mg/dl increase in HDL-cholesterol when hyperlipidemic patients taken 40 grams walnuts per day for three months. Our results also match with results of research study conducted by Blomhoff R et al [15]. They proved 7 mg/dl increase HDL-cholesterol when 20 grams of walnuts were advised to take by hypercholesteremic patients for three months. They made conclusion from their study's result that walnuts are rich in vitamin E which is one of the best fat (lipid) based antioxidants. Oxidative stress causes free radicals, and if they are not reduced or eliminated, they cause deterioration of the cells. This oxidative stress is also caused by normal cell metabolism, but is exasperated by a variety of environmental factors, including UV radiation, pollution, tanning, smoke, heavy-metals in the environment. Vitamin E content walnuts protects the cells and cell

Table 1: Showing mean values before and after consumption of green tea with its p-value.

Baseline Value	After Treatment Value	Change in Parameter	P-value
HDL-C: 31.80 ± 1.65 mg/dl	38.11 ± 1.54 mg/dl	6.3mg/dl	<0.01

Table 2: Showing control group's parameters mean values with its p-value and statistical significance.

Baseline Value	After Treatment Value	Change in Parameter	P-value
HDL-C: 33.39 ± 1.13 mg/dl	33.78 ± 2.67 mg/dl	0.4mg/dl	>0.05

Key: HDL-C stands for high-density lipoprotein cholesterol, \pm indicates SEM. P-value <0.01 indicates significant change and p-value >0.05 indicates non-significant change in parameter.

membranes from free-radical and environmental oxidative stress. They explained that Walnuts also elevate blood levels of melatonin and hence make for a good bedtime snack. Our results are in contrast with results of study conducted by Tapsell LC et al [16] who proved much lesser change in HDL-cholesterol when 10 grams of walnuts were used in 100 hyperlipidemic patients for 4 months. Their results proved only 1.94% mg/dl rise in HDL-cholesterol. Reason for this difference may be much lesser dose or concentration of substance used for pharmacodynamic effects. They have mentioned that structured "whole of diet" advice that included 30g of walnuts/day delivering substantial amounts of polyunsaturated fatty acid improved the lipid profile of patients with type 2 diabetes. Morgan JM et al [17] have explained that polyphenols are over 4,000 distinct species; many of these compounds have antioxidant activity *in vitro* but are unlikely to have antioxidant roles *in vivo*. Rather, they may affect cell-to-cell signaling, receptor sensitivity, inflammatory enzyme activity or gene regulation. Approximately 90% of the phenols in walnuts are found in the skin, including key phenolic acids, tannins, and Flavonoids. Cortes B et al [18] have mentioned that walnuts are a vital food for pregnant women. The good supply of proteins and omega-3 fats are good for baby's brain and eye development. Walnuts help breast-feeding women to maintain a healthy milk supply too. According to Morangoni F et al [19] walnuts have another health benefit. A solution containing 5% of Vitamin E obtained from walnuts decreased the healing time required and Vitamin E increases the breaking strength of wounds. Vitamin E is required for healthy collagen in the skin - which is the support system in the skin and helps the skin to remain firm and healthy.

Conclusion

After two months therapy with 30 grams of walnuts in 20 hyperlipidemic patients, it was observed that HDL-cholesterol was increased by 6.3mg/dl. We concluded that apart from other lipid profile's parameters, HDL-cholesterol might be increased significantly.

References

- Griell AE, Kris-Etherton PM, Hilpert KF, Zhao G, West SG, Corwin RL. An increase in dietary n-3 fatty acids decreases a marker of bone resorption in humans. *Nutr J.* 2007; 6: 2.
- Kelly JH Jr, Sabate J. Nuts and coronary heart disease: an epidemiological perspective. *Br J Nutr.* 2006; Suppl 2: S61-7.
- Rosenson RS, Brewer HB Jr, Ansell B, Barter P, Chapman MJ, Heinecke JW, et al. Translation of high-density lipoprotein function into clinical practice: current prospects and future challenges. *Circulation.* 2013; 128: 1256-1267.
- Patel G. Essential fats in walnuts are good for the heart and diabetes. *J Am*

- Diet Assoc. 2005; 105: 1096-1097.
5. Degoma EM, Rader DJ. Novel HDL-directed pharmacotherapeutic strategies. *Nat. Rev. Cardiol.* 2011; 8: 266-277.
 6. Riwanto M, Landmesser U. High density lipoproteins and endothelial functions: mechanistic insights and alterations in cardiovascular disease. *J. Lipid Res.* 2013; 54: 3227-3243.
 7. Tsai CJ, Leitzmann MF, Hu FB, Willett WC, Giovannucci EL. Frequent nut consumption and decreased risk of cholecystectomy in women. *Am J Clin Nutr.* 2004; 80: 76-81.
 8. Zhao G, Etherton TD, Martin KR, West SG, Gillies PJ, Kris-Etherton PM. Dietary {alpha}-Linolenic Acid Reduces Inflammatory and Lipid Cardiovascular Risk Factors in Hypercholesterolemic Men and Women. *J Nutr.* 2004; 134: 2991-2997.
 9. Anderson KJ, Teuber SS, Gobeille A, Cremin P, Waterhouse AL, Steinberg FM. Walnut polyphenolics inhibit *in vitro* human plasma and LDL oxidation. *Journal of Nutrition.* 2009; 131: 2837-2842.
 10. Gillen LJ, Tapsell LC, Patch CS, Owen A, Batterham M. Structured dietary advice incorporating walnuts achieves optimal fat and energy balance in patients with type 2 diabetes mellitus. *J Am Diet Assoc.* 2005; 105: 1087-1096.
 11. Bes-Rastrollo M, Sabate J, Gomez-Gracia E, Alonso A, Martinez JA, Martinez-Gonzalez MA. Nut consumption and weight gain in a Mediterranean cohort: The SUN study. *Obesity.* 2007; 15: 107-116.
 12. Fukuda T, Ito H, Yoshida T. Antioxidative polyphenols from walnuts (*Juglans regia L.*). *Phytochemistry.* 2003; 63: 795-801.
 13. Barter P. Lessons learned from the Investigation of Lipid Level Management to understand its Impact in Atherosclerotic Events (ILLUMINATE) trial. *Am. J. Cardiol.* 2009; 104: 10E-15E.
 14. Ros E, Nunez I, Perez-Heras A, Serra M, Gilabert R, Casals E, Deulofeu R. A walnut diet improves endothelial function in hypercholesterolemic subjects: a randomized crossover trial. *Circulation.* 2004; 109: 1609-1614.
 15. Blomhoff R, Carlsen MH, Andersen LF, Jacobs DR Jr. Health benefits of nuts: potential role of antioxidants. *Br J Nutr.* 2006; 96: S52-60.
 16. Tapsell LC, Gillen LJ, Patch CS, Batterham M, Owen A, Bare M, et al. Including Walnuts in a Low-Fat/Modified-Fat Diet Improves HDL Cholesterol-to-Total Cholesterol Ratios in Patients With Type 2 Diabetes. *Diabetes Care.* 2004; 27: 2777-2783.
 17. Morgan JM, Horton K, Reese D, Carey C, Walker K, Capuzzi DM. Effects of walnut consumption as part of a low-fat, low-cholesterol diet on serum cardiovascular risk factors. *Int J Vitam Nutr Res.* 2002; 72: 341-347.
 18. Cortes B, Nunez I, Cofan M, Gilabert R, Perez-Heras A, Casals E, et al. Acute effects of high-fat meals enriched with walnuts or olive oil on postprandial endothelial function. *J Am Coll Cardiol.* 2006; 48: 1666-1671.
 19. Marangoni F, Colombo C, Martiello A, Poli A, Paoletti R, Galli C. Levels of the n-3 fatty acid eicosapentaenoic acid in addition to those of alpha linolenic acid are significantly raised in blood lipids by the intake of four walnuts a day in humans. *Nutr Metab Cardiovasc Dis.* 2006; 123: 2728-2741.