

Special Article – Pulmonary Rehabilitation

Efficacy of Preoperative Comprehensive Pulmonary Rehabilitation in Patients with Lung Cancer

Kaymaz D^{1*}, Ergün P¹, Candemir I¹, Gülhan SE², Fındık G² and Demir N²¹Pulmonary Rehabilitation and Home Care Center, Ataturk Chest Diseases and Chest Surgery Training and Research Hospital, Ankara, Turkey²Clinic of Chest Surgery, Ataturk Chest Diseases and Chest Surgery Training and Research Hospital, Ankara, Turkey***Corresponding author:** Dicle Kaymaz MD, Department of Pulmonary Rehabilitation and Home Care Center, Atatürk Chest Diseases and Chest Surgery Training and Research Hospital 06280, Keçiören, Ankara, Turkey**Received:** October 04, 2016; **Accepted:** October 28, 2016; **Published:** November 02, 2016**Abstract****Introduction:** Preoperative pulmonary rehabilitation (PPR) can optimize individuals' exercise tolerance and overall medical stability before lung cancer resection surgery. Our aim was to evaluate the efficacy of pulmonary rehabilitation (PR) before lung cancer surgery.**Materials and Methods:** Fifteen consecutive patients with potentially resectable non small cell lung cancer (14 men) of mean age 66.6±5.87 (range 56-77) were included to this study. Short-term, compact, comprehensive, supervised PPR program consisted of bronchial hygiene, breathing control, energy conservation techniques, exercise training (endurance and strength), psychological support, nutritional support. Exercise capacity and VO₂ peak were evaluated by using incremental shuttle walk test (ISWT). Dyspnea sensation was assessed with MRC scale, psychological status with Hospital Anxiety and Depression (HAD) scale, the bioelectrical impedance analysis (BIA) was performed for the estimation of body composition on admission and after PPR.**Results:** After PPR significant beneficial changes were also observed in VO₂ peak (p=0.000), ISWT (p=0.000), MRC (p=0.004), anxiety (p=0.002), depression (p=0.001) scores. According to the results of evaluation of body composition mean fat free mass index (FFMI) was 19.1±1.6 kg/m² where it was 19.2±1.5 kg/m² at the end of PPR (p=0.184).**Conclusion:** In this study it was shown that short term, compact, comprehensive, multidisciplinary, PR programs before lung cancer surgery is potentially an attractive technique for optimizing preoperative exercise capacity, risk modification, dyspnea sensation, psychological status and conceivably, in turn, reducing postoperative respiratory complications.**Keywords:** Pulmonary Rehabilitation; Lung Cancer; Cancer patients**Introduction**

Lung cancer is a catastrophic and progressive disease that carries excessive social and economic burden. Eventhough the overall five year survival is poor, targeted therapeutics, early detection and multidisciplinary approaches to diagnosis and management have trend to improve outcomes. One of the recent interdisciplinary approach is pulmonary rehabilitation (PR) in perioperative period. PR has been shown to decrease post-operative complications as well as hospital length of stay in patients who underwent lung resection [1]. Low exercise tolerance is associated with poor thoracic surgical outcomes. Preoperative pulmonary rehabilitation (PPR) can optimize individuals' exercise tolerance and overall medical stability before lung cancer resection surgery. Pulmonary resection is the treatment of non small lung cancer (NSCLC) in early stage. The measurement of peak oxygen consumption (VO₂ peak) has been shown to be the strongest independent predictor of surgical complication rate. Specifically, NSCLC patients with a preoperative VO₂ peak ≥ 15 mL.kg⁻¹.min⁻¹ are at comparatively low risk of complications, whereas patients with ≤15 mL.kg⁻¹.min⁻¹ and ≤10 mL.kg⁻¹.min⁻¹ are at increased and very high risk of complications, respectively [2,3].

The aim of this study was to evaluate compact, short-term

comprehensive PR program effectiveness in NSCLC patients during preop period.

Materials and Methods

This study was performed in our Pulmonary Rehabilitation Center. Fifteen consecutive patients with potentially resectable NSCLC (14 men) of mean age 66.6±5.87 (range 56-77) were included to this study. Characteristics of the participants are shown in Table 1. For evaluation of exercise capacity of these patients, field tests [Incremental Shuttle Walking Test (ISWT)] was used. Peak VO₂ was calculated with a formulation of [4.19 + (walking distance x 0.025)] (mL.min⁻¹.kg⁻¹) [4,5].

Dyspnea sensation was assessed with MRC scale [6] and psychological status with Hospital Anxiety and Depression scale [7], the bioelectrical impedance analysis (BIA) was performed for the estimation of body composition [8] on admission and after PPR. Pulmonary rehabilitation programme was multidisciplinary, comprehensive, compact (5 consecutive days in a week) and lasted two weeks, totally 10 sessions. In patients with transportation problem, the programme was undertaken inpatient manner. PPR programme consisting of: (a) educational support, medication advices, bronchial hygiene techniques, and breathing control techniques, energy

Table 1: Characteristics of the Participants (n=15).

Variable	No	%
Age, mean±SD, year	66.6±5.87	
Male, %		93.3
FFMI, mean±SD, kg/m ²	19.1±1.6	
Smoking, mean±SD, pack/year	50.27±29.08	
Diagnosis	15	100
Nonsmallcelllungcancer		
Extent of resection		
Lobectomy	5	33.3
Pneumonectomy	1	6.66
Pulmonary Function Data		
FEV1 liters,	42±14	
FVC liters,	55±19	
FEV1/FVC	58±14	
Exercisecapacity data		
VO ₂ peak, mL.kg ⁻¹ .min ⁻¹	11.16±2.71	
ISWT, meters	279.07±110.30	

SD: Indicates standard deviation; FFMI: Fatfree mass index; FEV1: Forced expired volume; VO₂ peak: Peak oxygen consumption; ISWT: Incremental Shuttle Walk Test.

Table 2: PPR values of medically inoperabl patient's.

n=4	Before PPR program	After PPR program
ISWT (m)	199±108.6	272±69.9
VO ₂ Peak(mL .kg ⁻¹ .min ⁻¹)	9.17±2.7	11±1.7

Table 3: The values before and after PPR.

n=15	Before PPR program	After PPR program	p
MRC	2.67±0.81	2.20±0.56	0.004
ISWT(m)	279.07±110.30	347±106.53	0.000
VO ₂ PeakmL .kg ⁻¹ .min ⁻¹	11.16±2.71	12.85±2.66	0.000
FFMI (kg/m ²)	19.1±1.6	19.2±1.5	0.184
Anxiety	10.50±1.34	8.79±2.0	0.002
Depression	9.64±1.49	7.21±2.39	0.001

conservation, relaxation, and dietary advices. Educational sessions were delivered by two chest physicians, two physical therapists, a dietician, one respiratory nurses, and a psychologist. (b) exercise training, (c) a nutritional intervention, and (d) psychological counseling, if needed. The exercise training program was individually tailored to each patient. All exercise training sessions were supervised by physical therapist. Exercise included cycle ergometer training (15 min), treadmill training (15 min), upper and lower extremity strength training (5-10 min), breathing therapies (10-20 min), and relaxation therapies (5-10 min) for total 50-70 min/day. Patients underwent both cycle ergometer and treadmill training. Both workload for cycling and walking speed for treadmill ergometer were calculated from incremental shuttle walking test (ISWT) results using formulations and BORG dyspnea scores 4-6 were also used for prescribing exercise [9]. Patients were trained at 50% of peak workload and 50-80% of peak VO₂. Quadriceps resistance training was applied using free weights for 5 consecutive days in a week according to 1-repetition maximum starting at 50% for three sets and 10 repetitions per set in the 10 sessions. Upper extremity training consisted of one set, 10 repetitions per set totally 10 sessions. Loads were recorded as kilograms.

Exercises intensity increased according to the patient progress. Pulse oximetry was used to supervise patients during exercise. If

the SpO₂ fell below 90%, oxygen supplementation was provided to maintain SpO₂ ≥ 90%.

Outcomes

The primary outcome was change in VO₂ peak mL.kg⁻¹.min⁻¹ between baseline and immediately before pulmonary resection (presurgery). Secondary outcome was the evaluation of PPR efficacy in patients with lung cancer

Statistical analysis

The statistical analysis was performed using Statistical Package for Social Sciences (SSPS) version 15.0 (Statistical Package for Social Sciences (SSPS), Chicago, IL, USA). Data were given as mean±standard deviation. First, the variables were analyzed with Shapiro–Wilks test for normal distribution. Then, the pairedt test was used for variables with normal distribution and Wilcoxon signed rank test used for those variables without normal distribution. Spearman correlation analyses were performed. Statistical significance was determined as probability value of <0.05 was considered to be statistically significant.

Results

Five (33%) of the patients underwent lobectomy where one (6.66%) of them had pneumonectomy. In one patient who had metastasis in lenf node dissection by mediastinoscopy, has given cheomoradiotherapy. Four (26.7%) of them refused the operation. In four (26.7%) patients mean VO₂ peak was 9.2±2.7 mL .kg⁻¹ .min⁻¹ so they were accepted as medically inoperable with a FEV1 790±223 mL before PPR (Table 2). In this group of patients even VO₂ peak increased to a level of 1.84±0.9, they did not have surgical resection. The mean FEV1 and FVC of all patients were 1164±470 mL, 42±14% predicted, 2020±714 mL, 55±19 % predicted respectively. ISWT distances ranged from 46 m to 450 m (mean 279.07±110.30 m). The mean VO₂ peak calculated by using ISWT was 11.16±2.71 mL.kg⁻¹ .min⁻¹, Δ.VO₂ peak; 1.69±0.1 mL .kg⁻¹ .min⁻¹ (p <0.001) and Δ ISWT ; 68.26±4 m (p <0.001). Significant beneficial changes were also observed in VO₂ peak (p=0.000), ISWT(p=0.000), MRC (p=004) , anxiety (p=0.002), depression (p= 0.001) scores. According to the results of evaluation of body composition mean FFMI was 19.1±1.6 Kg/m² where it was 19.2±15kg/m² at the end of PPR (Table 3).

There was no statistically significant correlation between baseline lung function tests (FEV1 %, ml) and either baseline or changes in VO₂ peak, ISWT distance values.

Discussion

Surgery is the only curative-intent treatment for patients with resectable NSCLC. The principal finding of this study was that a short-term,compact, comprehensive, supervised PPR program was effective, feasible and safe among newly diagnosed NSCLC patients before lung cancer resection surgery.

Jones et al has shown that preoperative exercise could increase the numbers of candidates eligible for curative-intent pulmonary resection [10]. In their study they showed an increase in VO₂ peak according to cardiopulmonary exercise test (CPET). Eventhough CPET is gold standart for evaluating exercise capacity, in most countries often poorly available and it should be performed by expericenced stuff. ISWT which is readily available in most hospitals correlates well with

VO₂ peak in (CPET) [4] and has been widely used as a measure of exercise capacity in many clinical studies. The shuttle walk is a useful exercise test to assess potentially operable lung cancer patients with borderline lung function [10]. In our study after PPR, VO₂ peak which was calculated by using ISWT was increased significantly and by having risk modification six patients (40%) could be operated. In patients with lung cancer performance status is one the most important factor that influence the therapeutic choice. In our study in four (26.7%) patients mean VO₂ peak was 9.1±2.7 mL.kg⁻¹.min so they were accepted as medically inoperable with a FEV1 of 790±223 mL. After PPR eventhough risk level of these patients decreased VO₂ peak did not reach an acceptable level for surgery. So two of them underwent steriotactic radiotherapy, and two of them decided to have supportive therapy. Though surgery remains the treatment of choice for resectable lung carcinoma, postoperative complications of lung resection appear to be a major problem especially in patients with co-existing disease [11]. Incidence of postoperative complications after lung resection was reported as 24-48% while the mortality rates of lobectomy and pneumonectomy were 4% and 14%, respectively [11,12]. The most common complications and the main determinant of mortality and morbidity after lung resection are cardiopulmonary complications. Beckles et al stated that the pre-operative physiologic assessment of patients being considered for surgical resection of lung cancer must consider the immediate peri-operative risks from comorbid cardiopulmonary disease, the long-term risks of pulmonary disability, and the threat to survival due to inadequately treated lung cancer [11]. In our study one patient experienced 3 complications (respiratory failure, pulmonary embolism and death) whereas one patient experienced 1 complication (air leak). The average duration of hospital stay was 18,8 days. In this present study the complication rates in patients underwent surgical resection was 33.3%. Knowing that these patients also had the diagnosis of COPD as a comorbidity, we can say that PPR reduced the risk of complications. In comprehensive pulmonary rehabilitation programmes supervised exercise combined with stress management education in pulmonary rehabilitation may offer management strategies for persons with anxiety and depression. The prevalence rates of anxiety and depression in lung cancer survivors are 34% and 33%, respectively [13]. In a study exercise training has been shown positive effects on anxiety and depression in patients with lung cancer [14]. Our PPR programme was short, compact but comprehensive. The decrease in anxiety and depression scores of our patients (Δ Anxiety score=1.71, Δ Depression score=2.43 was statistically significant and over the level of MCID [15]. Most lung cancer patients suffer from nutritional issues. Although our patients FFMI values did not change at the end of the PPR, patients were able continue a healthy diet and educated on strategies to improve their nutritional needed if necessary.

There is some limitations of this study. First of all designing a randomised controlled study for candidates of lung cancer resection surgery is not ethical. Relatively small sample size of the patients underwent surgical resection is one the handicap of assessing the complication rates.

Conclusion

In conclusion short term, compact, comprehensive pulmonary rehabilitation programmes before lung cancer surgery is potentially an attractive technique for optimizing preoperative exercise capacity, risk modification and conceivably, in turn, reducing postoperative respiratory complications.

References

- Nagarajan K, Bennett A, Agostini P, Naidu B. Is preoperative physiotherapy/pulmonary rehabilitation beneficial in lung resection patients? *Interact Cardiovasc Thorac Surg*. 2011; 13: 300-302.
- Beckles MA, Spiro SG, Colice GL, Rudd RM. The physiologic evaluation of patients with lung cancer being considered for resectional surgery. *Chest*. 2003; 123: 105S-114S.
- Win T, Jackson A, Groves AM, Sharples LD, Charman SC, Laroche CM. Comparison of shuttle walk with measured peak oxygen consumption in patients with operable lung cancer. *Thorax*. 2006; 61: 57-60.
- Singh SJ, Morgan MD, Hardman AE, Rowe C, Bardsley PA. Comparison of oxygen uptake during a conventional treadmill test and the shuttle walking test in chronic airflow limitation. *Eur Respir J*. 1994; 7: 2016-2020.
- Singh SJ, Morgan MD, Scott S, Walters D, Hardman AE. Development of a shuttle walking test of disability in patients with chronic airways obstruction. *Thorax*. 1992; 47: 1019-1024.
- Mahler DA. Measurement of dyspnea Clinical ratings. In: Mahler DA, O'Donnell DE (eds). *Dyspnea*. 2nd Ed. Lung Biology in Health and Disease. Taylor and Francis, Boca Raton. 2005; 208: 147-165.
- Aydemir Ö. Validity and reliability of Turkish version of hospital anxiety and depression scale. *Türk Psikiyatri Dergisi*. 1997; 8: 280-287.
- Walter-Kroker A, Kroker A, Mattiucci-Guehlke M, Glaab T. A practical guide to bioelectrical impedance analysis using the example of chronic obstructive pulmonary disease. *Nutr J*. 2011; 10: 35.
- Bestall JC, Paul EA, Garrod R, Garnham R, Jones PW, Wedzicha JA. Longitudinal trends in exercise capacity and health status after pulmonary rehabilitation in patients with COPD. *Respir Med*. 2003; 97: 173-180.
- Jones LW, Peddle CJ, Eves ND, Haykowsky MJ. Effects of Presurgical Exercise Training on Cardiorespiratory Fitness Among Patients Undergoing Thoracic Surgery for Malignant Lung Lesions. *Cancer*. 2007; 110: 590-598.
- Beckles MA, Spiro SG, Colice GL, Rudd RM. American College of Chest Physicians the physiologic evaluation of patients with lung cancer being considered for resectional surgery. *Chest*. 2003; 123: S105-S114.
- Richter Larsen K, Svendsen UG, Milman N, Brenoe J, Petersen BN. Exercise testing in the preoperative evaluation of patients with bronchogenic carcinoma. *Eur Respir J*. 1997; 10: 1559-1565.
- Hopwood P, Stephens RJ. Depression in patients with lung cancer: prevalence and risk factors derived from quality-of-life data. *J Clin Oncol*. 2000; 18: 893-903.
- Chen H-M, Tsai C-M, Wu Y-C, Lin K-C. Randomised controlled trial on the effectiveness of home-based walking exercise on anxiety, depression and cancer-related symptoms in patients with lung cancer. *Bjancer*. 2015;112: 438-445.
- Puhan MA, Frey M, Büchi S, et al. The minimal important difference of the hospital anxiety and depression scale in patients with chronic obstructive pulmonary disease. *Health Qual Life Outcomes*. 2008; 6: 46.