

Research Article

Isolation and Evaluation of Endophytic Bacteria Against *Fusarium Oxysporum* f. sp. *Cucumerinum* Infecting Cucumber Plants

Ozaktan H^{1*}, Çakır B², Gül A¹, Yolageldi L², Akköprü A³, Fakhraei D¹ and Akbaba M¹

¹Department of Plant Protection, University of Ege Faculty of Agriculture, Turkey

²Department of Horticulture, University of Ege, Faculty of Agriculture, Turkey

³Department of Plant Protection University of Yüzüncü Yıl, Turkey

*Corresponding author: Birsen Çakır, Department of Horticulture, Faculty of Agriculture, Ege University, Bornova, Izmir, Turkey

Received: August 04, 2014; Accepted: February 02, 2015; Published: February 23, 2015

Abstract

The aim of this study was to isolate, within cucumber plants, Endophytic Bacterial (EB) isolates that can provide significant biological control of Fusarial wilt of cucumber caused by *Fusarium Oxysporum* f. sp. *Cucumerinum* (FOC) and enhance plant growth under *in vitro* conditions. The endophytic bacteria were isolated from the internal tissues of roots, leaves and stems of healthy cucumber plants. In this study, we isolated 112 EB strains from internal tissues of healthy cucumber plants grown in greenhouse and field conditions in Turkey. We determined several phenotypic properties of EB strains and found approximately equal numbers of Gram-negative and Gram-positive strains. These isolates were screened *in vitro* for their plant growth promoting traits such as production of Indol 3-Acetic Acid (IAA), Hydrogen Cyanide (HCN), siderophore, phosphate solubilization and antagonistic activity against FOC. More than 30% of the EB strains produced detectable levels (20-125 µg ml⁻¹) of IAA in culture filtrates. 46 % of EB strains exhibited siderophore production ranging from 3 to 19 mm zones. HCN production was more common trait of *Pseudomonas* strains (16%). Solubilization of phosphate was detected by 29% in the EB isolates. More than 53% of the EB strains inhibited the mycelial growth of FOC on PDA plates. The strains CC29/3 and CC25/2 were more active compared to other against FOC. Majority of isolated endophytes were not only able to suppress pathogenic fungi, but could also improve seed germination and plant growth. Finally, these strains may be candidates for biological control and plant growth promotion.

Keywords: *Fusarium oxysporum* f. sp. *cucumerinum*; Endophytic bacteria; Biological control

Abbreviations

FOC: *Fusarium Oxysporum* f. sp. *Cucumerinum*; EB: Endophytic Bacteria; IAA: Indol 3-Acetic Acid, HCN: Hydrogen Cyanide

Introduction

Fusarium Oxysporum f. sp. *Cucumerinum* (FOC), which is a soil-borne fungus, causes wilting and death of cucumber plants grown in greenhouse [1]. The symptoms include necrotic lesions of the stem base, foliar wilting and eventually plant death [1,2]. FOC has been identified in all cucumber growing regions around the world, including Turkey [3,4] and has been documented as an important economic threat to cucumber producers [3,5-7]. It is difficult to control of FOC since the pathogen could cause systemic invasion and move in the cucumber plant tissues by xylem vessels [1]. Chemical control methods which are effective against *Fusarium* root and stem rot of cucurbits are limited [8]. Therefore, control strategies focus mostly on preventing the pathogen from being introduced into disease-free areas, and the development of disease resistant varieties [9]. However, the development of new pathogenic races of FOC limits use of disease resistant varieties [3]. To date, there is no resistant cucumber cultivars to FOC. In response to environmental and health concerns about extended use of pesticides, there is considerable interest in finding alternative control approaches for use in integrated

pest management strategies for crop diseases.

Biological control offers potential alternatives to combat many soil-borne pathogen, including *Fusarium oxysporum* [10]. The biological control of *Fusarium* wilt of cucumber caused by FOC was obtained using siderophore fluorescent *pseudomonads* in Turkey [11].

The use of Endophytic Bacteria (EB) strains to control plant-pathogenic bacteria and fungi is receiving increasing attention as a sustainable alternative to synthetic pesticides. EB strains, which live inter- and intracellularly in plants without inducing pathogenic symptoms, interact with the host biochemically and genetically. EB may play many important beneficial roles in the metabolism and physiology of the host plant, including fixing atmospheric nitrogen, sequestering iron from the soil, solubilizing phosphates, synthesizing plant – growth hormones, and suppressing of ethylene production by 1-Amino Cyclopropane-1-Carboxylate (ACC) deaminase, degrading toxic compounds, inhibiting strong fungal activity and antagonizing bacterial pathogens [12-14]. The internal plant tissues provide a protective environment for endophytic bacteria, which colonize an ecological niche similar to plant pathogens. Endophytes, like *Pseudomonas*, *Agrobacterium*, *Bacillus*, *Burkholderia* and *Enterobacteria*, have been isolated from root nodules in various leguminous plants including alfalfa, clover, soybean pigeon pea, etc

[15] since 1902 [16-18]. Available reports indicated improved plant yield and health under greenhouse conditions (measured as an increase in root wet weight and nodulation) when co-inoculated with nodule endophytes compared to inoculation with rhizobia alone.

In order to reduce the input of pesticides and fertilizers and to make an eco-friendly agriculture, it will be important to develop inocula of biofertilizers, and biopesticides. The main aims of this study are: (i) to collect different EB from different area in Turkey, (ii) to screen these bacteria for a number of plant-beneficial traits, (iii) to test the selected potentially beneficial strains for their abilities to promote growth of cucumber and to control FOC caused by *Fusarium oxysporum* f. sp. *Cucumerinum*.

Materials and Methods

Isolation of endophytic bacteria

The endophytic bacteria were isolated from the internal tissues of roots, leaves and stems of healthy cucumber plants, which were surface-sterilized by sequential immersion in 70% ethanol for 5 min and a solution of 5% sodium hypochlorite for 10 min. Then the samples were rinsed three times in sterile distilled water to remove surface sterilizing agents prior to obtain bacterial isolates. Bacterial strains were isolated by two different techniques such as trituration of leaves and imprinting of stem and root tissues onto Tryptic Soy Agar (TSA). Surface sterility test was performed for each of the samples to ensure the elimination of surface microorganisms. If no bacterial growth occurred in the sterility test, the recovered bacteria were considered to be endophytes. Single colonies were isolated and maintained in pure cultures at -80°C in 15% (v/v) glycerol [19].

Phenotypic characterization of bacterial strains

Colonies of bacterial isolates were characterized for the following traits: color, form, elevation, margin, diameter, surface, opacity, and texture. The Gram reaction was performed by using a 3% KOH test [20]. Endophytic Bacteria (EB) were tested for Hypersensitive Response (HR) on tobacco leaves. EB strains, showed HR (+) on tobacco leaves were considered as possible plant pathogens and were not taken for further tests [21,22].

In vitro screening for antagonistic activity

All EB isolates were screened for their *in vitro* biocontrol activity toward FOC on Potato Dextrose Agar (PDA) using a dual-culture technique [23]. The plates inoculated with the pathogen alone were maintained as control. The mycelial disc (5 mm) from 7 days old culture of FOC was placed on one side of the plate containing PDA medium, and then EB strains were streaked on the opposite side of the plate by the help of sterilized inoculation needle. Three replications were performed for each treatment. The plates were incubated at room temperature for seven days. The inhibitory effects of EB strains on the linear growth of FOC were determined. The percent of inhibited FOC was calculated by comparison with fungal growth in control plates.

In vitro characterization of EB strains for plant growth promotion

The EB strains were screened by *in vitro* assays for the production of the following functional traits: hydrogen cyanide, HCN [24]; phosphate solubilization [25] and Indole 3- Acetic Acid, IAA [26];

plant growth promotion and siderophore production [27]. All experiments were replicated twice for each of the strains.

Effect on seed germination and seed vigor index

EB strains, which were found successful by *in vitro* assays for plant growth promotion and antagonistic activity to FOC, were further analyzed for seed emergence (cv. Gordion) and measured for Vigor Index (VI). For bacterization, seeds were surface sterilized with 1% sodium hypochlorite for 1 min and soaked in EB suspensions amended with 1% Carboxy Methyl Cellulose (CMC). After bacterization, the seeds were placed onto sterile filter paper moistened with Sterile Distilled Water (SDW) in petri plates (three plates with 10 seeds/plate) and incubated at room temperature. Control plates were arranged in a similar way, except that they were treated only with 1% CMC. For each isolate, effects on seed germination were measured by counting the number of fully germinated seeds per plate and comparing that with that of the control plates. After 5 days, the vigor index for each treatment was calculated by using the formula: VI = Percent germination X (seedling length + root length) as described previously [28].

Results and Discussion

Isolation of EB strains

We observed that the inner tissues of healthy cucumber plants were very rich for EB colonization. Trituration and imprinting of the plant tissues were the reliable and easy isolation techniques for EB. A total of 44 healthy cucumber plants including 34 plants grown in greenhouse and 10 plants grown in field were obtained from different sampling areas in Turkey. Surface sterility tests were performed for each sample to monitor the efficiency of the disinfection procedure during isolation. If no bacterial growth occurred in the sterility test, the recovered bacteria were considered to be endophytes. Here, 112 different endophytic colonizing bacterial strains were isolated from healthy cucumber plants grown in greenhouse and field conditions. On the basis of phenotypic identification tests such as some cultural, morphological and biochemical characteristics, a total of 104 endophytic bacterial strains were grouped into Gram (-) bacteria, and Gram (+) bacteria. Most of the EB strains were fluorescent *Pseudomonads*, Gram negative (66%) and rest Gram positive (Table 1). Among Gram-negative soil bacteria, *Pseudomonas* is the most abundant genus in the rhizosphere [29]. Root-associated *Pseudomonas* spp. strains have long been known to be beneficial to plants attribute to their Plant-Growth Promotion Effect (PGPE) or their potential as biological control agents. In addition, endophytic *Pseudomonas* spp. can also indirectly induce PGPE by controlling phytopathogens or pathogenic fungi using mechanisms such as producing antibiotic factors [30,31], enhancing competition for colonization sites [32], and induction of systemic resistance [33]. The diversity of EB reported here has many similarities with the EB isolated from other plants. *Agrobacterium*, *Arthobacter*, *Bacillus*, *Chryseobacterium*, *Enterobacter*, *Pseudomonas* and *Sterotrophomonas* were commonly identified from roots of cucumber, sugar beet, corn, and lemon [34].

In vitro plant growth promoting traits

One of the mechanisms of stimulation of plant growth by bacteria involves the production of phytohormones such as auxins,

Table 1: Overview of plant-beneficial traits of selected endophytic bacteria.

EB isolates	Bacterial Species	The percent inhibition of mycelial development of FOC ^{1,2}	IAA ³ (µg/ml)	Gram Staining	Florescent pigmentation	HCN ⁴	Siderophore production	Phosphate solubilization
CB1/1	2	38,1	31	-	-	-	1 mm	0
CB1/2	3	15,7	25	-	-	-	0 mm	0
CB2/1	4	44,7	30	+	-	-	0 mm	0
CB2/2	5	50	100	-	-	-	12 mm	0
CB2/3	6	57,8	44	-	-	-	10,5 mm	0
CB3	7	30,2	17	+	-	-	0 mm	0
CB4	8	28,9	37	-	-	-	5 mm	3 mm
CC4	9	26,6	27	-	-	-	1 mm	1 mm
CB5/2	11	40,7	26	-	-	-	7 mm	1 mm
CB7	12	34,2	22	-	-	-	5 mm	2 mm
CC7/1	13	56,7	12	-	+	-	13 mm	0 mm
CC7/2	14	53,9	30	-	-	-	5 mm	2,5 mm
CB8/1	15	27,6	37	-	-	-	5 mm	3 mm
CB8/2	16	18,5	32	+	-	-	10 mm	0
CB9/2	18	31,5	13	-	-	-	7 mm	1 mm
CB9/3	19	27,6	33	-	-	-	12 mm	0
CA10	20	39,4	6	+	+	-	0 mm	0 mm
CB10	21	32,8	6	-	-	-	12 mm	9 mm
CC13	22	0	6	+	-	-	0 mm	0
CA13/1	23	15,7	6	+	-	-	0 mm	0
CA13/2	24	0	5	-	-	-	0 mm	0
CB13	25	18,5	15	+	-	-	0 mm	0
CA15	26	0	3	-	-	-	0 mm	2 mm
CB15/1	27	0	5	+	-	-	0 mm	0
CB15/2	28	23,6	10	-	-	-	0 mm	0
CA17/1	29	14,4	12	-	-	-	0 mm	0
CA17/2	30	28,9	6	-	-	-	9 mm	0
CA17/3	31	26,3	9	-	+	+	7 mm	0
CA17/4	32	44,7	27	+	-	-	2 mm	0
CB17/1	33	0	6	+	-	-	0 mm	0
CB17/2	34	0	0	-	-	-	0	0
CA18	35	17,1	7	+	-	-	6,5 mm	0
CB18/1	36	0	6	+	-	-	0 mm	1 mm
CB18/2	37	14,4	7	-	-	-	7 mm	1 mm
CB20/1	38	0	7	+	-	-	0 mm	0
CB20/2	39	0	11	-	+	-	8 mm	3 mm
CB20/3	40	35,7	9	-	-	-	1 mm	1 mm
CC23/1	42	4,2	7	-	-	-	2 mm	2 mm
CC23/2	43	0	10	+	-	-	1 mm	2 mm
CA24	44	0	6	+	-	-	0mm	1 mm
CB24	45	30	18	+	-	-	1 mm	2 mm
CA25	46	14,2	32	-	-	-	3 mm	0
CB25	47	0	12	+	-	-	3 mm	1 mm
CC25/1	48	7	26	-	+	-	14 mm	0
CC25/2	49	64,2	34	-	+	-	9 mm	1 mm
CC26	50	58,5	16	-	-	-	0 mm	0
CA27/1	51	0	9	-	-	-	0 mm	0
CA27/2	52	8,5	9	-	+	+	11 mm	4 mm
CC27	53	18,5	9	-	+	+	13 mm	4 mm
CB27/1	54	0	7	+	-	-	0 mm	0
CB27/2	55	0	6	+	-	-	0 mm	0
CA28/1	56	45,7	13	-	-	-	12 mm	0
CA28/2	57	34,2	35	-	+	-	6 mm	0
CA28/3	58	15,7	15	-	+	-	19 mm	0
CB28	59	24,2	0	+	-	-	0 mm	0

CA29/1	60	27,1	25	-	+	-	3 mm	2 mm
CA29/2	61	27,1	14	-	+	-	16 mm	0
CC29/2	65	7	7	-	-	-	0 mm	0
CC29/3	66	62,8	12	+	-	-	0 mm	0
CC30	68	17,1	24	-	+	-	16 mm	2 mm
CB31	69	0	5	+	-	-	-	0
CA32/1	70	0	35	-	-	-	2 mm	0
CA32/2	71	50	35	+	-	-	8 mm	0
CA33/1	72	48	7	+	-	-	4mm	0
CA33/2	73	44	14	-	+	-	6 mm	3 mm
CA34	74	20	5	+	-	-	1 mm	0
CB34	75	0	5	+	-	-	1mm	0
CC35/1	76	14	16	-	+	-	8 mm	0
CC35/2	77	43	7	+	-	-	2 mm	0
CC35/3	78	0	8	-	-	-	12 mm	0
CA36	79	14	8	+	-	-	1 mm	0
CB36/1	80	43	125	-	-	-	7 mm	4 mm
CB36/2	81	25	20	+	-	-	10 mm	0
CC37/1	82	22,2	5	+	-	-	3 mm	0
CC37/2	83	15	45	-	-	-	7 mm	6 mm
CC37/3	84	42	15	+	-	-	5 mm	0
CA38	85	63	5	+	-	-	6 mm	0
CB38/1	86	0	6	+	-	-	4 mm	0
CB38/2	87	31	50	-	-	-	6 mm	1 mm
CA39/1	88	0	12	-	-	-	12 mm	0
CA39/2	89	32,6	13	+	-	-	4 mm	0
CB39	90	33,3	12	+	-	-	1 mm	0
CC39 /1	91	11,1	8	+	-	-	3 mm	0
CC39 /2	92	0	15	+	-	-	1 mm	0
CC39 /3	93	42	39	+	-	-	1 mm	0
C40	94	31,5	12	+	-	-	3 mm	0 mm
CB40 /1	95	16,4	11	-	+	-	11 mm	3 mm
CB40 /2	96	24,6	18	-	+	-	19 mm	0
CC40 /1	97	10,9	5	-	-	-	2 mm	0
CC40 /2	98	23,2	32	-	-	-	4 mm	1,5 mm
CA41 /1	99	17,8	6	+	-	-	1 mm	0
CA41 /2	100	9,5	11	-	-	-	2 mm	0
CA41 /3	101	17,8	9	+	-	-	2 mm	0
CC41 /1	102	26	8	+	-	-	0 mm	0
CC41 /2	103	19,1	41	-	-	-	3 mm	3,5 mm
CA42	104	10,9	8	-	-	-	0 mm	0
CB42 /1	105	17,8	26	+	-	-	1 mm	0
CB42 /2	106	17,8	6	+	-	-	0 mm	0
CC42 /1	107	20,5	41	-	-	-	1 mm	5 mm
CC42 /2	108	30,5	15	-	-	-	1 mm	1 mm
CC43	109	24,6	7	+	-	-	0 mm	1 mm
CA44 /1	110	34,2	6	+	-	-	0 mm	0
CA44 /2	111	26	8	-	-	-	1 mm	0
CC44	112	31,5	8	-	+	-	6 mm	1.5 mm

¹The values show the percent inhibition of mycelial development of FOC compared to non treated positive control plates of FOC

² The values are the mean of four replicate plates

³Auxin level after growth in medium supplemented with/without tryptophan

⁴Hydrogen cyanide production

⁵Mean vigor index value of non treated negative control was 4947




Figure 1: Vigority Index (VI) of germinated cucumber seeds coated with EB strains.

giberellins and cytokinins. Auxins are known to be essential for plant physiology directly affecting the root and shoot architecture. More than 30 % of the EB strains produced detectable levels (20–125 $\mu\text{g ml}^{-1}$) of IAA in culture filtrates (Table 1). IAA production was highest in the *Pseudomonas* followed by *Bacillus* isolates. Plant growth promotion mediated by endophytic bacteria may be exerted by several mechanisms, e.g. synthesis of siderophores, solubilisation of minerals such as phosphorous [12,13,35]. Siderophore production was exhibited by 46 % strains ranging from 3 to 19 mm zones on CAS agar (Table 1). Solubilization of phosphate was detected in 29 % of the EB strains ranging from 1 to 9 mm zones (Table 1). Most EB strains were HCN negative on TSA, with or without glycine. Only three *Pseudomonas* isolates showed detectable cyanide production by changing color from yellow to brown around its colonies. In this study, we showed that EB strains were very promising in respect to *in vitro* plant growth promotion parameters.

Effect on fungal growth

A total of 112 strains of EB were tested for their *in vitro* antagonistic activity against FOC on PDA plates. 53% of the EB strains showed antagonism against FOC on PDA plates producing inhibition zones by dual plate test (Table 1). The inhibitory rates varied from 20% to 64% depending on EB strains (Table 1). Most of the EB isolates, which were HCN negative on TSA showed antibiosis against FOC *in vitro*. Endophytic bacteria isolated from potato roots expressed high levels of hydrolytic enzymes such as cellulase, chitinase and glucanase [36]. Our results showed that the EB strains were effective against FOC and produced inhibitory metabolites other than hydrogen cyanide.

Effect on seed germination

Thirty eight isolates out of 112 EB, considered as successful for *in vitro* plant growth promoting traits and biocontrol activities toward FOC were analyzed for their effects on seed emergence on cucumber seeds and VI. Some of the EB strains had no apparent effect on seed germination and VI, where as others, when applied individually, caused suppression of seed germination *in vitro* and seedling growth compared to that of the control plates (Table 1). For example, among 38 EB strains, which were used in plate assay, strains CC37/2 and then CB38/2 were the best ones on enhancement of VI, while strains CA29/1, CA28/2, CB36/2 and Ca38 did not have any effect

on seed germination or VI (Figure 1). Consistent with our results, some bacterial endophytic isolates from healthy plants inhibited the growth of tomato seedlings in reinoculation assays, possibly through the production of certain metabolites [37]. In plate assay, CA33/2 and CB8/1 strains had the lowest values of VI (Figure 1). We observed that 40% of tested EB strains improved the VI of cucumber seedlings compared to that treated with CMC (1% w/v) only (Figure 1). Approximately 54 % of tested EB isolates had a strong potential for promoting seed germination and VI comparing to control plates. [19] described that the isolates of endophytic bacteria significantly improved seed germination and plant growth of oilseed rape and tomato.

Conclusion

In conclusion, EB can be isolated by surface-sterilized method. A total of 112 EB strains were recovered from cucumber plants. Present study showed high activity of EB strains against FOC. The strains CC29/3 and CC25/2 were more active compared to other strains against FOC. Furthermore, majority of isolated endophytes were not only able to suppress pathogenic fungi, but could also improve seed germination and plant growth. These strains may be candidates for biological control and plant growth promotion.

Acknowledgement

This study was supported by national grants from the Scientific and Technological Research Council of Turkey (TUBITAK-COST 111 O 505) and Ege University Science-Technology Research and Application Center (EBILTEM). The authors thank the contributions of the participants in the EU-COST Action FA1103.

References

- Owen JH. Fusarium wilt of cucumber. *Phytopathology*. 1955; 45: 435–439.
- Owen JH. Cucumber wilt, caused by *Fusarium Oxysporum* f. sp. *Cucumerinum* N.F. *Phytopathology*. 1956; 46: 153–157.
- Martyn RD. Fusarium wilt of cucumber. TA Zitter DL, Hopkins, CE Thomas, editors. In: *Compendium of cucurbit diseases*. St. Paul: The American Phytopathological Society Press. 1996.
- Wicks TJ, Volle D and Baker BT. The effect of soil fumigation and fowl manure on populations of *Fusarium oxysporum* f. sp. *cucumerinum* in greenhouse soil and on the incidence of cucumber wilt. *Agricultural Record*. 1978; 5: 4-8.

5. Jenkins SF, Wehner TC. Occurrence of *Fusarium oxysporum* f. sp. *cucumerinum* on greenhouse-grown *Cucumis sativus* seed stocks in North Carolina. *Plant Disease*. 1983; 67: 1024–1025.
6. Martínez R, Aguilar MI, Guirado ML, Álvarez A, Gómez J. First report of fusarium wilt of cucumber caused by *Fusarium oxysporum* in Spain. *Plant Pathology*. 2003; 52: 410.
7. Neshev G. Major soil-borne phytopathogens on tomato and cucumber in Bulgaria, and methods for their management. Rome: Food and Agriculture Organization of the United Nations (FAO). 2008: 1-22.
8. Song W, Zhou L, Yang C, Cao X, Zhang L, Liu Z. Tomato *Fusarium* wilt and its chemical control strategies in a hydroponic system. *Crop Protection*. 2004; 23: 243-247.
9. Lemanceau P, Alabouvette C. Suppression of *Fusarium* wilts by fluorescent *pseudomonads*: mechanisms and applications. *Biocontrol Science and Technology*. 1993; 3: 219-34.
10. Spadaro D and Gullino ML. Improving the efficacy of biocontrol agents against soilborne pathogens. *Crop Protection*. 2005; 24: 601-613.
11. Altın N, Bora T. Research on the biological control of *Fusarium* wilt of cucumber (*Fusarium oxysporum* f.sp. *cucumerinum*) in greenhouse PhD. Thesis in Plant protection. 2004; 111.
12. Hurek T, Reinhold-Hurek B. *Azoarcus* sp. strain BH72 as a model for nitrogen-fixing grass endophytes. *J Biotechnol*. 2003; 106: 169-178.
13. Whipps JM. Microbial interactions and biocontrol in the rhizosphere. *J Exp Bot*. 2001; 52: 487-511.
14. Ryan RP, Germaine K, Franks A, Ryan DJ, Dowling DN. Bacterial endophytes: recent developments and applications. *FEMS Microbiol Lett*. 2008; 278: 1-9.
15. Rajendran G, Sing F, Desai AJ, Archana G. Enhanced growth and nodulation of pigeon pea by co-inoculation of *Bacillus* strains with *Rhizobium* spp. See comment in PubMed Commons below *Bioresour Technol*. 2008; 99: 4544-4550.
16. Zakhia F, Jeder H, Willems A, Gillis M, Dreyfus B, de Lajudie P. Diverse bacteria associated with root nodules of spontaneous legumes in Tunisia and first report for *nifH*-like gene within the genera *Microbacterium* and *Starkeya*. *Microb Ecol*. 2006; 51: 375-393.
17. Kan FL, Chen ZY, Wang ET, Tian CF, Sui XH, Chen WX. Characterization of symbiotic and endophytic bacteria isolated from root nodules of herbaceous legumes grown in Qinghai-Tibet Plateau and in other zones of China. *Arch Microbiol*. 2007; 188: 103-115.
18. Li JH, Wang ET, Chen WF, Chen WX. Genetic diversity and potential for promotion of plant growth detected in nodule endophytic bacteria of soybean grown in Heilongjiang province of China. *Soil Biol Biochem*. 2008; 40: 238-246.
19. Nejad P, Johnson PA. Endophytic bacteria induce growth promotion and wilt disease suppression in oilseed rape and tomato. *Biol Control*. 2000; 18: 208–215.
20. Suslow TV, Schroth MN, Isaka M. Application of a rapid method for Gram differentiation of plant pathogenic and saprophytic bacteria without staining. *Phytopathology*. 1982; 72: 917–918.
21. Lelliot RA, Stead DE. *Methods for Diagnosis of Bacterial Diseases of Plants*. Blackwell. UK. 1987.
22. Schaad NW, Jones JB, Chun W. *Laboratory guide for identification of plant pathogenic bacteria*. Third Edition, APS: New York. 2001.
23. Trivedi P, Pandey A, Palni LM. *In vitro* evaluation of antagonistic properties of *Pseudomonas corrugata*. *Microbiol Res*. 2008; 163: 329-336.
24. Bakker AW, Schippers B. Microbial cyanide production in the rhizosphere in relation to potato yield reduction and *Pseudomonas* spp-mediated plant growth – stimulation. *Soil Biol. Biochem*. 1987; 19: 451-457.
25. Pikovskaya RE. Mobilization of phosphorus in soil in connection with vital activity of some microbial species. *Microbiologia*. 1948; 17: 362-370.
26. Bric JM, Bostock RM, Silverstone SE. Rapid in situ assay for indoleacetic Acid production by bacteria immobilized on a nitrocellulose membrane. *Appl Environ Microbiol*. 1991; 57: 535-538.
27. Schwyn B, Neilands JB. Universal chemical assay for the detection and determination of siderophores. *Anal Biochem*. 1987; 160: 47-56.
28. Abdul-Baki A, Anderson JD. Vigor determination in Soybean seed by multiple criteria. *Crop Sci*. 1973; 13: 630-633.
29. Bardas GA, Lagopodi A, Kadoglidou K, Tzavella-Klonari K. Biological control of three *Colletotrichum lindemuthianum* races using *Pseudomonas chlororaphis* PCL1391 and *Pseudomonas fluorescens* WCS365. *Biol Control*. 2009; 49: 139–145.
30. Jousset A, Rochat L, Scheu S, Bonkowski M, Keel C. Predator-prey chemical warfare determines the expression of biocontrol genes by rhizosphere-associated *Pseudomonas fluorescens*. *Appl Environ Microbiol*. 2010; 76: 5263-5268.
31. Vallet-Gely I, Novikov A, Augusto L, Liehl P, Bolbach G, Péchy-Tarr M, et al. Association of hemolytic activity of *Pseudomonas entomophila*, a versatile soil bacterium, with cyclic lipopeptide production. *Appl Environ Microbiol*. 2010; 76: 910–921.
32. Wensing AD, Braun S, Büttner P, Expert D, Völksch B, S Ullrich M, et al. Impact of siderophore production by *Pseudomonas syringae* pv. *syringae* 22d/93 on epiphytic fitness and biocontrol activity against *Pseudomonas syringae* pv. *glycinea* 1a/96. *Appl Environ Microbiol*. 2010; 76: 2704–2711.
33. Matilla MA, Ramos JL, Bakker PA, Doornbos R, Badri DV, Vivanco JM, et al. *Pseudomonas putida* KT2440 causes induced systemic resistance and changes in *Arabidopsis* root exudation. See comment in PubMed Commons below *Environ Microbiol Rep*. 2010; 2: 381-388.
34. Hallmann J, Quadt-Hallmann A, Mahaffee WF, Kloepper JW. Bacterial endophytes in agricultural crops. *Can. J. Microbiol*. 1997; 43: 895–914.
35. Cherrin L, Chet I. Microbial enzymes in biocontrol of plant pathogens and pests. Burns RG, Dick RP, Marcel Dekker, editors. In: *Enzymes in the environment: activity, ecology, and applications*. New York. 2002: 171–225.
36. Krechel A, Faupel A, Hallmann J, Ulrich A, Berg G. Potato-associated bacteria and their antagonistic potential towards plant-pathogenic fungi and the plant-parasitic nematode *Meloidogyne incognita* (Kofoid & White) Chitwood. *Can J Microbiol*. 2002; 48: 772-786.
37. van Peer R, Punte HL, de Weger LA, Schippers B. Characterization of Root Surface and Endorhizosphere *Pseudomonads* in Relation to Their Colonization of Roots. *Appl Environ Microbiol*. 1990; 56: 2462-2470.