

Research Article

# Comorbidity between Obsessive-Compulsive Disorder and Schizophrenia: Is there Evidence for a “Schizo-obsessive” Subtype?

Frias A<sup>1,2\*</sup>, Palma C<sup>1,2</sup> and Farriols N<sup>1,2</sup>

<sup>1</sup>RamonLlull University, Spain

<sup>2</sup>Adult Outpatient Mental Health Center, Spain

\*Corresponding author: Frias A, Adult Outpatient Mental Health Center, FPCEE Blanquerna, Cister 34, 08022, Barcelona, Spain

Received: June 28, 2014; Accepted: July 23, 2014;

Published: July 25, 2014

## Abstract

This systematic review displays main findings and research regarding the nosological status of patients diagnosed with OCD and schizophrenia. Particularly, results from psychopathology, psychosocial functioning, neurocognition and treatment are arranged. On the one hand, results obtained indicate that “schizo-obsessive” patients tend to show higher depressed mood, negative symptomatology and psychosocial impairment when compared with “schizophrenia” subjects. On the other hand, no consistent differences between both psychotic groups have been found in positive symptomatology, neurocognitive performance and treatment features. Of interest, some methodological constraints hamper generalization of research findings, especially type of “schizo-obsessive” subjects recruited (OCD vs. OCS comorbidity). Overall, results collected so far do not support that “schizo-obsessive” pathology represents an independent clinical entity.

**Keywords:** Schizo-obsessive disorder; Schizophrenia; Obsessive-compulsive disorder; Comorbidity, Theoretical study

## Abbreviations

ARMS: At-risk Mental State; CPT: Continuous Performance Test; CVLT: California Verbal Learning Test; OCD: Obsessive-compulsive Disorder; OCS: Obsessive-Compulsive Symptoms; TMT: Trail Making Test; WCST: Wisconsin Card Sorting Test; WMS: Wechsler Memory

## Introduction

Despite having been a research topic since the 19<sup>th</sup> century, the relationship between schizophrenia and OCD has only attained increasing recognition in the last 15 years [1-3]. Epidemiological studies have found that OCS are estimated to occur in up to 30% of patients with schizophrenia [4,5]. Likewise, OCD affects roughly 12% of this group, depending on the type of sample (clinical vs. community), setting (outpatient vs. inpatient), design study (cross-sectional vs. prospective), or assessment instruments (clinical interview vs. questionnaires) [6]. Overall, these prevalences are over 5 times higher than expected in subjects without schizophrenia, raising concerns regarding its etiopathogenic mechanisms and nosological status [7].

To date, several explanations have been suggested as ways of understanding this co-occurrence. First, OCD/OCS may be part of early stage of psychotic illness, specifically during the ARMS [2,8,9]. Second, one disorder might cause or increase the risk for the other [2,10,11]. Third, both disorders may share common or underlying risk factors such as neuropsychological and biological domains [12-15]. Finally, some current research postulates that the development of OCS in schizophrenia may be associated with atypical antipsychotic effects [16-18]. Irrespective of these possible theories, whether this

subgroup of “schizo-obsessive” patients may pose as a clinical entity with a distinct profile presents a long-standing trouble [19-23]. Accordingly, there is still a lack of recognition of “schizo-obsessive disorder” within current diagnosis reference manuals.

## Objective

Within this frame work, these systematic reviews aimed at delineating the clinical profile of “schizo-obsessive” patients and compare them with subjects diagnosed of “schizophrenia” or “OCD” alone. Based on the results obtained, we will elucidate whether subjects with this comorbidity may be posed as a distinct nosological category.

## Method

### Search strategy

A literature search was carried out through PsycINFO and PubMed databases from 1990 to June2014. Terms employed included indexing terms (e.g. MeSH) and free texts: [(OCDOR obsessive-compulsive) and (schizophrenia OR psychosis)].

### Selection criteria

Inclusion criteria included studies comprising subjects diagnosed of OCD and schizophrenia according to DSM-IV criteria, which were assessed for outcome measures related to this comorbidity. Overall, 60 studies fulfilled inclusion criteria and were finally collected, including both original articles and systematic reviews.

### Data extraction

Comparative data from socio demographic variables, psychopathology, quality of life and clinical variables, neuro cognitive profile and psycho pharmacological treatment were collected.

## Results

### Socio demographic variables

Concerning the socio demographic features, the “schizo-obsessive” subjects have displayed a lower degree of education, lesser rates of employment, and living fewer times with a partner than “schizophrenia” subjects [22,24,25].

### Psychopathological variables

Concerning the psychotic symptomatology and insight into psychosis, studies usually have found higher negative symptomatology among “schizo-obsessive” with OCD patients than “schizophrenia” subjects [4,20,26-30]. Conversely, those studies which collected “schizo-obsessive” subjects with OCS have not shown between-group differences [21,31-34]. Despite these previous results, to date those studies that have already included direct comparative analyses between “schizo-obsessive” patients with OCD vs. OCS have not yet obtained consistent evidences regarding more negative among the “schizo-obsessive” subjects with OCD instead of OCS [18,22,35]. With regard to the obsessive-compulsive features, most studies have not found differences between “schizo-obsessive” patients and “OCD” subjects respecting the severity of the symptomatology [36-39]. Concerning insight into OCD, studies have not yielded consistent findings [36,38,40].

Regarding the depressed mood, most research has obtained higher depressed mood in the “schizo-obsessive” group than in “schizophrenia” and/or “OCD” patients [4,20,25,35,38,41].

Finally, studies have not found statistically significant differences among the 3 groups concerning the global severity of the whole psychopathology [27,42,43].

### Quality of life and clinical variables

Regarding the quality of life, most studies have pointed out a lower quality of life among the “schizo-obsessive” subjects than in “schizophrenia” patients [4,32,38,44-47].

In addition, studies which have assessed clinical variables such as “number/length of hospitalizations”, “psychiatric emergencies”, and “previous suicide attempts” among the both psychotic groups have yielded inconsistent results. On the one hand, some studies have displayed a lack of differences in number of hospitalizations and psychiatric emergencies [4,25,38]. On the other hand, other inquiries have shown greater number/length of hospitalizations and psychiatric emergencies among “schizo-obsessive” patients [11,24,32]. In addition, “schizo-obsessive” subjects usually have committed statistically more suicide attempts than “schizophrenia” patients [4,48].

### Neuropsychological profile

Regarding the neuropsychological results, most previous studies that have included an “OCD” group have obtained a better performance in this group when compared with both psychotic groups [46,49,50]. Bearing in mind both psychotic groups, results have been inconsistent, depending on the neuro cognitive domain as well as the tests employed. Findings are described below.

Concerning visual spatial skills, both psychotic groups have shown either no between-group differences [47,49-51] or higher

performance among “schizo-obsessive” subjects [2,52]. Regarding working memory, studies have displayed a lack of statistically significant differences between both psychotic groups using the TMT [47,50,53]. However, other studies that assessed this domain using the WCST or the Stroop test yielded divergent findings [32,33,46,49,51,52,54]. With regard to processing speed, there was a lack of statistically significant differences between both psychotic groups [47,50,52,53], with the exception of some studies [51,54]. Concerning cognitive shifting abilities, most studies have yielded no between-group differences using the TMT [47,50,53], with the exception of one study [54]. Not with standing, studies which used other neuropsychological tests (e.g., alternation learning tasks, CPT, go-no go test and WCST) have obtained inconsistent results [32,33,46,49,51,55-57].

Regarding concentration, studies using the digit span subtest have not found statistically significant differences between both psychotic groups [50]. Conversely, research using the Stroop test has not yielded convergent results [47,50,51]. Concerning verbal memory, studies using the FAS have not yielded between-group differences [47,52]. Similarly, research with the same purpose using the WMS or the CVLT have yielded similar results [46,50,52].

### Psychopharmacological treatment

Concerning dosage and type of antipsychotic medications, studies that have recorded chlorpromazine equivalent doses [32,52,58] as well as the type of antipsychotics (typical vs. atypical)[22,51] have not displayed differences between both psychotic groups. These findings suggest that antipsychotic medication may not play a role on the psychopathological phenotype of “schizo-obsessive” subjects. Thus, potentially contaminating effects from this variable may be ruled out when interpreted previous findings.

## Conclusion

Findings from this systematic review do not overtly support that “schizo-obsessive” subjects were a distinct clinical profile when compared with “schizophrenia” patients. On the one hand, “schizo-obsessive” subjects seem to display higher depressed mood, negative symptomatology and psychosocial dysfunction. However, some methodological constraints affect results, especially type of “schizo-obsessive” subjects (with OCD or with OCS). On the other hand, neuropsychological, treatment and clinical variables have provided inconsistent results [59]. Likewise, some methodological flaws have partially determined these last findings, especially type of neuropsychological tests employed [60]. Overall, we cannot ensure that “schizo-obsessive” pathology represents an independent clinical entity. Further research is warranted for addressing this unresolved topic. By resolving these methodological constraints, findings from next studies may provide reliable evidence that could shed further light on the nosological status of “schizo-obsessive” patients.

## References

1. Poyurovsky M, Zohar J, Glick I, Koran LM, Weizman R, Tandon R, et al. Obsessive-compulsive symptoms in schizophrenia: implications for future psychiatric classifications. *Compr Psychiatry*. 2012; 53: 480-483.
2. Schirmbeck F, Zink M. Comorbid obsessive-compulsive symptoms in schizophrenia: contributions of pharmacological and genetic factors. *Front Pharmacol*. 2013; 4: 99.

3. Frías-Ibáñez A, Palma-Sevillano C, Farriols-Hernando N. Comorbidity between obsessive-compulsive disorder and schizophrenia: prevalence, explanatory theories, and nosological status. *Actas Esp Psiquiatr*. 2014; 42: 28-38.
4. de Haan L, Sterk B, Wouters L, Linszen DH. The 5-year course of obsessive-compulsive symptoms and obsessive-compulsive disorder in first-episode schizophrenia and related disorders. *Schizophr Bull*. 2013; 39: 151-160.
5. Swets M, Dekker J, van Emmerik-van Oortmerssen K, Smid GE, Smit F, de Haan L, et al. The obsessive compulsive spectrum in schizophrenia, a meta-analysis and meta-regression exploring prevalence rates. *Schizophr Res*. 2014; 152: 458-468.
6. Poyurovsky M, Weizman A, Weizman R. Obsessive-compulsive disorder in schizophrenia: clinical characteristics and treatment. *CNS Drugs*. 2004; 18: 989-1010.
7. Crino R, Slade T, Andrews G. The changing prevalence and severity of obsessive-compulsive disorder criteria from DSM-III to DSM-IV. *Am J Psychiatry*. 2005; 162: 876-882.
8. Fontenelle LF, Lin A, Pantelis C, Wood SJ, Nelson B, Yung AR. A longitudinal study of obsessive-compulsive disorder in individuals at ultra-high risk for psychosis. *J Psychiatr Res*. 2011; 45: 1140-1145.
9. Sterk B, Lankreijer K, Linszen DH, de Haan L. Obsessive-compulsive symptoms in first episode psychosis and in subjects at ultra high risk for developing psychosis; onset and relationship to psychotic symptoms. *Aust N Z J Psychiatry*. 2011; 45: 400-406.
10. Niendam TA, Berzak J, Cannon TD, Bearden CE. Obsessive compulsive symptoms in the psychosis prodrome: correlates of clinical and functional outcome. *Schizophr Res*. 2009; 108: 170-175.
11. Üçok A, Ceylan ME, Tihan AK, Lapçin S, Ger C, Tükel R. Obsessive compulsive disorder and symptoms may have different effects on schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; 35: 429-433.
12. Abbruzzese M, Bellodi L, Ferri S, Scarone S. Frontal lobe dysfunction in schizophrenia and obsessive-compulsive disorder: a neuropsychological study. *Brain Cogn*. 1995; 27: 202-212.
13. Kim MS, Kang SS, Youn T, Kang DH, Kim JJ, Kwon JS. Neuropsychological correlates of P300 abnormalities in patients with schizophrenia and obsessive-compulsive disorder. *Psychiatry Res*. 2003; 123: 109-123.
14. Kwon JS, Shin YW, Kim CW, Kim Y, Youn T, Han MH, et al. Similarity and disparity of obsessive-compulsive disorder and schizophrenia in MR volumetric abnormalities of the hippocampus-amygdala complex. *J Neurol Neurosurg Psychiatry*. 2003; 74: 962-964.
15. Martin V, Huber M, Rief W, Exner C. Comparative cognitive profiles of obsessive-compulsive disorder and schizophrenia. *Arch Clin Neuropsychol*. 2008; 23: 487-500.
16. Lin SK, Su SF, Pan CH. Higher plasma drug concentration in clozapine-treated schizophrenic patients with side effects of obsessive/compulsive symptoms. *Ther Drug Monit*. 2006; 28: 303-307.
17. Mahendran R, Liew E, Subramaniam M. De novo emergence of obsessive-compulsive symptoms with atypical antipsychotics in asian patients with schizophrenia or schizoaffective disorder: a retrospective, cross-sectional study. *J Clin Psychiatry*. 2007; 68: 542-545.
18. Sa AR, Hounie AG, Sampaio AS, Arrais J, Miguel EC, Elkis H. Obsessive-compulsive symptoms and disorder in patients with schizophrenia treated with clozapine or haloperidol. *Compr Psychiatry*. 2009; 50: 437-442.
19. Zohar J. Is there room for a new diagnostic subtype: the schizo-obsessive subtype? *CNS Spectr*. 1997; 2: 49-50.
20. Rajkumar RP, Reddy YC, Kandavel T. Clinical profile of "schizo-obsessive" disorder: a comparative study. *Compr Psychiatry*. 2008; 49: 262-268.
21. Cunill R, Castells X, Simeon D. Relationships between obsessive-compulsive symptomatology and severity of psychosis in schizophrenia: a systematic review and meta-analysis. *J Clin Psychiatry*. 2009; 70: 70-82.
22. Owashi T, Ota A, Otsubo T, Susa Y, Kamijima K. Obsessive-compulsive disorder and obsessive-compulsive symptoms in Japanese inpatients with chronic schizophrenia - a possible schizophrenic subtype. *Psychiatry Res*. 2010; 179: 241-246.
23. Docherty AR, Coleman MJ, Tu X, Deutsch CK, Mendell NR, Levy DL. Comparison of putative intermediate phenotypes in schizophrenia patients with and without obsessive-compulsive disorder: examining evidence for the schizo-obsessive subtype. *Schizophr Res*. 2012; 140: 83-86.
24. Berman I, Kalinowski A, Berman SM, Lengua J, Green AI. Obsessive and compulsive symptoms in chronic schizophrenia. *Compr Psychiatry*. 1995; 36: 6-10.
25. Seedat F, Roos JL, Pretorius HW, Karayiorgou M, Nel B. Prevalence and clinical characteristics of obsessive-compulsive disorder and obsessive compulsive symptoms in Afrikaner schizophrenia and schizoaffective disorder patients. *Afr J Psychiatry*. 2007; 10: 219-224.
26. Krüger S, Bräunig P, Höffler J, Shugar G, Börner I, Langkrär J. Prevalence of obsessive-compulsive disorder in schizophrenia and significance of motor symptoms. *J Neuropsychiatry Clin Neurosci*. 2000; 12: 16-24.
27. Sevincok L, Akoglu A, Topaloglu B, Aslantas H. Neurological soft signs in schizophrenic patients with obsessive-compulsive disorder. *Psychiatry Clin Neurosci*. 2004; 58: 274-279.
28. Byerly M, Goodman W, Acholonu W, Bugno R, Rush AJ. Obsessive compulsive symptoms in schizophrenia: frequency and clinical features. *Schizophr Res*. 2005; 76: 309-316.
29. Faragian S, Kurs R, Poyurovsky M. Insight into obsessive-compulsive symptoms and awareness of illness in adolescent schizophrenia patients with and without OCD. *Child Psychiatry Hum Dev*. 2008; 39: 39-48.
30. Poyurovsky M, Faragian S, Shabeta A, Kosov A. Comparison of clinical characteristics, co-morbidity and pharmacotherapy in adolescent schizophrenia patients with and without obsessive-compulsive disorder. *Psychiatry Res*. 2008; 159: 133-139.
31. Aoyama F, Iida J, Inoue M, Iwasaka H, Sakiyama S, Hata K, et al. Brain imaging in childhood- and adolescence-onset schizophrenia associated with obsessive-compulsive symptoms. *Acta Psychiatr Scand*. 2000; 102: 32-37.
32. Hwang MY, Opler LA. Management of schizophrenia with obsessive compulsive disorder. *Psychiatry Ann*. 2000; 30: 23-28.
33. Lysaker PH, Bryson GJ, Marks KA, Greig TC, Bell MD. Association of obsessions and compulsions in schizophrenia with neurocognition and negative symptoms. *J Neuropsychiatry Clin Neurosci*. 2002; 14: 449-453.
34. Nechmad A, Ratzoni G, Poyurovsky M, Meged S, Avidan G, Fuchs C, et al. Obsessive-compulsive disorder in adolescent schizophrenia patients. *Am J Psychiatry*. 2003; 160: 1002-1004.
35. de Haan L, Hoogenboom B, Beuk N, van Amelsvoort T, Linszen D. Obsessive-compulsive symptoms and positive, negative, and depressive symptoms in patients with recent-onset schizophrenic disorders. *Can J Psychiatry*. 2005; 50: 519-524.
36. Ozdemir O, Tükel R, Türksöy N, Uçok A. Clinical characteristics in obsessive-compulsive disorder with schizophrenia. *Compr Psychiatry*. 2003; 44: 311-316.
37. Poyurovsky M, Koran LM. Obsessive-compulsive disorder (OCD) with schizotypy vs. schizophrenia with OCD: diagnostic dilemmas and therapeutic implications. *J Psychiatr Res*. 2005; 39: 399-408.
38. Güleç G, Güneş E, Yenilmez C. Comparison of patients with schizophrenia, obsessive-compulsive disorder, and schizophrenia with accompanying obsessive-compulsive symptoms. *Türk Psikiyatri Derg*. 2008; 19: 247-256.
39. de Haan L, Dudek-Hodge C, Verhoeven Y, Denys D. Prevalence of psychotic disorders in patients with obsessive-compulsive disorder. *CNS Spectr*. 2009; 14: 415-417.
40. Matsunaga H, Kiriike N, Matsui T, Oya K, Iwasaki Y, Koshimune K, et al. Obsessive-compulsive disorder with poor insight. *Compr Psychiatry*. 2002; 43: 150-157.
41. Fawzi MH, Fawzi MM, Khedr HH, Fawzi MM. Tobacco smoking in Egyptian

- schizophrenia patients with and without obsessive-compulsive symptoms. *Schizophr Res.* 2007; 95: 236-246.
42. Poyurovsky M, Faragian S, Pashinian A, Levi A, Viosburd A, Stryjer R, et al. Neurological soft signs in schizophrenia patients with obsessive-compulsive disorder. *J Neuropsychiatry Clin Neurosci.* 2007; 19: 145-150.
43. Thomas N, Tharyan P. Soft neurological signs in drug-free people with schizophrenia with and without obsessive-compulsive symptoms. *J Neuropsychiatry Clin Neurosci.* 2011; 23: 68-73.
44. Poyurovsky M, Hramenkov S, Isakov V, Rauchverger B, Modai I, Schneidman M, et al. Obsessive-compulsive disorder in hospitalized patients with chronic schizophrenia. *Psychiatry Res.* 2001; 102: 49-57.
45. Lysaker PH, Lancaster RS, Nees MA, Davis LW. Patterns of obsessive-compulsive symptoms and social function in schizophrenia. *Psychiatry Res.* 2004; 125: 139-146.
46. Whitney KA, Fastenau PS, Evans JD, Lysaker PH. Comparative neuropsychological function in obsessive-compulsive disorder and schizophrenia with and without obsessive-compulsive symptoms. *Schizophr Res.* 2004; 69: 75-83.
47. Tiryaki A, Ozkorumak E. Do the obsessive-compulsive symptoms have an effect in schizophrenia? *Compr Psychiatry.* 2010; 51: 357-362.
48. Ongür D, Goff DC. Obsessive-compulsive symptoms in schizophrenia: associated clinical features, cognitive function and medication status. *Schizophr Res.* 2005; 75: 349-362.
49. Borkowska A, Pilaczyńska E, Rybakowski JK. The frontal lobe neuropsychological tests in patients with schizophrenia and/or obsessive-compulsive disorder. *J Neuropsychiatry Clin Neurosci.* 2003; 15: 359-362.
50. Tumkaya S, Karadag F, Oguzhanoglu NK, Tekkanat C, Varma G, Ozdel O, et al. Schizophrenia with obsessive-compulsive disorder and obsessive-compulsive disorder with poor insight: a neuropsychological comparison. *Psychiatry Res.* 2009; 165: 38-46.
51. Michalopoulou PG, Konstantakopoulos G, Typaldou M, Papageorgiou C, Christodoulou GN, Lykouras L, et al. Can cognitive deficits differentiate between schizophrenia with and without obsessive-compulsive symptoms? *Compr Psychiatry.* 2014; 55: 1015-1021.
52. Lee MJ, Shin YB, Sunwoo YK, Jung SH, Kim WH, Kang MH, et al. Comparative Analysis of Cognitive Function in Schizophrenia with and without Obsessive Compulsive Disorder. *Psychiatry Investig.* 2009; 6: 286-293.
53. Berman I, Merson A, Viegner B, Losonczy MF, Pappas D, Green AI. Obsessions and compulsions as a distinct cluster of symptoms in schizophrenia: a neuropsychological study. *J Nerv Ment Dis.* 1998; 186: 150-156.
54. Schirmbeck F, Rausch F, Englisch S, Eifler S, Esslinger C, Meyer-Lindenberg A, et al. Stable cognitive deficits in schizophrenia patients with comorbid obsessive-compulsive symptoms: a 12-month longitudinal study. *Schizophr Bull.* 2013; 39: 1261-1271.
55. Lysaker PH, Whitney KA, Davis LW. Associations of executive function with concurrent and prospective reports of obsessive-compulsive symptoms in schizophrenia. *J Neuropsychiatry Clin Neurosci.* 2009; 21: 38-42.
56. Patel DD, Laws KR, Padhi A, Farrow JM, Mukhopadhyaya K, Krishnaiah R, et al. The neuropsychology of the schizo-obsessive subtype of schizophrenia: a new analysis. *Psychol Med.* 2010; 40: 921-933.
57. Hermesh H, Weizman A, Gur S, Zalsman G, Shiloh R, Zohar J, et al. Alternation learning in OCD/schizophrenia patients. *Eur Neuropsychopharmacol.* 2003; 13: 87-91.
58. Ohta M, Kokai M, Morita Y. Features of obsessive-compulsive disorder in patients primarily diagnosed with schizophrenia. *Psychiatry Clin Neurosci.* 2003; 57: 67-74.
59. Meijer JH, Swets M, Keeman S, Nieman DH, Meijer CJ; GROUP investigators. Is a schizo-obsessive subtype associated with cognitive impairment? Results from a large cross-sectional study in patients with psychosis and their unaffected relatives. *J Nerv Ment Dis.* 2013; 201: 30-35.
60. Cunill R, Huerta-Ramos E, Castells X. The effect of obsessive-compulsive symptomatology on executive functions in schizophrenia: a systematic review and meta-analysis. *Psychiatry Res.* 2013; 210: 21-28.