

Special Article - Dream Anxiety Disorders

Nightmare Utilization by Successful Artists: A Case Report Series

Page JF*

Department of Psychiatry, University of Colorado School of Medicine, USA

*Corresponding author: Page JF, Department of Psychiatry, University of Colorado School of Medicine, PO Box 6, Arroyo Seco, NM 87514, USA

Received: June 29, 2017; Accepted: July 20, 2017;

Published: July 27, 2017

Abstract

Many successful artists use their nightmares in their work. This series of case reports (N=14) is a series of in-depth interviews with such artists, including an assessment of nightmare use in their work, any experience of trauma that may have contributed to their nightmares, and an assessment as to any past and current symptoms of PTSD. While some of these artists were best classified diagnostically as having Nightmare Disorder, a majority (11/14) of these successful artists had histories of significant trauma, and met DSM-V diagnostic criteria for PTSD. There is some evidence that these individuals did their best creative work after their experiences of trauma. These findings suggest that for the artist, nightmare expression rather than suppression (the objective of most PTSD therapies) might be a reasonable therapeutic option.

Introduction

Artists ranging from Goya, to Fusili and Picasso have used their nightmares in their work [1]. In our Sundance filmmaker work with dream use in creativity, we discovered significantly elevated levels of dream and nightmare recall and use when compared to clinical sleep and medical practice groupings, and the working/professional film making groups [2]. Sleep lab subjects reporting levels of creative interest and/or creative process were also found to report a higher incidence of nightmares than those reporting no creative interests [3]. These studies suggested the possibility that the report of nightmares might be a signifier or a marker for interest and involvement in the process of creativity. In a study of non-dreamers (a sleep laboratory grouping (N = 17) reporting no dream or nightmare recall by history or after multiple lab awakenings) the one documentable behavioral difference between this group and a grouping with minimal dream recall was a lack of interest and involvement in creative process [4]. This current study includes a series of fourteen in-depth interviews with successful visual artists who do creative work based at least in part on their nightmares. The interviews were minimally directed in order to obtain information as to recurrent nightmare content, any description of associated trauma, and any reported use of nightmares in their work. Due to the small group size, this is primarily a descriptive study designed to obtain answers to the following for this sample: 1) age, gender, and trauma exposure; 2) Post-Traumatic Stress Disorder (PTSD) symptom incidence; 3) characteristic nightmare content; and 4) incidence for the successful use of nightmares in artistic production.

Nightmares and POST-TRAUMATIC STRESS DISORDER (PTSD)

More than 70% of individuals being evaluated in the sleep lab report having nightmares more than once a month [4-6] (Table 1). Unlike other dreams, nightmares can be consistently defined by content that features agonizing dread, a sense of oppression, and the conviction of helpless paralysis [7]. Today, this classic definition for the nightmare has been both expanded and contracted, specifying that a nightmare is a disturbing mental experience generally occurring

during REM sleep that often results in awakening [8].

The classic nightmare almost always occurs in REMS except in individuals diagnosed with PTSD [9]. Distressing and recurrent nightmares are the most commonly reported symptom of PTSD, a waxing and waning chronic illness occurring after the experience of major physical or psychological trauma that can affect both waking and sleep into extreme old age [10,11]. Cognitive-behavioral therapies and medications can help some patients with the distress associated with PTSD [12]. Medications or behavioral approaches such as imagery or lucid control can reduce the intrusion of nightmares, hopefully reducing the destructive personal and social effects of PTSD [13]. However, many individuals who experience the irreconcilable physical or psychological trauma producing PTSD will continue to have symptoms throughout their lives. Many individuals with PTSD will never return to the level of functioning that they achieved before their experience of trauma [14]. Years after trauma, PTSD persists as a major risk factor for attempted and completed suicides [15,16]. A much higher risk of suicide post-trauma is present even in individuals with sub-threshold symptoms for PTSD [17]. In the short-term treatment of PTSD, reported nightmare frequency is the easiest to evaluate treatment outcome measure [11]. The therapeutic success of current medication and cognitive-behavioral psychological therapies is most often addressed as based on their ability to suppress and




Figure 1: Function and quality of life effects induced by major physical and psychological trauma and potential outcomes [2].

Table 1: Comparisons between PTSD groups of trauma severity; functional impairment; and nightmare content including Central Image (CI) intensity, narrative length; and nightmare (NM) incorporation into best work (artistic use) [means (range)].

PTSD CRITERIA	TRAUMA (1=NO RECALL – 5 = EXTREME)	FUNCTIONAL IMPAIRMENT (1=NONE – 5 = HOSPITAL OR SUICIDE ATTEMPT)	CENTRAL IMAGE INTENSITY (1 = MINIMAL -5 = PRIMARY)	NIGHTMARE LENGTH (1= IMAGE ONLY – 5 = LENGTHLY FULL STORY)	ARTISTIC USE (1 = NONE – 5 = NM BASED)
FULL (#7)	4.5 (3-5)	4.3 (4-5)	2.27 (1-5)	3.1 (1-5)	3.4 (1-5)
PARTIAL (#4)	4.0 (3-5)	3.0 (2-5)	3.5 (2-5)	3.75 (2-5)	3.75 (2-5)
NONE (#3)	1.4 (1-2)	1.0	3.0 (2-5)	3.3 (1-5)	4.3 (3-5)

decrease the frequency and distress of the associated nightmares [18].

Freud suggested that the psychological origin of creativity is based in trauma [19]. Freud suggested that the creative process could be therapeutic. The artist; “possesses the mysterious power of shaping some particular material until it becomes a faithful image of his phantasy.” So that the traumatized artist can potentially achieve; “...honour, power, and the love of women.” [20]. Ernest Hartmann noted that the only personality characteristic clearly associated with a higher incidence of nightmares is the characteristic of “thin-borders.” Individuals with thin borders are more likely to invest themselves in the creative experience, to produce creative products, and to lead creative roles in society with those reporting frequent nightmares testing as unusually imaginative [21]. In creative writing classes, those students with the most bizarre and fantastic dreams are the most likely to produce superior work [22].

Methodology

This project was presented in part at the 2010 Faulkner Society’s Words and Music meeting, in 2011 and 2013 at the International Society for the Study of Dreams, and in 2012 at Creativity and Madness (Santa Fe). The artists included in this project came based on personal contacts (primarily through the Sundance Film Labs), attendees at the above programs who volunteered after the presentation, and referrals from other interviewees involved in this project. All are successful artists who have, at some point, derived primary income based on their artistic role. All, even those past retirement age, were currently working in their field at the time of interview (2010-2015) except for one who is currently active duty military. The interviews were conducted by a physician Board certified in Sleep and Behavioral Sleep Medicine, and were transcribed and returned to each interviewee for correction and signed release. Three of the interviewees chose to submit their interview under a pseudonym. Art works were submitted by eight of the participants. Examples of that work have been published in a previous article [23].

Each subject voluntarily provided sufficient information to determine whether they met criteria for PTSD based on the current DSM-V criteria: 1) A traumatic exposure must result from one or more of the following scenarios, in which the individual directly experiences the traumatic event, witnesses the traumatic event in person, learns that the traumatic event occurred to a close family member or close friend, or experiences first-hand repeated or extreme exposure to aversive details of the traumatic event; 2) The disturbance, regardless of its trigger, causes clinically significant distress or impairment in the individual’s social interactions, capacity to work or other important areas of functioning. It is not the physiological result of another medical condition, medication,

drugs or alcohol; and 3) The experience includes the following diagnostic clusters: a) re-experiencing spontaneous memories of the traumatic event, recurrent dreams related to it, flashbacks or other intense or prolonged psychological distress, b) avoidance of distressing memories, thoughts, feelings or external reminders of the event, c) negative cognitions and mood including myriad feelings, from a persistent and distorted sense of blame of self or others, to estrangement from others or markedly diminished interest in activities, to an inability to remember key aspects of the event, and d) arousal including aggressive, reckless or self-destructive behavior, sleep disturbances, hyper- vigilance or related problems [24]. Using these criteria participants were divided into three groups: 1) nightmare disorder without a history of major trauma and without symptoms of PTSD [25]; 2) partial criteria for PTSD; and 3) full criteria for PTSD.

For analysis, categories of assessment were developed in the attempt to rate artistic use of nightmares, trauma severity, reported functional impairment, and the incorporation in art of contextual imagery. Artistic use of nightmares was rated from 1 = none to 5 = primary artistic focus. Trauma severity was rated from 1 = no clear recall of trauma through 5 = reported extreme trauma. Functional impairment was rated from 1 = none to 5 = volunteered history of associated suicide attempt or psychiatric hospitalization. Contextual imagery was rated from 1 = minimal through 5 = intense (the predominate characteristic of the reported nightmare). The length of nightmare report was also assessed varying from 1 = image with no reported story through 5 = lengthy report developed fully into a three act story.

Results

While the age of those interviewed varied markedly (18-90) for the age of nightmare onset, the mean age at the time of interview was almost identical (61.3). Ten subjects were visual artists (painters, photographers, and/or videographers) and four were writers. All of those interviewed were not currently taking psychotropic medication or using nightmare-suppressive cognitive or behavioral therapies. Reasons given included cost, drug side effects interfere with clarity and rapidity of thought or diminish their levels of enthusiasm, emotion and energy. Artistic use of nightmares was high for all the age groups.

As based on trauma exposure and symptoms of PTSD, the artists could be divided into three groups: (1) Nightmare disorder: No other symptoms of PTSD and no clear history of a major traumatic event and reporting recurrent nightmares since they were very young (#3) (21%); (2) PTSD meeting full criteria - a history of major trauma, definite and persistent functional impairments, as well as symptoms in all the diagnostic clusters (#7) (50%); and group (3) – PTSD

partial criteria, individuals who after their experience of major trauma, developed and continue to experience some, but not all, of the diagnostic cluster of symptoms associated with PTSD (#4) (29%) (Table 1).

Approximately 80% (11/14) of these successful artists using nightmares in their creative work had experienced major trauma and had at least some symptoms associated with PTSD. The rest (3/14) fit better into the diagnostic category referred to as nightmare disorder: individuals without a clear history of major trauma with frequent nightmares present since childhood that can lead to disrupted sleep and sometimes negative effects on next day mood and behavior. No significant difference in report length or extent of narrative development was evident based on PTSD symptom category.

The majority of the interviewees in this project (8/14 – 57%) described extreme traumatic experiences. These individuals were more likely than others interviewed to clearly have PTSD (6/8 – currently met full diagnostic criteria). This group had clearly experienced great distress as a result of their trauma. Six described suicidal ideation and two volunteered descriptions of suicide attempts during their interviews.

Nine of the interviews were with females and five with males. The mean age of trauma experience and nightmare onset was younger for the women respondents (20.8 years of age) than for the men (31.2 years of age). The mean age of nightmare onset for the women interviewed (#4) who had experienced the most severe documented trauma was only 7.5 years of age (range 3-11) where as for men the mean age of onset was 28.5 (range 18-50). The type of trauma was different as well. Three of the four women experienced the loss of a parent or sibling. Two of the three severely traumatized men experienced their trauma during war-associated combat. For those included in this project that had experienced lesser traumas, the gender based age difference was not nearly as profound: a mean of 27.5 for the women (range 4-60) and 32 for the men (range 6-50).

The content of the trauma-associated nightmares was reported as a replay of the actual event by two of the fourteen interviewees. Each of these individuals had nightmares that included metaphoric memories and associated emotional intrusions, closely parodying the actual experience of trauma. For the other individuals whose nightmares begin after the experience of major trauma, 5/9 or 55% had specific content elements from their experience of trauma that were apparent in their nightmares. Those reporting the most severe trauma were able to describe only minimal nightmare content during their interviews. Among the interviewees, 4/14 or 29% reported an overwhelming Central Image (CI) of a tidal wave as the primary image in their nightmare [26]. Other dream CI's included falling (2), spinning vortex (2), live burial, drowning, significant loss (2), and body alteration (1). For each PTSD grouping (full/partial/none) the intensity of central image, the extent of narrative content (dream length), as well as the severity of experienced trauma and the degree to which nightmares are incorporated into their art are compared in (Table 1).

Among the most successful group of interviewees (the 9/14 who had achieved awarded preeminence in their chosen fields), four meet full diagnostic criteria for PTSD, three meet partial criteria, and two

met no criteria. Among those who are somewhat less successful (#5), three met full, one partial, and one no criteria. The two interviewees whose nightmares were replays of their traumatic experience demonstrated the least capacity for an ability to integrate their experience into their art. Of the 11/14 with subjects with symptoms of PTSD, in 10 of 11 cases their most acclaimed work was accomplished after the experience of trauma.

Discussion

In our previous studies, questionnaire responses indicated an association between reported nightmare frequency and artistic / creative interest. The question arose as to whether some of those successful artists surveyed might have frequent nightmares as a symptom of experienced trauma and PTSD. This proposition, while consistent with Freudian theory, was to some degree counter-intuitive since some individuals with PTSD function poorly in their work and social roles [7]. This study indicates, however, that some creative individuals can successfully function and produce art of highest quality after the experience of significant trauma. In this small series, most, but not all, of this group of successful artists met criteria for the diagnosis of PTSD.

These interviews collaborate the PTSD literature in finding that the greater the severity of the trauma, the greater the likelihood for an individual to develop symptoms of PTSD and the functional impairment associated with that diagnosis (this finding is consistent with the PTSD literature) [7,14,15].

Successful artists with nightmares, including both those with PTSD and those without, were successful at the incorporation of their nightmares into the creation of some of their best work. As to specific nightmare content, CI's were commonly reported with tidal waves as primary content in 4 of 14 nightmares. Among those with the most intense central imagery, there was a tendency to report less narrative. There was the suggestion that individuals who had nightmares that are close replays of the actual traumatic event were less artistically successful.

Those interviewed with PTSD symptoms had been treated at some point with behavioral therapies or medications, prescribed to reduce their PTSD symptoms and improve their waking function. Most had used techniques (extinction therapy, imagery therapy, and/or lucid dream control) designed to suppress their nightmares? The painter Edvard Munch, hospitalized more than once in psychiatric hospitals, illustrates this point: "A German once said to me: 'But you could rid yourselves of your troubles.' To which I replied: 'They are part of me and my art. They are indistinguishable from me, and it would destroy my art. I want to keep those sufferings'" [27]. While none of these artists report that they are currently using any of the medications or the varieties of cognitive – behavioral therapy designed to suppress nightmares, three reported the periodic use of hypnotics, and antidepressants, as well as continued involvement in insight-based psychotherapy.

Many individuals being treated for PTSD discontinue their therapies and medications [28]. Some individuals, particularly those who have experienced less severe levels of trauma, can return to their normal level of functioning pre-trauma, but it is not currently clear as to which form of medication or therapy is most likely to help any

particular individual achieve normal functioning [18]. In some cases the discontinuation of medication and/or therapy may reflect a cure, but such was not apparent among this group. All of the groupings in this study with a history of significant trauma continued to have at least some symptoms of PTSD many years after the precipitating event. Many individuals with diagnosed PTSD will periodically achieve higher levels of function until episodic life-stressors, or new trauma induces a return of significant PTSD symptoms and produces a decline in function and quality of life [14] (Figure 1C). For those interviewed in this study, this pattern was commonly reported. Other PTSD affected individuals will learn to accept a level of functioning and quality of life post-trauma that is different, and less rich in quality of experience than the life they had lived before their trauma (Figure 1D). Others never recover physically and/or mentally from their experience of trauma (Figure 1E).

This series of interviews indicates that for artists an alternative approach exists for the individual confronted with the experience of irreconcilable trauma. That approach of creative immersion builds on the Freudian era's insights into the potential of art for individuals experiencing psychological decompensation [29]. An artist has the capacity to create – to project an artistic rendering outside of themselves. Creation can be transformative, incorporating the experience of trauma and sufferings into art. While these individuals continue to have at least some symptoms of PTSD including frequent and recurrent nightmares, through their art they have the potential for realignment after trauma. They can incorporate and project trauma-based angst and distress into their art. Such art can be of extraordinary quality, at a level even higher than possible before for the experience of trauma (Figure 1A and 1b). Such an improvement in their creative capacity can be considered an increase in their level of function in measurable terms that positively affects their quality of life producing commercial success, accolades and honors that for this group included two Academy Awards. No other approach to trauma, no other approach to the nightmare of PTSD, offers the possibility of improved function beyond levels present prior to trauma.

Conclusion

Although a small sample, it is apparent from this study that a significant proportion of successful artists using nightmares in their work have experienced major trauma, and that post trauma, many of these successful artists have symptoms consistent with the diagnosis of PTSD. Rather than attempting to suppress their symptoms post-trauma, these individuals have chosen to incorporate their nightmares and experience of trauma into their life's work. Some have been very successful in that endeavor, achieving levels of accomplishment post-trauma that are higher than pre-trauma. This suggests that for some creative individuals who have experienced irreconcilable trauma, nightmare incorporation and integration into artistic expression, rather than suppression with drugs and psychological therapies, can be a reasonable and even beneficial approach.

References

1. Van de Castle B. *Our Dreaming Mind*. New York. Ballantine Books. 1994.
2. Pagel JF, Kwiatkowski C, Broyles K. Dream Use in Film Making, *Dreaming*. 1999; 9: 247-296.
3. Pagel JF, Crow D, Sayles J. *Filmed dreams: cinematographic and story line characteristics of the cinematic dreamscapes of John Sayles*. *Dreaming*. 2003; 13: 43-48.
4. Pagel JF, Kwiatkowski CF. Creativity and dreaming: Correlation of reported dream incorporation into awake behavior with level and type of creative interest. *Creativity Research Journal*. 2003; 15: 199-205.
5. Pagel JF, Shocknasse S. Dreaming and insomnia: Polysomnographic correlates of reported dream recall frequency. *Dreaming*. 2007; 17: 140-151.
6. Pagel JF. *The Limits of Dream – A Scientific Exploration of the Mind / Brain Interface*, Academic Press (Elsevier) Oxford. 2008.
7. Jones E. *On the Nightmare*, London, The International Psycho-analytic Library. 1923; 20.
8. Zadra A, Donderi D. Nightmares and bad dreams: Their prevalence and relationship to well-being. *J Abnormal Psychology*. 2000; 109: 273-281.
9. Ross R, Ball W, Sullivan K, Caroff S. Dream disturbance as the hallmark of post-traumatic stress disorder. *Am J Psychiatry*. 1989; 146: 697-707.
10. Singareddy RK, Balon R. Sleep in posttraumatic stress disorder. *Ann Clin Psychiatry*. 2002; 14: 183-190.
11. Santiago P, Ursano R, Gray C, Pynoos RS, Spiegel D, Lewis-Fernandez R, et al. A systematic review of PTSD prevalence and trajectories in DSM-5 defined trauma exposed populations: intentional and non-intentional traumatic events. *PLOS One*. 8; e59236.
12. Bisson J, Roberts N, Andrew M, Cooper R, Lewis C. Psychological therapies for chronic Post-Traumatic Stress Disorder (PTSD) in adults. *Cochrane Database Syst Rev*. 2014; 12: CD003388.
13. Krakow B, J Pagel. *Imagery Therapy*, in *Dreaming and Nightmares*. Sleep Medicine Clinics Philadelphia, Saunders/Elsevier. 2010; 277-288.
14. Chopra M, Zhang H, Pless-Kaiser A, Moyer J, Llorente M, Oslin D, et al. PTSD is a chronic, fluctuating disorder affecting the mental quality of life in older adults. *Am J Geriatr Psychiatry*. 2015; 22: 86-97.
15. Ramsawh HJ, Fullerton CS, Mash HB, Ng TH, Kessler RC, Stein MB, et al. Risk for suicidal behaviors associated with PTSD, depression, and their comorbidity in the U. S. Army. *J Affect Disord*. 2014; 161: 116-122.
16. Pagel JF. Treating nightmares – Sleep Medicine and post-traumatic stress disorder. *J Clin Sleep Med*. 2015; 11: 9-10.
17. Marshall RD, Olfson M, Hellman F. Comorbidity, impairment, and suicidality in subthreshold PTSD. *Am J Psychiatry*. 2001; 58: 1467-1473.
18. Seda G, Sanchez-Ortuno MM, Welsh CH, Halbower AC, Edinger JD. Comparative meta-analysis of prazosin and imagery rehearsal therapy for nightmare frequency, sleep quality and post traumatic stress, *JCSM*. 2015.
19. Freud S. *Beyond the Pleasure Principle*, The Standard Edition. (Trans. & Ed.) J. Strachey, London, Hogarth 1916.
20. Freud S. *Creative Writers and Daydreaming*, in *Introductory Lectures on Psychoanalysis, the Standard Edition (Lecture XXIII – The Paths to the Formation of Symptoms)*. (1966) J. Strachey (Ed. & Trans.). New York. 1908; 375.
21. Hartmann E. *The Nightmare*, New York. Basic Books. 1984.
22. Schecter N, Schmeidler G, Staal M. Dream reports and creative tendencies in students in the arts, sciences and engineering. *J Consulting Psychology*. 1965; 29: 415-421.
23. Pagel. *The Creative Nightmare*, Dreamtime Spring. 2015.
24. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*. Arlington, VA, American Psychiatric Association. 2013.
25. Pagel JF, Nielsen T. Parasomnias: Recurrent Nightmares – The International Classification of Sleep Disorders – Diagnostic and Coding Manual (ICD-10). American Academy of Sleep Medicine. 2005.
26. Hartmann E. *The Nature and Functions of Dreaming*, Oxford UK, Oxford University Press. 2011; 7.

27. Svang R. Edvard Munch: The Man and his Art (trans. G. Culverwell) New York, Abbeville Press. 1979; 107.
28. Stein D. Ipser J. Seedal S. Pharmacotherapy for Post Traumatic Stress Disorder (PTSD). Cochrane Database Syst Rev. CD002795.
29. Pagel J. Dream Science: Exploring the Forms of Consciousness, Oxford UK, Academic Press. 2014.