

Mini Review

A Short Review on Violence against Women and Co-Occurring Mental Health Problems

Nur N*

Department of Public Health, University of Cumhuriyet, Turkey

***Corresponding author:** Naim Nur, Department of Public Health, University of Cumhuriyet, 58140, Sivas, Turkey**Received:** September 02, 2015; **Accepted:** December 28, 2015; **Published:** January 04, 2016**Abstract**

Violence Against Women (VAW) is a worldwide problem with significant adverse mental health consequences. However, little is known about VAW and co-occurring mental health problems and new studies are required on this subject. Here, a short systematic overview of the current literature (where available) are provided to underline the importance of problems of VAW and co-occurring mental health problems.

Keywords: Violence; Women; Mental health

Introduction

Although in many countries, Violence Against Women (VAW) has long been seen as a private problem in a family [1,2]; in the past few years it has been recognized as both a public-health and a serious human-rights matter [3-6]. The United Nations Declaration on the Elimination of Violence against Women in 1993 defines the violence against women as “any act of gender-based violence that results in, or is likely to result in, physical, sexual or psychological harm or suffering to women, including threats of such acts, coercion or arbitrary deprivation of liberty, whether occurring in public or in private life” [7]. As a result, the United Nations Secretary General released a study related to VAW, in 2006, which highlighted that ‘Violence against women persists in every country in the world as a pervasive violation of human rights and a major impediment to achieving gender equality’[8].

In the past decade increasing attention has been focused on the women’s mental health manifestation caused by VAW. VAW now widely has been known as a worldwide problem with significant adverse mental health consequences. The mental health status of women has been closely linked with healthiness and wellness of her children. As reported in recent available data malnutrition, low birth weight and other developmental problems are strongly associated with an increased risk of children’s health whose mothers suffer from mental disorders [9-12].

In this short review, a simply Pub Med search was done for the terms of “violence against women” and “health” yielded 2.217 titles dating back to the late of 1970s. There were only two publications about the terms in late of 1970s. In titles of that’s publications the term of VAW was appeared as “the social construction of a private event” and “the myth of masochism”. While in 1980s, there were a few publications focused on VAW, over the next two decades with a considerable increase in researches appearing in the 1990s and 2000s.

Globally, VAW is occurring at alarming rates. According to recent estimates nearly one in three women has experienced one form of physical and/or sexual violence against women in their lifetimes within the context of an intimate relationship [13]. Moreover, VAW is not only a substantial health problem by virtue of its direct effects,

such as physical and sexual, but also that it might contribute to the overall burden of disease as a risk factor for women’s mental health problems.

The relationship between violence and mental health problems has been well-documented with anxiety disorder, depression, Post-Traumatic Stress Disorder (PTSD), sleep disturbances as well as suicidal ideation and attempts [6,14,15]. For instance, a recent review article by Beydoun et al., noted that 11% of women who experienced any form of violence had a major depressive disorder or symptoms were reported by 26% of victimized women [16].

Previous studies indicated worse outcomes related to greater frequency and severity of violence [17], while others showed that even low violence levels (e.g. shoving, pushing, hitting etc.) are related with symptoms of depression [18,19]. For example, more than 50% of women who experienced any type of violence suffered some mental health consequences and approximately 75% of the women who experienced severe violence had one or more diagnosed mental health disorders [20].

In a recent survey argued that the associations between physical, psychological, or sexual violence and poor mental health may be mediated by the continuous stress of live conditions [21]. It has been suggested that there are several factors for the presence of VAW which can be divided as individual factors, personal relationship and factors associated with community norms. It is obviously shown that these factors include a low education of the members of the family and low socio-economic status of women, presence of social support, and childhood domestic violence history [22,23]. Furthermore some evident suggests that women with poor mental health situation are more vulnerable to experiences some form of violence due to their impaired perception, cognition, insight and judgment/decision or inability to resolve the potentially threatening situations [24].

The associations between VAW and mental health are complex and to effectively address these matters we need to more understand the relationships. The aim of this short review was is to summarize existing evidence on violence against women and co-occurring mental health problems. Thus, investigation of the recent available systematic reviews and meta-analyses will allow the researchers to better understand the processes of how women’s experiences of

violence affect their mental health and to develop an appropriate intervention way to prevent violence against women.

VAW and Co-Occurring Mental Health Problems

Recent literature worldwide suggested that, women who have experienced violence at some stage in their lives are two to three times more likely to develop depression and almost four times more likely to have suicidal attempt than women who were not abused [16,25].

Literature researches have shown that psychological, physical or sexual way, the VAW has been associated with greater prevalence of various mental disorders such as anxiety disorders, substance abuse and health disorders mood [6,26-29]. For instance, as showed by Falloot and Harris, virtually all women with mental health problems have a violence history [30].

A recent review study focused on mental health consequences of VAW by Lagdon et al., [31] found that in 39 of 58 studies among women who experiencing physical and/or psychological violence, the most common mental health problem was depression. Also, they reported that in 33 of these 58 studies PTSD was an important mental health consequence. In the same review study noted that of the 18 studies that examined the connections among violence, substance use and other mental health problems; nine of these indicated co-occurring PTSD and the remainder reported anxiety disorder, sleep disturbance, suicide attempt, and substance abuse.

Systematic reviews of the published literature indicated that while there is a variation in rate of VAW across various study settings; VAW was associated with a range of mental health disorders including depression, anxiety, PTSD and sleep disturbance. The relative significance of VAW in its impact on mental health disorder is shown in a recent systematic review and meta-analysis study by Trevillion et al., [32] who found that there was a greater rate and increased odds of experiencing violence for women with anxiety, depression or PTSD. Moreover in another study by Vos et al., [33] found that 34.7 % of the total VAW burden was attributable to depression. This is in comparison to 27.3 % attributable to anxiety and 10.7 % to suicide [33].

The associations between violence against women and mental health problems are more likely to be complex and bi-directional. For example, Devries and colleagues were conducted systematic review and meta-analysis including longitudinal studies to examine the relationship between violence and mental health problems. In their study they suggested that women who experienced violence were more likely to have depressive symptoms and of suicide attempts. At the same time, they noted that women with existing depressive symptoms are more likely to experience violence later in their life [13,34].

The complexity of the relationship between violence against women and mental health problems presents a challenge to healthcare providers in order to help the women with these problems (including depression, anxiety, PTSD and sleep disturbance). This challenge is a result of many factors such as relatively small number of available longitudinal researches, variability among the type of violence, the severity and frequency of violence experiences, differences in the

mental health assessment and the recency of mental health problems [32-35].

Conclusion

Despite the common and reciprocity of violence and co-occurring mental health problems in women's lives, concrete recommendations in responding to these issues are quite lacking. Further longitudinal research using consistent definitions and measures are needed to clarify the temporal nature of these relationships and to growing recognition among health care providers and researchers of the extent of these co-occurring problems.

VAW and mental health problems are complex and bidirectional issue. Because of the nature of this issue of violence and mental health problems, women who experience violence with mental health problems have complex needs. It is clear that understanding the complex relationship between violence and mental health will enable healthcare providers to respond to those needs appropriately, effectively, and compassionately. To address the learning needs of healthcare providers who work with women who experience violence with mental health problems requires better communication, co-ordination across services and systematic assessment of new education and training initiatives. Also, it is important to consider the specific needs of woman at the health care center and to keep the individual of her when designing individualized treatment plans.

References

- Andersson N, Cockcroft A, Ansari U, Omer K, Ansari NM, Khan A, et al. Barriers to disclosing and reporting violence among women in Pakistan: findings from a national household survey and focus group discussions. *J Interpers Violence*. 2010; 25: 1965-1985.
- Gracia E, Herrero J. Perceived neighborhood social disorder and attitudes towards reporting domestic violence against women. *J Interpers Violence*. 2007; 2: 737-752.
- Jewkes R, Levin J, Penn-Kekana L. Risk factors for domestic violence: findings from a South African cross-sectional study. *Soc Sci Med*. 2002; 55: 1603-1617.
- Kishor S, Johnson K. Reproductive health and domestic violence: are the poorest women uniquely disadvantaged? *Demography*. 2006; 43: 293-307.
- Koenig MA, Stephenson R, Ahmed S, Jejeebhoy SJ, Campbell J. Individual and contextual determinants of domestic violence in North India. *Am J Public Health*. 2006; 96: 132-138.
- Ellsberg M, Jansen HA, Heise L, Watts CH, Garcia-Moreno C; WHO Multi-country Study on Women's Health and Domestic Violence against Women Study Team. Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study. *Lancet*. 2008; 371: 1165-1172.
- United Nations Assembly General Declaration on the Elimination of Violence against Women, 1993.
- The Secretary-General's in-depth study on all forms of violence against women, 2006.
- Harpham T, Huttly S, De Silva MJ, Abramsky T. Maternal mental health and child nutritional status in four developing countries. *J Epidemiol Community Health*. 2005; 59: 1060-1064.
- Patel V, Prince M. Maternal psychological morbidity and low birth weight in India. *Br J Psychiatry*. 2006; 188: 284-285.
- Walker SP, Wachs TD, Gardner JM, Lozoff B, Wasserman GA, Pollitt E, et al. International Child Development Steering Group. Child development: risk factors for adverse outcomes in developing countries. *Lancet*. 2007; 369: 145-157.

12. Nur N. Association between domestic violence and miscarriage: a population-based cross-sectional study among women of childbearing ages, Sivas, Turkey. *Women Health*. 2014; 54: 425-438.
13. Devries KM, Mak JY, García-Moreno C, Petzold M, Child JC, Falder G, et al. Global health. The global prevalence of intimate partner violence against women. 2013; 340: 1527-1528.
14. Ellsberg M, Emmelin M. Intimate partner violence and mental health. *Glob Health Action*. 2014; 7: 25658.
15. Howard LM, Trevillion K, Agnew-Davies R. Domestic violence and mental health. *Int Rev Psychiatry*. 2010; 22: 525-534.
16. Beydoun HA, Beydoun MA, Kaufman JS, Lo B, Zonderman AB. Intimate partner violence against adult women and its association with major depressive disorder, depressive symptoms and postpartum depression: a systematic review and meta-analysis. *Soc Sci Med*. 2012; 75: 959-975.
17. Vogel LC, Marshall LL. PTSD symptoms and partner abuse: low income women at risk. *J Trauma Stress*. 2001; 14: 569-584.
18. Stein MB, Kennedy C. Major depressive and post-traumatic stress disorder comorbidity in female victims of intimate partner violence. *J Affect Disord*. 2001; 66: 133-138.
19. Cascardi M, O'Leary KD. Depressive symptomatology, self-esteem, and self-blame in battered women. *J Fam Violence*. 1992; 7: 249-259.
20. Roberts GL, Williams GM, Lawrence JM, Raphael B. How does domestic violence affect women's mental health? *Women Health*. 1998; 28: 117-129.
21. Howard LM, Trevillion K, Khalifeh H, Woodall A, Agnew-Davies R, Feder G. Domestic violence and severe psychiatric disorders: prevalence and interventions. *Psychol Med*. 2010; 40: 881-893.
22. James L, Brody D, Hamilton Z. Risk factors for domestic violence during pregnancy: a meta-analytic review. *Violence Vict*. 2013; 28: 359-380.
23. Flury M, Nyberg E, Riecher-Rössler A. Domestic violence against women: Definitions, epidemiology, risk factors and consequences. *Swiss Med Wkly*. 2010; 140: w13099.
24. Simonelli A, Pasquali CE, De Palo F. Intimate partner violence and drug-addicted women: from explicative models to gender-oriented treatments. *Eur J Psychotraumatol*. 2014; 5.
25. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. Geneva, Switzerland: World Health Organization.
26. Devries K, C Watts, Yoshihama M, Kiss L, Schraiber LB, Deyessa N, et al. Violence against women is strongly associated with suicide attempts: evidence from the WHO multi-country study on women's health and domestic violence against women. *Soc Sci Med*. 2011; 73: 79-86.
27. Ansara DL, Hindin M. Psychosocial consequences of intimate partner violence for women and men in Canada. *J Interpers Violence*. 2011; 26: 1628-1645.
28. Berrios DC, Grady D. Domestic violence. Risk factors and outcomes. *West J Med*. 1991; 155: 133-135.
29. Howard LM, Trevillion K, Khalifeh H, Woodall A, Agnew-Davies R, Feder G. Domestic violence and severe psychiatric disorders: prevalence and interventions. *Psychol Med*. 2010; 40: 881-893.
30. Fallot R, M Harris. *Integrated Service Teams for Women Survivors with Alcohol and other Drug Problems and Co-Occurring Mental Disorders, in Responding to Physical and Sexual Abuse in Women with Alcohol and Other Drug and Mental Disorders*. Haworth Press: Binghampton, USA. 2004.
31. Lagdon S, Armour C, Stringer M. Adult experience of mental health outcomes as a result of intimate partner violence victimisation: a systematic review. *Eur J Psychotraumatol*. 2014; 5.
32. Trevillion K, Oram S, Feder G, Howard LM. Experiences of domestic violence and mental disorders: a systematic review and meta-analysis. *PLoS One*. 2012; 7: e51740.
33. Vos T, Astbury J, Piers LS, Magnus A, Heenan M, Stanley L, et al. Measuring the impact of intimate partner violence on the health of women in Victoria, Australia. *Bull World Health Organ*. 2006; 84: 739-744.
34. Devries KM, Child JC, Bacchus LJ, Mak J, Falder G, Graham K, et al. Intimate partner violence victimization and alcohol consumption in women: a systematic review and meta-analysis. *Addiction*. 2014; 109: 379-391.
35. Bundock L, Howard LM, Trevillion K, Malcolm E, Feder G, Oram S. Prevalence and risk of experiences of intimate partner violence among people with eating disorders: a systematic review. *J Psychiatr Res*. 2013; 47: 1134-1142.