

## Mini Review

# Tissue Regeneration and Healing by ROS-Mediated NOX2 and Ca<sup>2+</sup> Ion Uptake

Lin C-S<sup>1,2</sup>, Wuputra K<sup>1</sup> and Yokoyama KK<sup>1,3,4\*</sup><sup>1</sup>Graduate Institute of Medicine, Kaohsiung Medical University, Taiwan<sup>2</sup>Department of Biological Sciences, National Sun Yat-sen University, Taiwan<sup>3</sup>Center for Stem Cell Research, Kaohsiung Medical University Taiwan<sup>4</sup>Center of Infectious Diseases and Cancer Research, Kaohsiung Medical University, Taiwan**\*Corresponding author:** Kazunari K Yokoyama, Graduate Institute of Medicine, Kaohsiung Medical University, 100 Shih-Chuan 1st Rd., San-Ming District, Kaohsiung 80708, Taiwan**Received:** May 23, 2018; **Accepted:** June 11, 2018;**Published:** June 18, 2018**Abstract**

Unrestrained generation of Reactive Oxygen Species (ROS) is thought to contribute to tissue damage and remodeling in mammalian tissues and cells. However, the locally specific production of ROS plays a critical role in the wound-healing response and in the repair of tissues. Recently, two articles reported the new and unexpected role of ROS generated by the NADPH oxidase 2 complexes in peripheral neural tissues, and mitochondrial ROS (mtROS) production by membrane injury-induced Ca<sup>2+</sup> uptake, which triggers the activation of the Ras homolog family member A (RhoA) to facilitate actin remodeling in injured tissues. These data suggest caution regarding the use of indiscriminate antioxidation therapy after tissue or cell injury, as it may hinder oxidative regenerating signaling, therefore limiting the healing and regeneration of particular local tissues.

**Keywords:** Axonal regeneration; Mitochondrial redox signaling; NOX2 complex; Reactive oxygen species; Repair injured skeletal muscle cells

**Introduction**

Reactive Oxygen Species (ROS) included superoxide anions (O<sub>2</sub><sup>-</sup>), hydroxyl radical (·OH), and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) are generated by reduced reaction of the oxygen [1]. O<sub>2</sub><sup>-</sup> regarded as the primary ROS, is generated abundantly in cells and is rapidly dismutated to H<sub>2</sub>O<sub>2</sub>, which then be converted to (·OH) (Figure 1). ROS are relatively unstable and exhibit a short half-life. The primary origin of intracellular O<sub>2</sub><sup>-</sup> is four organelles such as (i) mitochondria, (ii) Endoplasmic Reticulum (ER), (iii) peroxisomes and (iv) phagosomes, which display localized generation of ROS [2]. In mitochondria Electron Transport System (ETC), ATP is synthesized coupling electron transfer of O<sub>2</sub> in both complex I and II of the ETC, which generates O<sub>2</sub><sup>-</sup> [1]. In the ER, H<sub>2</sub>O<sub>2</sub> is generated by posttranslational oxidative modification during the process of protein folding [2,3]. In peroxisomes, ROS are generated during the metabolic pathways, such as amino acid metabolism, α- and β-oxidation of long-side chain fatty acids and other reactions [4,5]. Another origin of ROS is the membrane-bound NADPH oxidase NOX family and Dual Oxidase (DUOX). These enzymes include seven family members [6] and localize in the membranes of organelle to generate extracellular ROS (eROS). They convert O<sub>2</sub> to O<sub>2</sub><sup>-</sup> by using NADPH as an electron donor [7]. NOX proteins contain an intracellular flavin domain at carboxyl region, which exhibits the binding sites for NADPH and FAD (flavin adenine dinucleotide), and six transmembrane-α-helices domains, which include four histidine residues as heme-binding at amino-terminal portion. NOX proteins act as carrier to transport electrons across the membrane.


In cells, the balance between production of ROS and rescue by antioxidation is crucial for surviving. The antioxidation cascade is represented as enzymes such as (i) Superoxide Dismutase (SOD), (ii) catalases, (iii) Glutathione Peroxidases (GPxs) and (iv) Peroxiredoxins (PRxs) [3]. SOD family catalyzes the dismutation of reactive O<sub>2</sub><sup>-</sup> anions to H<sub>2</sub>O<sub>2</sub> [6,7]. Antioxidant enzymes included

catalases, GPxs, and PRxs are responsible to convert H<sub>2</sub>O<sub>2</sub> to H<sub>2</sub>O and O<sub>2</sub>. The GPxs reduced H<sub>2</sub>O<sub>2</sub> by oxidized glutathione to maintain the level of reduced glutathione for ROS homeostasis [8]. PRx-family has a redox-sensitive cysteine which is inactivated by oxidation *via* H<sub>2</sub>O<sub>2</sub>. Thioredoxin functions as an electron donor to refuel the level of reduced PRx [9].

**ROS and Redox Signaling**

It is widely believed that ROS are deleterious to the mammalian cells by causing oxidative stress *via* the indiscriminate damaging of nucleic acids, proteins, and lipids. The reactive O<sub>2</sub><sup>-</sup> and ·OH radicals can induce damages irreversibly and produce the cellular dysfunction. However, recent evidence has shown that ROS can also involve in signaling functions, known as “Redox signaling” [10]. For example, proteins that participate in a variety of processes, such as the phosphatases PTEN and PTP1B, MAPK kinases, and redox-dependent transcription factors like FOXO4, can be regulated by H<sub>2</sub>O<sub>2</sub> through reversible cysteine oxidation [2,11]. H<sub>2</sub>O<sub>2</sub> oxidizes the thiol of cysteine residue to produce reactive sulfenic acid (-SOH), which can form intra-and intermolecular disulfide (-S-S-) bonds or cyclic sulfonamide (-S-N) or can undergo hyperoxidation to produce sulfinic (-SO<sub>2</sub>H) or sulfonic (-SO<sub>3</sub>H) acids [12]. The reversible modifications may therefore alter the protein structure, function, or activation process of the target. Together with cysteine thiols, H<sub>2</sub>O<sub>2</sub> can also oxidize several other amino acid residues, such as methionine, lysine, tyrosine, histidine, arginine, and proline [13].

One mechanism aimed at explaining how target proteins are recruited during the oxidation process by H<sub>2</sub>O<sub>2</sub>. The colocalization of ROS and targets are occurred, and then the redox related molecules close to the ROS were also accumulated at the same sites [2]. For instance, NOX colocalizes with targets such as phosphatases or kinases, at the membranes, thereby affecting receptor-typed tyrosine kinase signaling [14]. Another mechanism suggests that H<sub>2</sub>O<sub>2</sub> oxidizes




scavenging enzymes like PRx or GPx, which transfer the oxygen atom to target proteins. This signaling is detected for the  $H_2O_2$ -dependent oxidation of the Apoptosis Signaling Kinase 1 (ASK1), and p38 associated phosphorylation, which is dependent on the formation of an ASK1-PRx1 disulfide intermediate [15]. A third mechanism is that the transient inactivation of scavenger by hyperoxidation or post-translational modification results in the accumulation of  $H_2O_2$ , thus allowing the oxidation reactions of target proteins [16]. Therefore, *via* the localized changes in the redox capacity, cells can regulate the ROS to select each signaling [17]. Moreover, depending on the locus of Cys residues in protein, not all of these Cys residues are susceptible to  $H_2O_2$ -dependent oxidation equally [18]. Another system of ROS control involves in the  $H_2O_2$  transport across membranes *via* aquaporin, which are integral proteins and are involved in the transport of the extracellular  $H_2O_2$  produced by NADPH Oxidases (NOXs) across the membrane, to mediate intracellular signaling cascades as described [19]. Thus,  $H_2O_2$  generates two different signaling networks for (i) regular deleterious functions and (ii) intracellular signaling cascades, for tissue repair.

## ROS-Mediated Regeneration and Repair

Recently, two articles demonstrated the evidence of ROS merits for neurite regeneration and wound closure after tissue damages as the new connection. Hervera et al. [20] studied the regeneration of sensory neurons in the Dorsal Root Ganglion (DRG) that protrudes two axonal branches; the peripheral branches and the sensory Central Nervous System (CNS) branch (Figure 2A). A prior conditioning injury on peripheral branch can stimulate axonal regeneration after CSN damage, but the molecular mechanism remains to be elucidated.

Using ROS-sensing dyes, the authors showed that a regeneration-permissive Sciatic Nerve Crush (SNC) increased ROS production in the distant DRG but not a case of a permissive Dorsal Column Axotomy (DCA). They found neurite outgrowth on DRG from mice *ex vivo* after either a conditioning SNC or a local introduction of  $H_2O_2$  to the nerve, indicating that ROS are critical for DRGs to grow the long axons. Furthermore, *in vivo* experiments by Sciatic Nerve Axotomy (SNA) and the conditioning lesion paradigm (DCA after SNC)



support that redox signaling is needed and is sufficient for peripheral axonal regeneration. Importantly, when  $H_2O_2$  conditioning preceded a spinal cord injury, the sensory motor recovery of the animals was significantly improved [20-21].

After SNC, the authors excluded the involvement of mitochondrial ROS. Instead, injury-induced recruitment of macrophages released exosome vesicles that contained NOX2 were taken up by DRGs. These exosome-borne NOX2 were transported to the soma, resulting in the oxidation and inactivation of Phosphatase and the Tensin homolog (PTEN), thus stimulating PI3K/Akt Kinase cascades and DRG outgrowth. The application of the ROS scavenger NAC (N-acetyl-cysteine) to the sciatic nerve limited DRG outgrowth after spinal cord injury, consistent with the results of direct  $H_2O_2$

injection into the sciatic nerve SNA and conditioning H<sub>2</sub>O<sub>2</sub> treatment induced prominent alterations in gene expression that were related to inflammatory signaling and to the pathways regulated by NOX signaling. After SNC injury, the phosphorylated p47phox and catalytic gp91phox in NOX subunits enriched along the axon and in the DRG. By which importin  $\beta$ 1 and dynein were required for transporting NOX2 to the soma, leading to the commitment of ROS to cysteine oxidation, to inactivate PTEN and to activate PI3K/Akt signaling. In turn, this signal induces neurite outgrowth. Thus, the delivery of NOX2 oxidase from macrophages to injured neurons through exosomes stimulates ROS signaling and axon recovery.

In the second article, Horn et al. [22] reported the interplay between Ca<sup>2+</sup> influx and mitochondria-generated ROS (mtROS) to stimulate actin-mediated closure of damaged membranes that was required for the survival of injured mouse muscle cells and human epithelial cells (Figure 2B). A membrane injury results in a steep influx of Ca<sup>2+</sup> and excess Ca<sup>2+</sup> which may trigger a cell death signaling, is scavenged by lysosomes and mitochondria, with the Sarcoplasmic Reticulum Calcium Adenosine Triphosphatase (SERCA) pumping Ca<sup>2+</sup> into the sarcoplasmic reticulum. The Mitochondrial Calcium Uniporter (MCU-1) mediates Ca<sup>2+</sup> uptake into the mitochondria leading to a local flare-up in mtROS production and activation of RhoA to facilitate actin remodeling for membrane repair. Membrane proximal lysosomes undergo Ca<sup>2+</sup>-triggered exocytosis, which may also function to help exclude excess Ca<sup>2+</sup>. Mitochondrial Ca<sup>2+</sup> uptake provides an injury indicator that is capable of directing either cell survival through activation of mtROS-RhoA axis to enhance cytoskeletal remodeling for wound closure, or cell death cascades through induction of the mitochondrial permeability. Thus, quenching mtROS in mice muscles that had been exercised resulted in serious damage to myofibers and reduced muscle force [23].

## Conclusion

The two articles reviewed here demonstrate that globally quenching ROS generation using antioxidants, which are a popular nutritional supplement, may have detrimental, context-specific effects that must be balanced against their potential merits.

## Acknowledgement

We thank CC Ku and JB Pan for editing of this manuscript. This work was supported by the grants from the Ministry of Science and Technology (MOST 106-2320-B-037-001-MY3, MOST 106-2320-B-037-028; MOST 106-2314-B-037-017), the National Health Research Institutes (NHRI-Ex107-10720SI); and Kaohsiung Medical University (KMU-TP103A04, KMU-TP103G03, KMU-TP104E23, KMU-DT104001, KMU-M106001).

## References

- Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J*. 2009; 417: 1-13.
- Dickinson BC, Chang CJ. Chemistry and biology of reactive oxygen species in signaling or stress responses. *Nat Chem Biol*. 2011; 7: 504-511.
- Reczek CR, Chandel NS. ROS-dependent signal transduction. *Curr Opin Cell Biol*. 2015; 33: 8-13.
- Antonenkov VD, Grunau S, Ohlmeier S, Hiltunen JK. Peroxisomes are oxidative organelles. *Antioxid Redox Signal*. 2010; 13: 525-537.
- Fransen M, Nordgren M, Wang B, Apanasets O. Role of peroxisomes in ROS/RNS-metabolism: implications for human disease. *Biochim Biophys Acta*. 2012; 1822: 1363-1373.
- Ritsick DR, Edens WA, McCoy JW, Lambeth JD. The use of model systems to study biological functions of Nox/Duox enzymes. *Biochem Soc Symp*. 2004; 71: 85-96.
- Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nat Rev Immunol*. 2004; 4: 181-189.
- Brigelius-Flohé R, Maiorino M. Glutathione peroxidases. *Biochim Biophys Acta*. 2013; 1830: 3289-3303.
- Rhee SG, Woo HA, Kil IS, Bae SH. Peroxiredoxin functions as a peroxidase and a regulator and sensor of local peroxides. *J Biol Chem*. 2012; 287: 4403-4410.
- Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Curr Biol*. 2014; 24: R453-462.
- Veal EA, Day AM, Morgan BA. Hydrogen peroxide sensing and signaling. *Mol Cell*. 2007; 26: 1-14.
- Miki H, Funato Y. Regulation of intracellular signaling through cysteine oxidation by reactive oxygen species. *J Biochem*. 2012; 151: 255-261.
- Stadtman ER. Protein oxidation and aging. *Science*. 1992; 257: 1220-1224.
- Ushio-Fukai M. Localizing NADPH oxidase-derived ROS. *Sci STKE*. 2006; 2006: re8.
- Jarvis RM, Hughes SM, Ledgerwood EC. Peroxiredoxin 1 functions as a signal peroxidase to receive, transduce, and transmit peroxide signals in mammalian cells. *Free Radic Biol Med*. 2012; 53: 1522-1530.
- Woo HA, Yim SH, Shin DH, Kang D, Yu DY, Rhee SG. Inactivation of peroxiredoxin I by phosphorylation allows localized H<sub>2</sub>O<sub>2</sub> accumulation for cell signaling. *Cell*. 2010; 140: 517-528.
- Toledano MB, Planson AG, Delaunay-Moisan A. Reining in H<sub>2</sub>O<sub>2</sub> for safe signaling. *Cell*. 2010; 140: 454-456.
- Poole LB, Karplus PA, Claiborne A. Protein sulfenic acids in redox signaling. *Annu Rev Pharmacol Toxicol*. 2004; 44: 325-347.
- Miller EW, Dickinson BC, Chang CJ. Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling. *Proc Natl Acad Sci USA*. 2010; 107: 15681-15686.
- Hervera A, De Virgiliis F, Palmisano I, Zhou L, Tantardini E, Kong G, et al. Reactive oxygen species regulate axonal regeneration through the release of exosomal NADPH oxidase 2 complexes into injured axons. *Nat Cell Biol*. 2018; 20: 307-319.
- Krämer-Albers EM. Exosomes deliver ROS for regeneration. *Nat Cell Biol*. 2018; 20: 225-226.
- Horn A, Van der Meulen JH, Defour A, Hogarth M, Sreetama SC, Reed A, et al. Mitochondrial redox signaling enables repair of injured skeletal muscle cells. *Sci Signal*. 2017; 10.
- Cooper ST. Ca<sup>2+</sup> and mitochondrial ROS: Both hero and villain in membrane repair. *Sci Signal*. 2017; 10.