

Mini Review

The Functions of Chicken Embryo Extract

Kevin Li^{1,2*}¹College of Life Science and Medicine, Northwest University, Xi'an, China²Sharon High School, Sharon, MA, USA

*Corresponding author: Kevin Li, College of Life Science and Medicine, Northwest University, Xi'an 710068, China; Sharon High School, Sharon, MA, 02067, USA

Received: March 18, 2022; Accepted: April 15, 2022;

Published: April 22, 2022

Introduction

Chicken embryo eggs are regarded as a high-quality natural nutrition supplement and sickness cure in China (where they are known as Mao Dan) and other Southeast Asian nations [1]. Pregnant women, the elderly and even cancer patients are advised to consume it [2]. Because of its anti-wrinkle activity and stimulation for cell culture, the embryo extract is also frequently employed in the cosmetics industry as well as scientific research field [3]. The functions of chicken embryo extract are summarized in this publication.

Nutrition Supplement

Chick embryo eggs have long been known to be a rich source of nutrients and to contain a variety of bioactive substances [4]. Proteins, amino acids, carbohydrates, lipids, fatty acids, and vitamins have all been found in them [5]. During egg hatching, the nutrients within the embryo alter [6-8]. Seed germination in plants appears to represent a similar transition in the natural world. Beside the abundant amino acids and vitamins, the chick embryo eggs also contain phospholipids and growth factors [9,10].

Antioxidant Activity

Egg white protein hydrolysate has been shown in several studies to have high antioxidant activity [8]. Certain hydrophobic amino acids and basic amino acids have been shown to improve peptide scavenging activity [11]. More hydroxyl radical scavenging chemicals are created during egg hatching. The study's findings show that chicken embryo hydrolysate has a higher antioxidant content and nutritional value than an unhatched egg. Chick embryo eggs were found to have strong antioxidant activity and to delay aging [12,13]. Separated from the egg white protein hydrolysate, two types of peptides have high radical-scavenging activity [14].

The ability of chick embryo egg hydrolysates (CEEH) to scavenge free radicals was concentration-dependent. CEEH also has a higher antioxidant capacity and nutritional value than an unhatched egg. Furthermore, the antioxidant activity of the 11-day and 14-day CEEH was higher, indicating that the incubation medium-term chick embryo eggs have a higher nutritional value [15]. This study's findings may provide significant insight into the possible use of chick embryo eggs in the functional food business.

The antioxidant activity of peptides in fertilized eggs increased as

the incubation time rose. In addition, the natural antioxidant peptide HLFPGPGKKDPV was isolated and described [16].

Immune Function Enhancement

Chick embryo contains interleukin-4 (IL-4) and interleukin-2 (IL-2) [17,18], which could activate the immune system. On the immunosuppressed mouse, chicken embryo eggs act as a robust immunopotentiator, which could be advantageous for the successful application of chicken embryo eggs as a superior immunostimulatory tonic [19].

In *in vivo* mouse tests, crude chicken embryos homogenate has been indicated to have immunomodulatory effects in recent pharmacological investigations [20]. Spleen (spleen lymphocyte proliferation and IL-2 production) and macrophage function (macrophage phagocytosis and NO production) were improved by chicken embryo eggs. The lipid-soluble fraction was mostly important for spleen function, whereas the water-soluble fraction was primarily responsible for macrophage behavior, both of which were dose-dependent. The function of spleen lymphocytes and peritoneal macrophages is improved by chicken embryo extracts [19].

Scholars confirmed that CEEH (chicken embryo egg hydrolysates) can increase the spleen index, promote mouse lymphocyte proliferation, increase hemolysin activity and macrophage phagocytic capacity, and prolong exhaustive lethal swimming time [21]. In aging rats, the chick embryo and nutrient cocktail improved immunological function, increased antioxidant enzyme activity, and repaired organ damage in a synergistic manner [22].

Anticancer Activity

Animal embryo extract has been shown to have anticancer properties in several investigations. Tumor cells were treated with animal embryonic extracts and then transplanted into an embryo, demonstrating that the embryonic milieu supported tumor cell reprogramming [23,24]. Tumor cells *in vitro* were inhibited and their apoptosis was promoted by extracts from frog oocytes and *Danio rerio* (zebrafish) embryos. The link between the extracts' action and DNA methylation modification in tumor cells has been established.

CEE treatment killed the majority of cultivated melanoma cells, but those that survived produced a subclone of actively proliferating cells that showed stronger resistance and proliferative potential than the original line and exhibited stem cell markers and possible tumor development markers [25]. A unique protein complex derived from chicken liver embryo was proved to decrease the invasiveness of HepG2 cancer cells [26]. CEE may promote the reversion of metastatic phenotypes of osteosarcoma cells [27].

Supplement for Cell Culture

Hayden et al. separated the chicken embryo extract fraction into high and low molecular weight components on Sephadex G-25 as early as 1966. In clones of cartilage and pigmented retina cells,

the medium supplemented with the low molecular weight fraction could support complete differentiation. With the inclusion of a big molecular weight component, growth rates could be boosted [28]. Several factors have been discovered in chick embryos, including stem cell factor (SCF), nerve growth factor (NGF), and epidermal growth factor (EGF) [17,18]. These factors perform a variety of physiological functions, such as maintaining neuron survival and healing tissue damage. As a result, animal embryonic extract is employed as a growth factor cocktail for growing diverse stem cells.

As an addition to the medium in cell growth *in vitro*, chicken embryo extract is thought to be a source of functional and structural proteins [29,30]. It has also been utilized to support the growth and development of neuronal, neuroepithelial, and embryonic stem cells in culture [31,32]. A particular neural crest stem cell medium (NCSCM) is necessary to develop neural crest stem cells (NCSC) *in vitro*, which incorporates the most complicated section of the chick embryo extract, which provides crucial growth factors for the NCSC [33]. Treatment with CEE has been shown to improve the morphology and growth rate of the slower-growing SC-1 cell line [34].

Anti-Aging

Aging is a natural process that causes changes in the biological system as well as tissue and organ degeneration. Several functional foods have been discovered to have powerful anti-aging properties, making them suitable candidates for anti-aging therapy [35].

Jia Ma et al. found that supplementing with dietary chicken embryo plus nutritional mixture improves cognitive deterioration in D-gal caused aging rats [36]. Meanwhile, another study from the same group, which used a bone marrow-derived mesenchymal stem cell model, found that chicken embryos may have anti-aging properties. All of these studies could lead to the development of new anti-aging therapies [22]. The efficacy of different sections of the chicken embryo against aging induction was tested using a D-gal-induced aging mice model. The chicken embryo possesses significant antifatigue properties, which may help to reduce the emergence of age-related disorders [37].

References

- Yang S, et al. Purification and Characterization of an Antioxidant Protein from Fertilized Eggs. *Korean journal for food science of animal Resources*. 2016; 36: 791-798.
- Magat M. Balut: "Fertilized duck eggs and their role in Filipino culture". *Western Folklore*. 2002; 61: 63-96.
- Landecker H. *Culturing life: How cells became technologies*. Cambridge, Mass: Harvard University Press. 2007.
- Duan X, Li M, Ji B, Liu X, Xu X. Effect of fertilization on structural and molecular characteristics of hen egg ovalbumin. *Food Chemistry*. 2017; 221: 1340-1345.
- Lidong L, Jianxing Y, Xiaoli M, Hengsong C. Study on nutrients improvement and efficacy factors of embryonated eggs during different incubation periods. *Food Sci*. 2004; 11: 287-290.
- Farkas K, Ratchford IA, Noble RC, Speake BK. Changes in the size and docosahexaenoic acid content of adipocytes during chick embryo development. *Lipids*. 1996; 31: 313-321.
- Firling CE, Severson AR, Hill TA. Aluminum effects on blood chemistry and long bone development in the chick embryo. *Arch Toxicol*. 1994; 68: 541-547.
- Chen C, Chi YJ. Antioxidant, ACE inhibitory activities and functional properties of egg white protein hydrolysate. *J. Food Biochem*. 2011; 36: 383-394.
- Chehelcheraghi F, Eimani H, Sadraie SH. Improved viability of random pattern skin flaps with the use of bone marrow mesenchymal-derived stem cells and chicken embryo extract. *Iranian Journal of Basic Medical Sciences*. 2015; 18: 764-772.
- Farjah GH, Fazli F. The effect of chick embryo amniotic fluid on sciatic nerve regeneration of rats. *Iranian Journal of Veterinary Research*. 2015; 16: 167-171.
- Sarmadi BH, Ismail A. Antioxidative peptides from food proteins: a review. *Peptides*. 2010; 31: 1949-1956.
- Clara Dombre, Nicolas Guyot, Thierry Moreau, Philippe Monget, Mylene da Silva, Joel Gautron, et al. Egg serpins: The chicken and/or the egg dilemma. *Seminars in cell & developmental biology*. 2017; 62: 120-132.
- Carlson ME, Conboy IM. Loss of stem cell regenerative capacity within aged niches. *Aging Cell*. 2007; 6: 371-382.
- Chen Chen, Yu-Jie Chi, Ming-Yang Zhao, Lei LV. Purification and identification of antioxidant peptides from egg white protein hydrolysate. *Amino Acids*. 2012; 43: 457-466.
- Hao Sun, Ting Ye, Yuntao Wang, Ling Wang, Yijie Chen, Bin Li. Antioxidant activities of chick embryo egg hydrolysates. *Food Science & Nutrition*. 2013; 58-64.
- Xiang Duan, Denis Ocen, Fengfeng Wu, Mei Li, Na Yang, Jin Xu, et al. Purification and characterization of a natural antioxidant peptide from fertilized eggs. *Food Research International*. 2014; 56: 18-24.
- Robak T. Stem cell factor. *Acta haematologica Polonica*. 1994; 25: 205-214.
- Vincent Jonchere, Sophie Rehault-Godbert, Christelle Hennequet-Antier, Vonick Sibut, Larry A. Cogburn, Yves Nys, et al. Gene expression profiling to identify eggshell proteins involved in physical defense of the chicken egg. *BMC genomics*. 2010; 11: 57-76.
- Xi Li, Yujie Su, Jun Sun, Yanjun Yang. Chicken embryo extracts enhance spleen lymphocyte and peritoneal macrophages function. *Journal of Ethnopharmacology*. 2012; 144: 255-260.
- Meiqi Wang, Larry J Guilbert, Jie Li, Yingqi Wu, Peter Pang, Tapan K Pasu, et al. A proprietary extract from North American ginseng (*Panax quinquefolium*) enhances IL-2 and IFN- γ productions in murine spleen cells induced by Con-A. *International Immunopharmacology*. 2004; 4: 311-315.
- Qiong Wu, Hai-Dong Yao, Zi-Wei Zhang, Bo Zhang, Fan-Yu Meng, Shi-Wen Xu, et al. Possible correlation between selenoprotein W and Myogenic regulatory factors in chicken embryonic myoblasts. *Biol Trace Elem Res*. 2012; 150: 166-172.
- Jia Ma, Yanru Guo, Jialei Hu, Yue Pan, Xia Qi, Huaxin Wang, et al. The positive effect of chick embryo and nutrient mixture on bone marrow-derived mesenchymal stem cells from aging rats. *Scientific Reports*. 2018; 8: 7051.
- Diez-Torre A, Andrade R, Eguizabal C, López E, Arluzea J, Silió M, et al. Reprogramming of melanoma cells by embryonic microenvironments. *Int. J. Dev. Biol*. 2009; 53: 1563-1568.
- Hendrix MJ, Sefror EA, Sefror RE, Kasemeier-Kulesa J, Kulesa PM, Postovit LM. Reprogramming metastatic tumour cells with embryonic microenvironments. *Nat. Rev. Cancer*. 2007; 7: 246-255.
- Suraeva NM, Morozova LF, Samoilov AV, Burova OS, Golubeva VA, Baryshnikova MA, et al. Changes in the morphological and immunological characteristics of Mel Ibr melanoma cells in response to chicken embryo extract. *Bull. Exp. Biol. Med*. 2015; 159: 520-523.
- Malwina Sosnowska, Marta Kutwin, Barbara Strojny, Piotr Koczoń, Jarosław Szczepaniak, Jaśmina Bałaban, et al. Graphene oxide nanofilm and chicken embryo extract decrease the invasiveness of HepG2 liver cancer cells. *Cancer Nanotechnology*. 2021; 12: 2-35.
- Xiaodong Mu, Bolat Sultankulov, Riddhima Agarwal, Adel Mahjoub, Trevor Schott, Nicholas Greco, et al. Chick Embryo Extract Demethylates Tumor Suppressor Genes in Osteosarcoma Cells. *Clin Orthop Relat Res*. 2014; 472: 865-873.

28. Hayden G Coon, Robert D Cahd. Differentiation *in vitro*: Effects of Sephadex Fractions of Chick Embryo Extract. *Science*. 1966; 153: 1116-1119.
29. Essid N, Chambard JC, Elgaaied AB. Induction of epithelial-mesenchymal transition (EMT) and Gli1 expression in head and neck squamous cell carcinoma (HNSCC) spheroid cultures. *Bosn J Basic Med Sci*. 2018; 18: 336-346.
30. Chehelcheraghi F, Eimani H, Sadraie SH, Torkaman G, Amini A, Shemshadi H. Improved viability of random pattern skin flaps with the use of bone marrow mesenchymal-derived stem cells and chicken embryo extract. *Iran J Basic Med Sci*. 2015; 18: 764-772.
31. Surayeva NM, Samoilov AV. Production of pharmaceutical proteins using transgenic poultry. *Vestn. Ross. Onkol. Nauch. Tsentr*. 2009; 20: 19-26.
32. Kalyani A, Hobson K, Rao MS. Neuroepithelial stem cells from the embryonic spinal cord: isolation, characterization, and clonal analysis. *Dev. Biol*. 1997; 186: 202-223.
33. Pajtler K, Bohrer A, Maurer J, Schorle H, Schramm A, Eggert A, et al. Production of chick embryo extract for the cultivation of murine neural crest stem cells. *J. Vis Exp*. 2010; 45.
34. SA Christman, BW Kong, MM Landry, DN Foster. Chicken Embryo Extract Mitigates Growth and Morphological Changes in a Spontaneously Immortalized Chicken Embryo Fibroblast Cell Line. *Poultry Science*. 2005; 84: 1423-1431.
35. Jinze Xu, Arnold Y Seo, Darya A Vorobyeva, Christy S Carter, Stephen D Anton, Angela MS Lezza, et al. Beneficial effects of a Q-ter based nutritional mixture on functional performance, mitochondrial function, and oxidative stress in rats. *PLoS One*. 2010; 5: e10572.
36. Jia Ma, Huaxin Wang, Bing Liu, Yujia Shan, Huimin Zhou, Xia Qi, et al. Combination of chick embryo and nutrient mixture prevent D-galactose-induced cognitive deficits, immune impairment and oxidative stress in aging rat model. *Scientific Reports*. 2019; 9: 4092.
37. Hsin-Tai Hong, Tsung-Hsien Hsu, Shao-Wen Hung, Chien-Chao Chiu, Chun-Yun Wu, Chia-Chi Chen, et al. Different Parts of the Chicken Embryo Egg Improve D-Galactose-Induced Aging in a Mice Model. *BioMed Research International*. 2021: 6654683.